

Basic Study

Green tea polyphenols ameliorate non-alcoholic fatty liver disease through upregulating AMPK activation in high fat fed Zucker fatty rats

Yi Tan, Jane Kim, Jing Cheng, Madeleine Ong, Wei-Guo Lao, Xing-Liang Jin, Yi-Guang Lin, Linda Xiao, Xue-Qiong Zhu, Xian-Qin Qu

Yi Tan, Jane Kim, Jing Cheng, Madeleine Ong, Wei-Guo Lao, Xing-Liang Jin, Yi-Guang Lin, Linda Xiao, Xue-Qiong Zhu, Xian-Qin Qu, School of Medical and Molecular Biosciences, University of Technology Sydney, NSW 2007, Australia

Jing Cheng, Xing-Liang Jin, Xue-Qiong Zhu, Xian-Qin Qu, The Second Affiliated Hospital of Wenzhou Medical University, Wenzhou 32527, Zhejiang Province, China

Author contributions: Tan Y, Qu XQ and Zhu XQ substantially contributed to the conception and design of the study; Tan Y, Kim J, Cheng J, Ong M, Lao WG, Jin XL, Lin YG, Xiao L contributed to acquisition, analysis and interpretation of data; all authors drafted the article and made critical revisions related to the intellectual content of the manuscript, and approved the final version of the article to be published.

Institutional review board statement: All procedures in animal study were reviewed and approved by the Animal Care and Ethics Committee of the University of Technology Sydney.

Institutional animal care and use committee statement: All procedures involving animals were reviewed and approved by the University of Technology Sydney Animal Ethics Committee (UTS ACEC 2009-325A), following guidelines issued by the National Health and Medical Research Council of Australia.

Conflict-of-interest statement: To the best of our knowledge, no conflict of interest exists.

Data sharing statement: Technical appendix, statistical code, and dataset available from the corresponding author at Xianqin.qu@uts.edu.au. No additional data are available.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this

work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Xian-Qin Qu, PhD, MD, School of Medical and Molecular Biosciences, University of Technology Sydney, 15 Broadway, NSW 2007, Australia. xianqin.qu@uts.edu.au
Telephone: +61-2-95147852
Fax: +61-2-95147852

Received: January 23, 2017

Peer-review started: January 27, 2017

First decision: March 3, 2017

Revised: March 13, 2017

Accepted: May 4, 2017

Article in press: May 4, 2017

Published online: June 7, 2017

Abstract

AIM

To investigate protective effects and molecular mechanisms of green tea polyphenols (GTP) on non-alcoholic fatty liver disease (NAFLD) in Zucker fatty (ZF) rats.

METHODS

Male ZF rats were fed a high-fat diet (HFD) for 2 wk then treated with GTP (200 mg/kg) or saline (5 mL/kg) for 8 wk, with Zucker lean rat as their control. At the end of experiment, serum and liver tissue were collected for measurement of metabolic parameters, alanine aminotransferase (ALT) and aspartate aminotransferase

(AST), inflammatory cytokines and hepatic triglyceride and liver histology. Immunoblotting was used to detect phosphorylation of AMP-activated protein kinase (AMPK) acetyl-CoA carboxylase (ACC), and sterol regulatory element-binding protein 1c (SREBP1c).

RESULTS

Genetically obese ZF rats on a HFD presented with metabolic features of hepatic pathological changes comparable to human with NAFLD. GTP intervention decreased weight gain (10.1%, $P = 0.052$) and significantly lowered visceral fat (31.0%, $P < 0.01$). Compared with ZF-controls, GTP treatment significantly reduced fasting serum insulin, glucose and lipids levels. Reduction in serum ALT and AST levels (both $P < 0.01$) were observed in GTP-treated ZF rats. GTP treatment also attenuated the elevated TNF α and IL-6 in the circulation. The increased hepatic TG accumulation and cytoplasmic lipid droplet were attenuated by GTP treatment, associated with significantly increased expression of AMPK-Thr172 ($P < 0.05$) and phosphorylated ACC and SREBP1c (both $P < 0.05$), indicating diminished hepatic lipogenesis and triglycerides out flux from liver in GTP treated rats.

CONCLUSION

The protective effects of GTP against HFD-induced NAFLD in genetically obese ZF rats are positively correlated to reduction in hepatic lipogenesis through upregulating the AMPK pathway.

Key words: Non-alcoholic fatty liver disease; Green tea polyphenols; Hepatic lipogenesis; Inflammatory cytokines; AMP-activated protein kinase

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The aim of this study was to examine the protective effects and molecular mechanism of green tea polyphenols (GTP) on non-alcoholic fatty liver disease (NAFLD)-induced by high fat diet (HFD) in genetically obese Zucker fatty (ZF) rats. The data of the study has demonstrated: (1) HFD-ZF rats is an optimal rodent model of NAFLD for testing novel/natural agents for this disease; (2) GTP treatment ameliorated metabolic and histopathological abnormalities in HFD-ZF rats with NAFLD; and (3) GTP exerted a protective effect against hepatic steatosis and liver injury by attenuating inflammatory cytokines and inhibiting lipogenesis through upregulating AMPK activation. Therefore, GTP proves to be an effective natural agent for preventing NAFLD and reducing the risk of its severe complications such as nonalcoholic steatohepatitis, cirrhosis and hepatic cellular carcinoma.

Tan Y, Kim J, Cheng J, Ong M, Lao WG, Jin XL, Lin YG, Xiao L, Zhu XQ, Qu XQ. Green tea polyphenols ameliorate non-alcoholic fatty liver disease through upregulating AMPK activation in high fat fed Zucker fatty rats. *World J Gastroenterol*

2017; 23(21): 3805-3814 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v23/i21/3805.htm> DOI: <http://dx.doi.org/10.3748/wjg.v23.i21.3805>

INTRODUCTION

Non-alcoholic fatty liver disease (NAFLD), characterised by excess accumulation of fat in the liver, has a dramatically rising incidence parallel with the increasing population of metabolic syndrome worldwide^[1,2]. NAFLD is emerging as a potential health burden in society because hepatic lipid accumulation can trigger hepatocyte damage, inflammation and fibrogenesis leading to more serious liver disorders such as nonalcoholic steatohepatitis (NASH), cirrhosis and hepatic cellular carcinoma (HCC)^[3]. In addition, increased liver lipids [triglyceride (TG), diglyceride, and ceramides] exacerbate hepatic insulin resistance subsequently leading to type 2 diabetes and cardiovascular complications^[4,5]. Therefore, identifying strategies to reduce excessive fat accumulation in the liver, especially to decrease hepatic TG, is critical for treatment of NAFLD and prevention of its potential consequences such as cardiometabolic disease.

Currently, no pharmacological therapy is approved for NAFLD. Several drug treatments for the management of NAFLD have been proposed, but none have shown considerable efficacy on the whole spectrum of liver damage^[6]. Lifestyle interventions involving exercise and weight loss are the only accepted treatments for this disease but are often difficult to maintain for NAFLD sufferers. Thus, there is an urgent need to discover agents targeted at elevated hepatic lipids and safe for long-term administration.

Green tea (*Camellia sinensis*) contains natural catechins that is a heterogeneous class of polyphenolic compounds, mainly (-)-epigallocatechin-3-gallate (EGCG), with beneficial effects related to their antioxidant and anti-inflammatory properties^[7]. Recently, green tea polyphenols (GTP) has received considerable attention for its metabolic effects against metabolic syndrome and type 2 diabetes^[8]. Several lines of studies suggest that the anti-diabetic effects of GTP are probably due to suppressing appetite, modifying dietary fat emulsification in the gastrointestinal tract, inhibiting gastrointestinal lipolysis and reducing nutrient absorption^[9-11]. Our previous studies have demonstrated that GTP has direct effects on glucose and lipid metabolism through enhancing glycogen synthesis and inhibiting lipogenesis in hepatocytes^[12]. Based on the results of this *in vitro* study, we hypothesised that GTP can reduce hepatic lipid accumulation against fatty liver *in vivo*.

NAFLD is a complex metabolic disease. Many epidemiological studies have demonstrated that overnutrition or inappropriate diet leads to insulin resistance, dyslipidaemia and chronic inflammation,

which significantly contribute to the development of NAFLD^[13,14], in particular for those who have tendency for obesity genetically^[15]. To evaluate effects of GTP on a rodent model that mimics most pathophysiological features of NAFLD in humans, Zucker fatty (ZF) rat, a genetic model of metabolic syndrome with obesity^[16] was used in the study and fed with a high fat diet (HFD). There is increasing evidence that increased liver TG content is highly correlated to increased *de novo* lipogenesis^[17]. AMP-activated protein kinase (AMPK) activity plays a vital role in mediating hepatic lipogenesis^[18]. Recent studies have shown that AMPK activity is reduced by the factors implicated in the development of NAFLD, such as inflammation and obesity^[19,20]. Therefore, inhibition of hepatic lipogenesis by AMPK activation is envisaged as a viable therapeutic strategy to prevent the initiation and progression of NAFLD^[21]. Whether GTP improves both genetic and dietary factors implicated in NAFLD through activation of AMPK pathway, however, has not previously been addressed. In the current study, we tested the hypothesis that GTP would confer an effective therapy against obese- and HFD-induced NAFLD in rats through upregulating the AMPK pathway. Thus, protein molecules involved in AMPK-mediated hepatic lipogenesis were detected followed by a metabolic study in HFD-ZF rats with or without GTP treatment.

MATERIALS AND METHODS

LC-MS analysis of polyphenols in green tea

The polyphenols extracted from green tea (98% purity) were analyzed using an Agilent 1200 series liquid chromatograph/mass selective detector equipped with QTOF 6510 mass spectrometer (Agilent Technologies). Mass spectrometry data were acquired and quantified with an EGCG standard. The concentration of other polyphenols were calculated using the EGCG standard (CAS No: 989-51-5, MF: C₂₂H₁₈O₁₁, MW: 458.378 and Purity: ≥ 99% (HPLC)).

Animal and treatment

Male Zucker fatty (fa/fa) rats (ZF) and their lean littermates, aged 5-7 wk and approximately 160 g body weight, were supplied by the Monash Animal Research Platform (Monash University, Vic, Australia). The animal protocol designed to minimize pain or discomfort to the animals was approved by the Animal Care and Ethics Committee (ACEC # 2009-325A) of the University of Technology Sydney and was in accordance with the National Health and Medical Research Council of Australia Guidelines on Animal Experimentation. Rats were acclimatised in communal cages at 22 °C, with a 12 h light-dark cycle (lights on 0700 h) for 1 wk and had *ad libitum* access to a standard chow diet (Gordon's Specialty Stock Feed, Sydney, Australia) and water. ZF rats (*n* = 24) were fed a high-fat diet (HFD) for 2 wk. The energy

percentage composition of the HFD was 59% fat, 20% carbohydrate, and 21% protein, with equal quantities of fibre, vitamins, and minerals to the standard chow diet. The HFD was replenished daily. After two weeks of HFD feeding, ZF rats were divided into two groups. The control group (ZF-Con; *n* = 12) was administered saline (5 mL/kg of body weight) and ZF rats in the GTP group (ZF-GTP; *n* = 12) was administered GTP at 200 mg/kg of body weight daily for 8 wk *via* oral gavage. HFD feeding was continued throughout the 8 wk treatment period. Lean Zucker rats (Lean, *n* = 12) were fed a standard laboratory chow diet throughout the treatment period.

Determination of metabolic parameters, liver enzymes and insulin sensitivity

At the end of treatment, blood samples were collected from the tail of rats after an overnight fast (12 h). Fasting serum total cholesterol (TC), high-density lipoprotein cholesterol (HDL-C), triglycerides (TG) and non-esterified fatty acids (NEFAs) were analysed using enzymatic colorimetric kits obtained from Roche Diagnostic (Mannheim, Germany) and Wako Pure Chemical Industries (Osaka, Japan) respectively. Low-density lipoprotein cholesterol (LDL-C) concentrations were calculated by Friedewald's formula: LDL-C (mmol/L) = TC - HDL-C - TG/2.2. Fasting serum insulin concentration was measured using a RIA kit (Linco Diagnostic Services Inc, St. Charles). Fasting serum glucose, alanine aminotransferase (ALT) and aspartate aminotransferase (AST) were determined by spectrophotometric analysis using commercial kits (Dialab, Vienna, Austria). All experimental assays were carried out according to the manufacturer's instruction.

Tissue collection and measurement of liver triglycerides

At the end of the experiment, animals were anesthetized using inhalant aesthetic gas (isoflurane and nitrous oxide) after 12 h of fasting. Liver and visceral fat (epididymal and peri-renal adipose tissues) were quickly excised, washed by ice-cold PBS then stored at -80 °C for subsequent histological and molecular assays.

Approximately 50 mg of liver tissues was homogenized at 4 °C in RIPA lysis buffer (Sigma-Aldrich, St. Luis, MO, United States). Lipids from the liver homogenate were extracted using chloroform/methanol method (2:1), evaporated, and dissolved in 1 mL ethanol (Sigma-Aldrich, St. Luis, MO, United States). TG concentration was assayed using kits from Roche Diagnostic (Mannheim, Germany) following the manufacturers' instructions.

Liver histological analysis

A small portion of frozen liver tissue was cut and embedded with pre-cooled optimal cutting compound (Torrance, CA, United States) for cryostat sectioning at 6 μm. The sections were mounted on microscope

Table 1 Effects of green tea polyphenols on parameters associated with obesity and metabolism

Paradigm parameter	Lean	ZF-Con	ZF-GTP
Initial body weight (g)	258.6 ± 9.5	369.8 ± 12.8 ^b	370.4 ± 13.7
	344.8 ± 8.4	449.6 ± 17.6 ^b	404.1 ± 10.8
Body weight (g)			
Food intake (g/d/rat)	23.13 ± 0.04	17.01 ± 0.33 ^a	16.52 ± 0.51
Liver weight (g)	11.84 ± 0.41	13.51 ± 0.82 ^a	12.73 ± 0.74
Visceral adipose (g)	1.96 ± 0.57	9.16 ± 0.17 ^a	6.32 ± 0.21 ^d
Glucose (mmol/L)	4.38 ± 0.54	7.54 ± 1.70 ^a	6.60 ± 1.22 ^c
Insulin (μU/L)	0.55 ± 0.04	77.7 ± 14.3 ^c	38.64 ± 11.2 ^c
SHBG (nmol/L)	102.2 ± 6.9	82.6 ± 4.8 ^a	91.2 ± 5.6 ^c
Serum lipids levels			
Total cholesterol (mmol/L)	2.90 ± 0.07	5.42 ± 0.21 ^b	4.04 ± 0.23 ^c
Triglyceride (mmol/L)	0.11 ± 0.01	0.75 ± 0.02 ^c	0.49 ± 0.04 ^d
HDL (mmol/L)	2.64 ± 0.22	1.37 ± 0.21 ^b	2.03 ± 0.07 ^c
LDL (mmol/L)	1.37 ± 0.2	4.28 ± 0.42 ^c	2.46 ± 0.36 ^c
NEFAs (mmol/L)	0.76 ± 0.04	1.29 ± 0.03 ^a	1.02 ± 0.07 ^c

Zucker fatty (ZF) rat were fed high-fat diet for 2 wk then treated with saline (5 mL/kg of body weight), or green tea polyphenols (GTP, 200 mg/kg of body weight) for 8 wk. Metabolic parameters are represented as mean ± SEM from 12 rats each group. ^a*P* < 0.05, ^b*P* < 0.01 and ^c*P* < 0.001 vs lean rats; ^d*P* < 0.05 and ^e*P* < 0.01 vs ZF-Con rats. NEFA: Non-esterified fatty acids; SHBG: Sexual hormone binding globulin.

slides then fixed with 10% formaldehyde solution for 48 h. The samples were then stained with Haematoxylin and Eosin (H&E) or Oil Red O (ORO, Sigma-Aldrich, St. Luis, MO, United States) to investigate architecture of the liver and hepatic lipid droplets. Stained ORO slides were visualized with the Olympus microscope and images were captured with an Olympus digital camera (DP70, Tokyo, Japan) using Image-Pro6.2 software (Media Cybernetics, Inc. MD, United States). Lipid droplets were quantified at least 5 different high-power fields in a blinded way. For each group, liver samples from 6–8 rats were prepared and stained and six fragments from each liver were further analysed. All slides were scanned at an absolute magnification of 400X using Image-Pro6.2 software (Media Cybernetics, Inc. MD, United States) under a light microscopy (Olympus, BX51 microscope, Tokyo, Japan). Morphometric results are presented as area fractions—the percentage of specific counts in relation to total number of counted points.

Measurement of proinflammatory cytokines in circulation

Serum levels of TNF- α , interleukins (IL-6 and IL-10) and interferon gamma (IFN γ) were measured with a standard sandwich enzyme immunoassay method using the Bio-Plex Pro mouse cytokine assay kit (Bio-rad Laboratories, Hercules, CA, United States) for the simultaneous quantification of TNF- α , ILs and IFN γ according to the manufacturer's instructions. Resulting TNF- α , ILs and IFN γ levels were determined using a Bio-Plex MAGPIX array reader, which identified and quantified each specific reaction based on bead colour and fluorescent signal intensity and expressed as

pictogram per millilitre.

Western blotting

One hundred milligram of frozen liver samples was homogenized in 1 mL lysis buffer (Roche Diagnostics, Indianapolis, IN, United States). Then, the homogenates were centrifuged at 3000 *g* for 10 min at 4 °C. The supernatants were collected and the protein concentration of each supernatant was determined by the Bradford method. Total protein (20 μg) was separated to 7.5% SDS-polyacrylamide gel electrophoresis then transferred to 0.45 μmol/L polyvinylidenedifluoride (PVDF) membrane and immunoblotted with primary antibodies against phospho-AMPK α (Thr172), AMPK α , phospho-ACC (Ser79) and ACC (Cell Signaling Technology Inc, MA), SREBP-1c and β -actin (Santa Cruz, CA, United States) at 1:1000 dilution and secondary antibodies (Santa Cruz, CA, United States) at 1:10000 dilution. Blots were then developed with enhanced chemiluminescence (ECL) (Pierce, IL, United States) according to manufacturer's instructions. The protein bands were visualized by ChemiDoc XRS systems (Bio-Rad Laboratories, CA, United States). The density of bands was quantified with Quality One 4.6.1 software (Bio-Rad, CA, United States) and the quantified results from 8 rats of each group were calculated as percentage of control lean rats for statistical analysis.

Statistical analysis

All values are expressed as mean ± SEM. Comparisons across the three groups were performed using one-way ANOVA followed by post-hoc analysis of Tukey's test to determine significant differences between the two groups using Prism version 6 (GraphPad Inc, San Diego, CA, United States). *P* value < 0.05 was considered statistically significant.

RESULTS

Effects of GTP on body weight and metabolic parameters

Average food intake remained consistent and similar between ZF control rats and GTP treatment at approximately 17 g/rat per day for the duration of the study period. The body weight and visceral fat of ZF rat were significantly higher than their lean control, associated with hyperglycaemia and hyperlipidaemia (Table 1). No significant differences in liver weight were noted between ZF control and GTP-ZF rats. GTP treatment prevented HFD-induced weight gain by 10.1% (*P* = 0.052) and significantly decreased visceral fat by 31.0% (*P* < 0.01) and lowered fasting serum glucose and insulin levels (Table 1, ZF-Con vs ZF-GTP group, both *P* < 0.05). GTP treatment also significantly reduced the elevated serum levels of cholesterol, TG and NEFAs by 25.4% (*P* < 0.05), 34.6% (*P* < 0.01), and 20.9% (*P* < 0.05), respectively, when compared




Figure 1 Effect of green tea polyphenols on alanine transaminase (A) and aspartate transaminase (B) in high-fat diet Zucker fatty rats. Serum alanine transaminase (ALT) and aspartate transaminase (AST) concentrations were measured after 8 wk of daily treatment with saline (5 mL/kg; Lean, ZF-Con) or GTP (200 mg/kg; ZF-GTP). Data are means \pm SEM, $n = 12$ rats/group. ^b $P < 0.01$ vs Lean rats and ^d $P < 0.01$ vs ZF-Con rats.

with the ZF-Con rat. HFD feeding caused a 48.1% reduction of serum HDL level (ZF-Con vs Lean group, $P < 0.01$) while HDL level was significantly higher in GTP treated ZF rats than ZF-Con rats (Table 1, $P < 0.05$).

Effects of GTP on hepatic ALT and AST levels in HFD fed ZF rats

When compared with the lean control, serum hepatic enzyme (ALT and AST) levels in HFD-fed ZF rats were increased by 6.2 fold (Figure 1A; 29.20 ± 4.01 U/L vs 189.89 ± 20.69 U/L, $P < 0.001$) and 3 fold (Figure 1B; 54.71 ± 17.88 U/L vs 183.14 ± 13.91 U/L, $P < 0.001$), respectively. GTP treatment significantly attenuated the elevated ALT and AST values (both $P < 0.01$) when compared with control ZF rats (Figure 1).

Effect of GTP on hepatic steatosis and TG content of HFD ZF rats

The macrovesicular steatosis and fat droplets were assessed in H&E and ORO-stained sections by light microscopy and digital image analysis (DAI). More microvesicular steatosis was observed on H&E stained sections of ZF control rats than ZF-GTP rats and no macrovesicular steatosis in the lean control (Figure 2A). Fat filtration and hepatic steatosis were further confirmed with ORO-stained cryosections because ORO staining analysis is much higher sensitivity and specificity for hepatic steatosis. The elevated lipid accumulation was revealed by many lipid droplets in ORO-stained cryosections of ZF control rats compared to the lean control (Figure 2B). DIA results (Figure 2C) showed significant reduction in lipid droplets with GTP treated ZF rats, indicating that GTP greatly prevented lipid infiltrations and hepatic steatosis.

Consistent with ORO-staining results, measurement of TG content in the liver showed that HFD ZF rats presented significantly higher hepatic TG by 4-fold compared to lean rats (13.65 ± 4.51 μ mol/g vs 3.36 ± 0.72 μ mol/g; $P < 0.01$). Results in Figure 2D showed that GTP treatment significantly reduced hepatic TG content in HFD fed ZF rats (9.05 ± 2.87 μ mol/g vs 13.65 ± 4.51 μ mol/g; $P < 0.01$).

Effects of GTP on inflammatory cytokines in circulation

Circulating inflammatory cytokines were assessed by enzyme immunoassay. Figure 3 shows significantly increased TNF- α and IL-6 in HFD-ZF control rats compared with the Lean control ($P < 0.01$ and $P < 0.05$ respectively), whereas, GTP treatment markedly decreased the elevation of TNF- α and IL-6 (Figure 3A and B, both $P < 0.05$). IL-10 and IFN γ levels were no significant difference among three groups (Figure 3C and D).

Effects of GTP on de novo lipogenesis pathway in the liver of ZF rats

AMPK and ACC are key enzymes that regulate *de novo* lipogenesis in the liver and contribute considerably to overall metabolism of lipids^[15-16]. Results in Figure 4 show that expression of phospho-AMPK α (Thr172) was significantly lower in HFD fed ZF rats compared with the lean control rats ($P < 0.05$), and GTP administration significantly enhanced expression of phospho-AMPK α (Thr172) ($P < 0.05$) in HFD ZF rats. A reduction of ACC (Ser79) phosphorylation was observed in the liver of HFD ZF rats, and this was reversed by GTP treatment in a similar manner to that phospho-AMPK α (Thr172) expression.

SREBP-1c, a key transcription factor regulating the gene expression of key enzymes implicated in upregulating *de novo* lipogenesis in the liver^[22] was also detected by immunoblotting. HFD-fed ZF rats presented an approximate 1.75 fold increase in SREBP-1c expression in the liver compared to the lean control (Figure 5B; $P < 0.01$). GTP treatment abolished overexpression of SREBP-1c protein compared to ZF control rats (Figure 5B; $P < 0.01$).

DISCUSSION

Hepatic steatosis is the first stage of NAFLD, which is mostly associated with increased *de novo* lipogenesis and obesity, however, it can progress to a more advanced form of the disease, NASH, hepatic necrosis, fibrosis and cirrhosis and hepatocellular carcinoma.




Figure 2 Reduction of hepatic steatosis and triglyceride content with green tea polyphenols treatment. A: Images were stained with H&E to visualise macrovesicular steatosis in liver sections at $\times 400$ magnification. Compared to the livers in lean control rats, a larger unstained area that consisted of fat vacuoles was seen in ZF control rats; B: Representative ORO staining of liver samples from lean, ZF-Con and ZF-GTP rats. A larger red area that consisted of lipid droplets (B, C: ORO staining; $\times 400$ magnification) and higher triglyceride contents (D) in the livers of ZF-control rats. GTP treatment significantly reversed these changes in the liver. Data are expressed as mean \pm SME from 6-8 rats each group. ^b $P < 0.01$, ^e $P < 0.001$ vs Lean group; ^d $P < 0.01$ vs ZF-Con group. GTP: Green tea polyphenols; ZF: Zucker fatty.

Therefore, it is important to control NAFLD at the initial stage. Here, we report for the first time that GTP can protect against HFD-induced NAFLD in genetically obese rats. HFD-fed Zucker fatty rats recapitulated the key metabolic features seen in humans with NAFLD, including central obesity, hyperlipidemia, and hyperinsulinemia and elevated fasting glucose levels. Hepatic manifestations of NAFLD in the rats were evidenced by the elevated ALT and AST levels, increased lipids droplets and TG content in the liver. The parameters associated with obesity and metabolisms shown in Table 1 indicate that HFD-fed ZF rat is an optimal model of NAFLD to test preventive and therapeutic agents against NAFLD.

The key finding of this study is that GTP, as a natural agent, protected liver damage by HFD feeding and genetic obese, evidenced by improved hepatic enzymes (AST and ALT) levels and significantly

lowered TG content in the liver of HFD-ZF rats. Results attained from histological analysis showed that GTP also markedly reduced lipid droplets in the liver of HFD-ZF rats. Elevated serum ALT and AST levels are the primary abnormality seen in patients with NAFLD and other chronic liver disease in human^[23]. These finding indicate that GTP can protect hepatocytes against HFD- and obesity-induced hepatic steatosis. Traditionally, beneficial effects of GTP on the liver have been attributed to their antioxidant properties^[8]. Accumulating evidence has revealed that NAFLD is strongly related to inflammation. For example, IL-6, as a proinflammatory cytokine, has been demonstrated as a potential mediator leading to NAFLD. TNF α has been implicated in liver fibrosis and advanced stages of NAFLD in human^[24,25]. In this study, HFD fed obese rats presented systemic inflammation by markedly raised circulating TNF α and IL-6 levels. Although no change




Figure 3 Effects of green tea polyphenols on inflammatory cytokines in circulation. TNF- α (A), IL-6 (B), IL-10 (C) and IFN γ (D) levels in serum of lean, ZF-Con and ZF-GTP rats. Data are mean \pm SEM from 12 rats each group. ^a $P < 0.05$, ^b $P < 0.01$ vs lean group; ^c $P < 0.05$ vs ZF-Con group. GTP: Green tea polyphenols; ZF: Zucker fatty.




Figure 4 Effects of green tea polyphenols on *de novo* lipogenesis pathway in the liver of Zucker fatty rats. Proteins extracted from the livers of lean, Zucker fatty (ZF)-Con. and ZF-GTP rats were separated by SDS-PAGE and immunoblotted with anti-p-AMPK- α , anti-AMPK- α , and anti-p-ACC: Anti-ACC. β -Actin was used to confirm equal protein loading. Upper panels (A and C): Immunoblot bands of detected proteins; lower panels (B and D): Quantitative analysis of corresponding proteins. Data are expressed as mean \pm SME ($n = 8$). ^a $P < 0.05$ vs lean, ^c $P < 0.05$ vs ZF-Con. GTP: Green tea polyphenols; ACC: Acetyl-CoA carboxylase.

was found in anti-inflammatory cytokine IL-10, raised TNF α and IL-6 were attenuated by GTP treatment. This finding is consistent with a recent study in which Xiao *et al.*^[26] reported EGCG, a major active compound

in GTP, against HFD-induced liver fibrosis associated with inflammation in rats through deregulation of TGF/SMAD, PI3 K/Akt/FoxO1, and NF- κ B pathways. Taken together, intervention with GTP and EGCG can




Figure 5 Expression of SREBP-1c in the liver of Zucker fatty rats. Immunoblots (A) and quantitative analysis of (B) SREBP-1c in liver tissue of chow fed lean rats and HFD fed Zucker fatty (ZF) rats treated with saline (5 mL/kg of body weight) or with GTP (200 mg/kg of body weight). β -Actin was used to confirm equal protein loading. Data are expressed as mean \pm SEM ($n = 8$). ^b $P < 0.01$ vs lean, ^d $P < 0.01$ vs ZF-Con. GTP: Green tea polyphenols.

effectively prevent and reverse inflammation-mediated liver damage and fibrosis in rats, suggesting that GTP is beneficial in humans with NAFLD-induced by dietary factors and genetic obesity.

In addition to anti-inflammatory and antioxidant action, protective effects of GTP on NAFLD may be related to improved hyperinsulinemia and hyperlipidaemia. In NAFLD, there is excess of (free fatty acids) FFAs/NEFAs delivered to the liver due to increased adipose lipolysis under insulin resistant status, which lead to hepatic steatosis by over conversion of FFAs into TG and cause hepatocytes injury-induced by lipotoxicity^[27]. In this study, serum NEFAs level was significantly higher in HFD-ZF rats than the lean control, whereas, GTP treatment significantly attenuated the elevated NEFAs levels, which in turn, could reduce substrate influx to the liver for hepatic TG synthesis and also may diminish unoxidized fatty acids accumulation, therefore, prevent lipoperoxidative stress-induced hepatic injury^[27,28]. Furthermore, reduction of hepatic lipids in GTP treated rats may be also attributed for improved hyperglycemia. Consequently, glucose influx to the liver was reduced and ultimately decreased *de novo* lipogenesis (DNL) in the liver.

Hepatic lipogenesis is the metabolic pathway by which acetyl-CoA carboxylase (ACC) converts acetyl-CoA derived from glucose to fats, which involves sub-processes of fatty acid synthesis and subsequent triglyceride synthesis utilizing carbohydrate in the liver^[17]. In healthy individuals, DNL contributes < 5% to liver TG content, whereas in individuals with NAFLD, DNL contributes > 26% of liver TG^[17]. The significant increase (approximately five fold) in DNL suggests that this pathway may play an important role in the development of NAFLD. Thus, controlling hepatic DNL highlights a potential therapeutic utility for alleviating NAFLD and preventing its potential consequences

(NASH and HCC). DNL is regulated by complex enzyme cascades/pathways. Among these, AMPK-mediated phosphorylation of target substrates plays a critical role in the regulation of hepatic lipogenesis^[21]. The activity of AMPK is stimulated by insulin through promoting its phosphorylation at Thr172^[29], subsequently, AMPK activation leads to phosphorylation of ACC1 at Ser79 and ACC2 at Ser221, causing a reduction in ACC activity that lowers malonyl-CoA from acetyl-CoA, which leads to decreased hepatic TG synthesis^[30,31].

We have previously reported that GTP inhibits lipogenesis associated with enhanced expression of phosphorylated AMPK α and ACC proteins in hepatocytes^[12]. To elucidate molecular mechanisms by which GTP improves hepatic steatosis *in vivo*, AMPK-ACC pathway was examined by immunoblotting analysis. We found that obese rats exposed to HFD expressed less AMPK Thr172 associated with reduced phosphorylated ACC-Ser79 protein in the liver than the lean control rats. GTP intervention significantly enhanced protein expression of AMPK Thr172 and ACC Ser79 compared to saline-treated ZF rats, indicating that decreased hepatic TG and reversed liver histopathology are likely through diminishing hepatic lipogenesis in GTP-ZF rats. In addition to suppression of *de novo* lipogenesis, increased AMPK activity associated with inactivation of ACC by GTP may prevent hepatic injury by increasing fatty acid oxidation. Several lines of studies have demonstrated that reduction in malonyl-CoA secondary to AMPK α 1-mediated ACC inactivation enhance carnitine palmitoyltransferase I (CPT I) activity, which in turn, increases in mitochondrial fatty acid oxidation by increasing CPT I flux^[29,32]. Thus, by enhancing AMPK activation, reduced hepatic TG in GTP treated rats is due, at least in part, to suppression of hepatic lipogenesis and fatty acid oxidation in the liver through upregulating AMPK activation.

SREBP-1c is a key transcription factor regulating the gene expression of key enzymes implicated in hepatic lipogenesis and triglyceride-rich lipoprotein secretion^[33]. In this study, SREBP1c levels were lower in HFD-ZF rats and GTP enhanced expression SREBP-1c, which is likely to reduce hepatic triglyceride biosynthesis and out flux. This finding indicates that GTP inhibited lipogenesis also through inactivation of SREBP-1c. This finding explained the molecular mechanism of decreased serum cholesterol and TG levels in GTP-treated ZF rats.

Significant reduction of serum ALT and AST levels and hepatic TG content in GTP-treated HFD ZF rats strongly supports that GTP can protect HFD-induced liver damage. Mechanisms are likely to include anti-inflammation and suppression *de novo* lipogenesis through enhancing AMPK activation and promoting fatty-acid oxidation thereby reducing hepatic lipotoxicity. In summary, we provide the first evidence that consistent administration of GTP protect genetically obese rodent against HFD induced hepatic steatosis. This protection was associated

with increased activation of AMPK pathway to control hepatic *de novo* lipogenesis and hepatic TG secretion. Our results encourage further study into the effects and safety for GTP conferring protection against NAFLD in human.

ACKNOWLEDGMENTS

We are grateful to Zuyi Lushen Kangyuan Co (Meitan, Guizhou, China) for supporting this study by providing GTP. The authors also would like to thank Dr Yali Sun's advice for chemical analysis of GTP.

COMMENTS

Background

Non-alcoholic fatty liver disease (NAFLD) has a dramatically rising incidence and is emerging as a potential health burden in society because hepatic lipid accumulation can trigger hepatocyte damage, inflammation and fibrogenesis leading to more serious liver disorders. There is an urgent need to discover agents for NAFLD treatment and prevention. Green tea polyphenols (GTP) has received considerable attention for its metabolic effects against metabolic syndrome and type 2 diabetes. The previous studies have demonstrated that GTP has direct effects on glucose and lipid metabolism *in vitro* study. Thus, we hypothesise that GTP can reduce hepatic lipid accumulation against fatty liver *in vivo*. The authors tested this hypothesis in high fat diet (HFD) fed genetically Zucker obese rat.

Research frontiers

Currently, no pharmacological therapy is approved for NAFLD and there is also a recent surge in interest in naturally derived products for the treatment of metabolic disorders. There have been no previous studies into GTP on animal models of NAFLD from both dietary and genetic factors.

Innovations and breakthroughs

The authors provide the first evidence that consistent administration of GTP protect genetically obese rodent against HFD induced hepatic steatosis. This protection was associated with increased activation of AMP-activated protein kinase pathway to control hepatic *de novo* lipogenesis and hepatic TG secretion.

Applications

These results encourage further study into the effects and safety for GTP conferring protection against NAFLD in human.

Peer-review

In this article, the authors showed beneficially effects of GTP on NAFLD, including improved lipid profiles and drastically reduced visceral fat, improved liver function and reduced TG accumulation in the liver. The overall study is solid and well designed. The results are consistent with the proposed hypothesis.

REFERENCES

- Williams CD, Stengel J, Asike MI, Torres DM, Shaw J, Contreras M, Landt CL, Harrison SA. Prevalence of nonalcoholic fatty liver disease and nonalcoholic steatohepatitis among a largely middle-aged population utilizing ultrasound and liver biopsy: a prospective study. *Gastroenterology* 2011; **140**: 124-131 [PMID: 20858492 DOI: 10.1053/j.gastro.2010.09.038]
- Wong VW, Chu WC, Wong GL, Chan RS, Chim AM, Ong A, Yeung DK, Yiu KK, Chu SH, Woo J, Chan FK, Chan HL. Prevalence of non-alcoholic fatty liver disease and advanced fibrosis in Hong Kong Chinese: a population study using proton-magnetic resonance spectroscopy and transient elastography. *Gut* 2012; **61**: 409-415 [PMID: 21846782 DOI: 10.1136/gutjnl-2011-300342]
- Pais R, Charlotte F, Fedchuk L, Bedossa P, Lebray P, Poynard T, Ratziu V. A systematic review of follow-up biopsies reveals disease progression in patients with non-alcoholic fatty liver. *J Hepatol* 2013; **59**: 550-556 [PMID: 23665288 DOI: 10.1016/j.jhep.2013.04.02]
- Samuel VT, Liu ZX, Qu X, Elder BD, Bilz S, Befroy D, Romanelli AJ, Shulman GI. Mechanism of hepatic insulin resistance in non-alcoholic fatty liver disease. *J Biol Chem* 2004; **279**: 32345-32353 [PMID: 15166226]
- Savage DB, Petersen KF, Shulman GI. Disordered lipid metabolism and the pathogenesis of insulin resistance. *Physiol Rev* 2007; **87**: 507-520 [PMID: 17429039 DOI: 10.1152/physrev.00024.2006]
- Rotman Y, Sanyal AJ. Current and upcoming pharmacotherapy for non-alcoholic fatty liver disease. *Gut* 2017; **66**: 180-190 [PMID: 27646933 DOI: 10.1136/gutjnl-2016-312431]
- Sabu MC, Smitha K, Kuttan R. Anti-diabetic activity of green tea polyphenols and their role in reducing oxidative stress in experimental diabetes. *J Ethnopharmacol* 2002; **83**: 109-116 [PMID: 12413715 DOI: 10.1016/S0378-8741(02)00217-9]
- Zhao B. Antioxidant effects of green tea polyphenols. *Kexue Tongbao* 2003; **48**: 315-319
- Raederstorff DG, Schlachter MF, Elste V, Weber P. Effect of EGCG on lipid absorption and plasma lipid levels in rats. *J Nutr Biochem* 2003; **14**: 326-332 [PMID: 12873714]
- Shishikura Y, Khokhar S, Murray BS. Effects of tea polyphenols on emulsification of olive oil in a small intestine model system. *J Agric Food Chem* 2006; **54**: 1906-1913 [PMID: 16506852 DOI: 10.1021/jf051988p]
- Cha KH, Song DG, Kim SM, Pan CH. Inhibition of gastrointestinal lipolysis by green tea, coffee, and gomchui (*Ligularia fischeri*) tea polyphenols during simulated digestion. *J Agric Food Chem* 2012; **60**: 7152-7157 [PMID: 22730927 DOI: 10.1021/jf301047f]
- Kim JJ, Tan Y, Xiao L, Sun YL, Qu X. Green tea polyphenol epigallocatechin-3-gallate enhance glycogen synthesis and inhibit lipogenesis in hepatocytes. *Biomed Res Int* 2013; **2013**: 920128 [PMID: 24066304 DOI: 10.1155/2013/920128]
- Luo Y, Burrington CM, Graff EC, Zhang J, Judd RL, Suksaranjit P, Kaewpoowat Q, Davenport SK, O'Neill AM, Greene MW. Metabolic phenotype and adipose and liver features in a high-fat Western diet-induced mouse model of obesity-linked NAFLD. *Am J Physiol Endocrinol Metab* 2016; **310**: E418-E439 [PMID: 26670487 DOI: 10.1152/ajpendo.00319.2015]
- Kanwar P, Nelson JE, Yates K, Kleiner DE, Unalp-Arida A, Kowdley KV. Association between metabolic syndrome and liver histology among NAFLD patients without diabetes. *BMJ Open Gastroenterol* 2016; **3**: e000114 [PMID: 27933201 DOI: 10.1136/bmjgast-2016-000114]
- Mollard RC, Sénéchal M, MacIntosh AC, Hay J, Wicklow BA, Wittmeier KD, Sellers EA, Dean HJ, Ryner L, Berard L, McGavock JM. Dietary determinants of hepatic steatosis and visceral adiposity in overweight and obese youth at risk of type 2 diabetes. *Am J Clin Nutr* 2014; **99**: 804-812 [PMID: 24522441 DOI: 10.3945/ajcn.113.079277]
- Okamoto T, Kanemoto N, Ohbuchi Y, Okano M, Fukui H, Sudo T. Characterization of STZ-Induced Type 2 Diabetes in Zucker Fatty Rats. *Exp Anim* 2008; **57**: 335-345 [PMID: 18633156]
- Lambert JE, Ramos-Roman MA, Browning JD, Parks EJ. Increased *de novo* lipogenesis is a distinct characteristic of individuals with nonalcoholic fatty liver disease. *Gastroenterology* 2014; **146**: 726-735 [PMID: 24316260 DOI: 10.1053/j.gastro.2013.11.049]
- Steinberg GR, Kemp BE. AMPK in Health and Disease. *Physiol Rev* 2009; **89**: 1025-1078 [PMID: 19584320 DOI: 10.1152/physrev.00011.2008]
- Hasanvand A, Amini-Khoei H, Hadian MR, Abdollahi A, Tavangar SM, Dehpour AR, Semiei E, Mehr SE. Anti-

- inflammatory effect of AMPK signaling pathway in rat model of diabetic neuropathy. *Inflammopharmacology* 2016; **24**: 207-219 [PMID: 27506528 DOI: 10.1007/s10787-016-0275-2]
- 20 **Xu XJ**, Gauthier MS, Hess DT, Apovian CM, Cacicedo JM, Gokce N, Farb M, Valentine RJ, Ruderman NB. Insulin sensitive and resistant obesity in humans: AMPK activity, oxidative stress, and depot-specific changes in gene expression in adipose tissue. *J Lipid Res* 2012; **53**: 792-801 [PMID: 22323564]
- 21 **Smith BK**, Marcinko K, Desjardins EM, Lally JS, Ford RJ, Steinberg GR. Treatment of nonalcoholic fatty liver disease: role of AMPK. *Am J Physiol Endocrinol Metab* 2016; **311**: E730-E740 [PMID: 27577854 DOI: 10.1152/ajpendo.00225.2016]
- 22 **Ferré P**, Fofelle F. Hepatic steatosis: a role for de novo lipogenesis and the transcription factor SREBP-1c. *Diabetes Obes Metab* 2010; **12** Suppl 2: 83-92 [PMID: 21029304 DOI: 10.1111/j.1463-1326.2010.01275.x]
- 23 **Arora A**, Sharma P. Non-invasive Diagnosis of Fibrosis in Non-alcoholic Fatty Liver Disease. *J Clin Exp Hepatol* 2012; **2**: 145-155 [PMID: 25755423 DOI: 10.1016/S0973-6883(12)60103-0]
- 24 **Kishimoto T**. IL-6: from its discovery to clinical applications. *Int Immunol* 2010; **22**: 347-352 [PMID: 20410258 DOI: 10.1093/intimm/dxq030]
- 25 **Lesmana CR**, Hasan I, Budihusodo U, Gani RA, Krisnuhoni E, Akbar N, Lesmana LA. Diagnostic value of a group of biochemical markers of liver fibrosis in patients with non-alcoholic steatohepatitis. *J Dig Dis* 2009; **10**: 201-206 [PMID: 19659788 DOI: 10.1111/j.1751-2980.2009.00386.x]
- 26 **Xiao J**, Ho CT, Liong EC, Nanji AA, Leung TM, Lau TY, Fung ML, Tipoe GL. Epigallocatechin gallate attenuates fibrosis, oxidative stress, and inflammation in non-alcoholic fatty liver disease rat model through TGF/SMAD, PI3 K/Akt/FoxO1, and NF-kappa B pathways. *Eur J Nutr* 2014; **53**: 187-199 [PMID: 23515587 DOI: 10.1007/s00394-013-0516-8]
- 27 **Guturu P**, Duchini A. Etiopathogenesis of nonalcoholic steatohepatitis: role of obesity, insulin resistance and mechanisms of hepatotoxicity. *Int J Hepatol* 2012; **2012**: 212865 [PMID: 22792473 DOI: 10.1155/2012/212865]
- 28 **Jones JG**. Hepatic glucose and lipid metabolism. *Diabetologia* 2016; **59**: 1098-1103 [PMID: 27048250 DOI: 10.1007/s00125-016-3940-5]
- 29 **Chiu CY**, Chan IL, Yang TH, Liu SH, Chiang MT. Supplementation of chitosan alleviates high-fat diet-enhanced lipogenesis in rats via adenosine monophosphate (AMP)-activated protein kinase activation and inhibition of lipogenesis-associated genes. *J Agric Food Chem* 2015; **63**: 2979-2988 [PMID: 25756465 DOI: 10.1021/acs.jafc.5b00198]
- 30 **Cho YS**, Lee JI, Shin D, Kim HT, Jung HY, Lee TG, Kang LW, Ahn YJ, Cho HS, Heo YS. Molecular mechanism for the regulation of human ACC2 through phosphorylation by AMPK. *Biochem Biophys Res Commun* 2010; **391**: 187-192 [PMID: 19900410 DOI: 10.1016/j.bbrc.2009.11.029]
- 31 **Choi S**, Choi Y, Choi Y, Kim S, Jang J, Park T. Piperine reverses high fat diet-induced hepatic steatosis and insulin resistance in mice. *Food Chem* 2013; **141**: 3627-3635 [PMID: 23993530 DOI: 10.1016/j.foodchem.2013.06.028]
- 32 **McGarry JD**, Takabayashi Y, Foster DW. The role of malonyl-coa in the coordination of fatty acid synthesis and oxidation in isolated rat hepatocytes. *J Biol Chem* 1978; **253**: 8294-8300 [PMID: 711753]
- 33 **Gorgani-Firuzjaee S**, Meshkani R. SH2 domain-containing inositol 5-phosphatase (SHIP2) inhibition ameliorates high glucose-induced de-novo lipogenesis and VLDL production through regulating AMPK/mTOR/SREBP1 pathway and ROS production in HepG2 cells. *Free Radic Biol Med* 2015; **89**: 679-689 [PMID: 26456051 DOI: 10.1016/j.freeradbiomed.2015.10.036]

P- Reviewer: Liagre B, Zhang JD **S- Editor:** Qi Y **L- Editor:** A
E- Editor: Zhang FF

