

Genomic Epidemiology of NDM-1-Encoding Plasmids in Latin American Clinical Isolates Reveals Insights into the Evolution of Multidrug Resistance

Ricaurte Alejandro Marquez-Ortiz^{1,2}, Leanne Haggerty^{2,7}, Narda Olarte³, Carolina Duarte⁴, Ulises Garza-Ramos⁵, Jesus Silva-Sanchez⁵, Betsy E. Castro¹, Eby M. Sim², Mauricio Beltran⁴, María V. Moncada¹, Alberto Valderrama³, Jaime E. Castellanos⁶, Ian G. Charles^{2,8}, Natasha Vanegas^{1,2}, Javier Escobar-Perez^{1,*}, and Nicola K. Petty^{2,*}

¹Bacterial Molecular Genetics Laboratory, Universidad El Bosque, Bogotá, D.C., Colombia

²The iThree Institute, University of Technology Sydney, New South Wales, Australia

³El Tunal Hospital E.S.E, Bogotá, Colombia

⁴Grupo de Microbiología, Instituto Nacional de Salud, Bogotá, Colombia

⁵Instituto Nacional de Salud Pública (INSP), CISEI, Cuernavaca, Morelos, México

⁶Grupo de Patogénesis Infecciosa, Universidad Nacional de Colombia, Bogotá, D.C., Colombia

⁷Present address: European Bioinformatics Institute, Wellcome Trust Genome Campus, Cambridge, United Kingdom

⁸Present address: Institute of Food Research, Norwich Research Park, Norwich, United Kingdom

*Corresponding authors: E-mails: labgenmolecular@unbosque.edu.co; nicola.petty@uts.edu.au.

Accepted: June 21, 2017

Data deposition: The annotated, complete NDM-1-encoding plasmids sequenced in this study are available in the DDBJ/EMBL/GenBank public databases under the GenBank accession numbers KX832927 (p16Pre36-NDM), KX832926 (p16Pre36-2), CP017672 (pRB151-NDM), KX832928 (p06-1619-NDM), and KX832929 (p06-1619-2). The trimmed and filtered MiSeq sequencing reads for all genomes sequenced in this study have been deposited in the Sequence Read Archive under Bioproject accession number PRJNA342046 with sample accession numbers for each strain listed in supplementary data set 1, Supplementary Material online.

Abstract

Bacteria that produce the broad-spectrum Carbapenem antibiotic New Delhi Metallo- β -lactamase (NDM) place a burden on health care systems worldwide, due to the limited treatment options for infections caused by them and the rapid global spread of this antibiotic resistance mechanism. Although it is believed that the associated resistance gene *bla*_{NDM-1} originated in *Acinetobacter* spp., the role of *Enterobacteriaceae* in its dissemination remains unclear. In this study, we used whole genome sequencing to investigate the dissemination dynamics of *bla*_{NDM-1}-positive plasmids in a set of 21 clinical NDM-1-positive isolates from Colombia and Mexico (*Providencia rettgeri*, *Klebsiella pneumoniae*, and *Acinetobacter baumannii*) as well as six representative NDM-1-positive *Escherichia coli* transconjugants. Additionally, the plasmids from three representative *P. rettgeri* isolates were sequenced by PacBio sequencing and finished. Our results demonstrate the presence of previously reported plasmids from *K. pneumoniae* and *A. baumannii* in different genetic backgrounds and geographically distant locations in Colombia. Three new previously unclassified plasmids were also identified in *P. rettgeri* from Colombia and Mexico, plus an interesting genetic link between NDM-1-positive *P. rettgeri* from distant geographic locations (Canada, Mexico, Colombia, and Israel) without any reported epidemiological links was discovered. Finally, we detected a relationship between plasmids present in *P. rettgeri* and plasmids from *A. baumannii* and *K. pneumoniae*. Overall, our findings suggest a Russian doll model for the dissemination of *bla*_{NDM-1} in Latin America, with *P. rettgeri* playing a central role in this process, and reveal new insights into the evolution and dissemination of plasmids carrying such antibiotic resistance genes.

Key words: metallo-beta-lactamase, genomics, antibiotic resistance, *Providencia rettgeri*, mobile genetic elements, bacterial evolution.

Introduction

In the first global report on antimicrobial resistance issued by the World Health Organization very high resistance rates were found in the bacteria that are the main causes of community and health care associated infections, including *Enterobacteriaceae* and *Acinetobacter baumannii* (World Health Organization 2014). For instance, the prevalence of *Enterobacteriaceae* resistant to broad-spectrum β -lactam type antibiotics often used as “last-line-of-defence” agents, such as carbapenems, are persistently increasing across the world (Rhombert and Jones 2009; Prabaker and Weinstein 2011; van Duijn et al. 2011; CDC 2013). Infections caused by carbapenem-resistant bacteria increase health care costs by requiring hospitalization of patients and increase the risk of mortality (Lemos et al. 2014; World Health Organization 2014). It is therefore important to gain greater insight into how resistance spreads, recognizing in doing so that resistance can spread either vertically, through distribution of clones of established “successful” resistant bacterial species, or horizontally, through dispersal of mobile genetic elements (e.g., transposons, plasmids, and prophages) carrying genes for antimicrobial resistance (Woodford et al. 2011). The horizontal transfer of antibiotic resistance genes between bacteria can contribute rapid expansion in the suite of resistance mechanisms present in a bacterial strain.

One mechanism for resistance that can be acquired from the mobile gene pool is the capability for drug modification, an example of which is the group of metallo- β -lactamases. These enzymes have great impact on public health due to their broad substrate range, and are increasing in frequency in clinically important Gram-negative bacteria (Palzkill 2013). A new member of this group of enzymes was identified in 2008 in a patient treated in New Delhi, India (Yong et al. 2009), and named New Delhi Metallo- β -lactamase (NDM). Initially, NDM dissemination was epidemiologically linked to the Indian subcontinent, though the complexity of transmission of this antibiotic resistance determinant became apparent rapidly, due to the presence of NDM-encoding genes (bla_{NDM}) in diverse Gram-negative bacteria, both fermenters (*Enterobacteriaceae*) and nonfermenters (*Acinetobacter baumannii* and *Pseudomonas aeruginosa*) (Johnson and Woodford 2013). Subsequently, NDM-positive strains were isolated in multiple countries on all continents in a great variety of bacterial genera without any epidemiological or molecular links to the strains circulating in the Indian subcontinent (Johnson and Woodford 2013).

The bla_{NDM} gene is usually carried by conjugative plasmids, although the host plasmid characteristics can vary greatly in attributes such as size, incompatibility group, gene content, and organization. As conjugative plasmids are self-transmitting, this underscores the point that these resistance genes can spread independently of clones of the original bacterial host. In *Acinetobacter* spp., although the bla_{NDM} gene has been reported to be located in the chromosome (Espinal

et al. 2011), it mainly resides in a family of plasmids known as pNDM-BJ01-like, named after the first completely sequenced *Acinetobacter* spp. NDM-plasmid, reported in 2012 (Hu et al. 2012). The pNDM-BJ01-like plasmids are highly conserved, with >99% nucleotide identity extending over at least 85% of the 47-kb pNDM-BJ01 length; they do not belong to any reported incompatibility (Inc) group (Hu et al. 2012); they have a Type IV Secretion System (T4SS); and they have a region of replication and transfer genes separated by a variable region containing a Tn125 composite transposon (Hu et al. 2012). It is within Tn125 that bla_{NDM} is located, along with other genes conserved in the order 5'- bla_{NDM} - ble_{MBL} - $trpF$ - tat - dct - $groES$ - $groEL$ - $ISCR21$ - Δpac -3', flanked upstream and downstream by IS*Aba125*, though the Tn125 structure has been found to be truncated in some strains. Among non-*Acinetobacter* bacteria—with the exception of *Enterobacter aerogenes* (Chen et al. 2015) where bla_{NDM-1} was found in a pNDM-BJ01-like plasmid— bla_{NDM} is carried in a great variety of plasmids belonging to diverse Inc groups (FII, FIB, AVC2, HI1A, HI1B, LM, N, N2, X3, R, T as well as unclassified plasmids) (Johnson and Woodford 2013; Khong et al. 2016). Despite the diversity of NDM-encoding plasmids in non-*Acinetobacter*, the immediate genetic context of the bla_{NDM} gene remains the same in all known cases to date, in that it is always found within Tn125 or its remnants (Poirel et al. 2011; Partridge and Iredell 2012; Wailan, Paterson, et al. 2016). However, Tn125 is often surrounded by other transposons (Tn) or insertion sequence (IS) elements, including IS*Kpn14*, IS26, IS5, ISCR1 or Tn3-like elements, which are frequently found in *Enterobacteriaceae* and may be involved in the further dissemination of bla_{NDM} through a combination of transposition and homologous recombination (Toleman et al. 2012; Khong et al. 2016; Wailan, Sidjabat, et al. 2016). Although the bla_{NDM} gene is believed to have originated in an *Acinetobacter* spp. as the result of the fusion of an aminoglycoside resistance gene with a pre-existing metallo- β -lactamase (Toleman et al. 2012) and later transferred to *Enterobacteriaceae*, aside from the Tn125 remnants and the case of the *Enterobacter aerogenes* harboring a pNDM-BJ01-like plasmid, little is known about this transmission from *Acinetobacter* spp. to *Enterobacteriaceae*, nor about how the diverse bla_{NDM} positive plasmids in *Enterobacteriaceae* evolved.

NDM-positive *Enterobacteriaceae*, particularly *Providencia* spp. play an increasingly important role in multidrug resistant infections and dissemination of bla_{NDM} around the world, as evidenced by the rapidly accumulating reports of isolation of this bacteria harboring this gene (Carvalho-Assef et al. 2013; Mataseje et al. 2014; Pollett et al. 2014; Tada et al. 2014; Carmo Junior et al. 2015; Manageiro et al. 2015; Nachimuthu et al. 2015; Wailan, Paterson, et al. 2016). Previously, we reported the first South American bla_{NDM-1} outbreak, which occurred in Colombia in *Klebsiella pneumoniae*, as well as an NDM-1-positive *Providencia rettgeri* outbreak in Mexico, both

of which occurred in 2011–2012, without any link to the Indian subcontinent (Barrios et al. 2013; Escobar Perez et al. 2013). Shortly thereafter, we commenced a surveillance study across Colombia, and found the majority of NDM-positive bacteria isolated were *P. rettgeri*. Here we describe the use of whole genome sequencing (WGS) to investigate the dissemination dynamics of *bla*_{NDM-1}-positive plasmids among *Enterobacteriaceae* and *A. baumannii* clinical isolates from this surveillance study, as well as from the previous outbreaks in Mexico and Colombia. Our results demonstrate interesting genetic links between NDM-1-positive *P. rettgeri* from distant geographic locations, and between their plasmids and those present in *K. pneumoniae* and *Acinetobacter* spp. isolates, providing insights into the central role of *P. rettgeri* in antibiotic resistance dissemination in Latin America.

Materials and Methods

Isolate Collection and Culture Conditions

Twenty-one NDM-1-positive clinical isolates were included in this study (supplementary data set 1, Supplementary Material online, strains used in this study and statistics of assemblies): *P. rettgeri* (14), *K. pneumoniae* (6), and *A. baumannii* (1). Of these, 12 (11 *P. rettgeri* and one *A. baumannii*) were isolated from samples obtained in a surveillance study for carbapenem resistant bacteria that was conducted over a period of 20 months, from September 2012 to April 2014, in three different hospitals in three distant cities in Colombia (Bogota, Cali, and Bucaramanga) (supplementary data set 1, Supplementary Material online). The other nine clinical isolates were obtained from two, previously described clinical outbreaks generated by *bla*_{NDM-1} positive *K. pneumoniae* and *P. rettgeri*, respectively, reported in Colombia and Mexico (Barrios et al. 2013; Escobar Perez et al. 2013). Clinical and epidemiological features of the all *bla*_{NDM-1} positive isolates are listed in supplementary data set 1, Supplementary Material online. *Escherichia coli* transconjugants were obtained from six representative samples using as donor the *bla*_{NDM-1} positive clinical isolate and as recipient the sodium azide-resistant *E. coli* J53 strain. Equal amounts of a four hours Luria–Bertani (LB) (Oxoid Limited) broth culture of both donor and recipient, were mixed and 100 μ l were placed onto a LB agar plate, then conjugation was allowed for 16 h at 37°C. Subsequently, the NDM-1-positive sodium azide-resistant *E. coli* transconjugants were selected using LB agar plates supplemented with ceftazidime (30 μ g/ml) and sodium azide (100 μ g/ml) (Sigma–Aldrich Co. LLC.). The *E. coli* species (*uidA* gene) and the *bla*_{NDM-1} gene were verified in the transconjugants by PCR. Possible donor strain contamination in the transconjugants was ruled out by PCR using specific primers to the genes *khe* for *K. pneumoniae*, *gyrB* for *A. baumannii*, and *dnaA* for *P. rettgeri* (see supplementary table S1, Supplementary Material online). Otherwise indicated, all NDM-1-positive bacteria were routinely grown in brain hearth

infusion agar or broth supplemented with ceftazidime (30 μ g/ml) as a selective pressure for guarantee of plasmid permanence.

Whole Genome Sequencing

Total DNA was extracted from 21 *bla*_{NDM-1} positive clinical isolates and six transconjugants (supplementary data set 1, Supplementary Material online) using the PureLink[®] Genomic DNA mini kit from ThermoFisher. Multiplexed total DNA libraries were prepared using the Nextera XT Library Preparation Kit and 300-bp paired end sequencing was performed on the Illumina MiSeq platform using the MiSeq v3 600-cycle reagent kit. Sequencing reads were trimmed and filtered using cutadapt v1.1.7 (Martin 2011) to remove adapters, and PRINSEQ-lite v0.20.4 (Schmieder and Edwards 2011) to remove any low quality reads with average read quality less than Q20, low quality trailing ends with base quality less than Q20 and short reads <87 bp. Reads were then de novo assembled using SPAdes v3.5.0 (Bankevich et al. 2012) with default settings and the assemblies were improved to high-quality draft genome standard (Chain et al. 2009) by scaffolding using SSPACE v2.0 (Boetzer et al. 2011), gap filling using GapFiller v1.10 (Boetzer and Pirovano 2012) and removal of contigs shorter than 300 bp. Details of the sequencing data, assemblies and accession numbers for each of these genomes are listed in supplementary data set 1, Supplementary Material online. The *bla*_{NDM-1} gene variant was verified by comparing the genome assemblies against the reported sequence (accession NC_015872) using BLASTn (Altschul et al. 1990). Assembly for each strain was searched for matches to any known *bla*_{NDM-1}-positive plasmids, by BLASTn (Altschul et al. 1990) against an extensive database compiled from all fully sequenced *bla*_{NDM-1}-carrying plasmids deposited in the NCBI nucleotide repository (a total of 141 complete plasmids as at 18 May 2017; supplementary data set 2, Supplementary Material online).

Phylogenetic Analysis of *Providencia rettgeri* Clinical Isolates

Since a MLST scheme for the phylogenetic characterization of *P. rettgeri* isolates does not exist, was built a phylogenetic tree based on the core-genome SNPs determined from the assembled contigs of the 14 *P. rettgeri* genomes sequenced in this study, plus the draft genome of *P. rettgeri* Dmel1 (NZ_AJSB000000000.1), the most complete published *P. rettgeri* genome available at the time, as an out-group control. To build the phylogenetic tree, partially assembled genomes were annotated using Prokka v1.11 (Seemann 2014) and an alignment of concatenated core genes (genes present in all genomes with $\geq 90\%$ of nucleotide identity) was created with Roary (Page et al. 2015) using PRANK (Loytynoja 2014). Poorly aligned positions and divergent regions were eliminated using Gblocks (Talavera and Castresana 2007).

Table 1General features of *bla*_{NDM-1}-positive plasmids harboured in the strains included in this study.

Plasmid	Size (bp)	Inc Group	Host	Resistance Gene Profile	GenBank Accession No	Reference
pNDM-BJ01	47,274	Not assigned	<i>Acinetobacter</i> spp.	<i>aph(3')</i> -Vla, <i>bla</i> _{NDM-1}	NC_019268	Hu et al. (2012)
p6234-178kb	178,193	IncA/C2	<i>K. pneumoniae</i>	<i>aph(3')</i> -Vla, <i>aacA29</i> , <i>aadA2</i> , <i>bla</i> _{NDM-1} , <i>bla</i> _{CARB-2} , <i>mph(E)</i> , <i>msr(E)</i> , <i>catB3</i> , <i>cmlA1</i> , <i>sul2</i> , <i>sul1</i>	NZ_CP010391	Rojas et al. (2016)
p16Pre36-NDM	244,116	Not assigned	<i>P. rettgeri</i>	<i>aadA1</i> , <i>aph(3')</i> -Ia, <i>bla</i> _{NDM-1} , <i>sul2</i> , <i>sul1</i> , <i>tet(B)</i> , <i>dfrA1</i>	KX832927	This study
p16Pre36-2	43,191	Not assigned	<i>P. rettgeri</i>	<i>aac(3)-IIa</i> , <i>bla</i> _{TEM-1B}	KX832926	This study
pRB151-NDM	108,417	Not assigned	<i>P. rettgeri</i>	<i>bla</i> _{NDM-1}	CP017672	Marquez-Ortiz et al. (2017)
p06-1619-NDM	54,712	Not assigned	<i>P. rettgeri</i>	<i>aph(3')</i> -Vla, <i>bla</i> _{NDM-1}	KX832928	This study
p06-1619-2	90,666	Not assigned	<i>P. rettgeri</i>	No resistance genes	KX832929	This study
pPrY2001	113,295	Not assigned	<i>P. rettgeri</i>	<i>aph(3')</i> -Vla, <i>armA</i> , <i>aacA4</i> , <i>bla</i> _{NDM-1} , <i>aac(6')</i> IIb-cr, <i>mph(E)</i> , <i>msr(E)</i> , <i>sul1</i>	NC_022589	Mataseje et al. (2014)

Note.—Plasmids sequenced in this study are shown in bold letters. The *bla*_{NDM-1}-positive pPrY2001 plasmid reported previously in a *P. rettgeri* from Canada was also included.

Finally, the phylogenetic tree was created using RAxML version 8.2.9 (Stamatakis 2014) running 1,000 bootstrap replicates under the generalized time reversible model (GTRCAT). Finally, the consensus tree was plotted using Dendroscope (Huson and Scornavacca 2012). Branch lengths are expressed in units of changes/nucleotide position (scale bar).

Complete Plasmid Sequencing

Total DNA was extracted from three representative *P. rettgeri* isolates (16Pre36, RB151, and 06-1619) using the UltraClean[®] Microbial DNA Isolation Kit (Mo Bio Laboratories, Inc.). BluePippin (Sage Science) 20-kb size-selected libraries were prepared, then sequenced using one SMRT cell each on the PacBio RS II platform (Pacific Biosciences) using P6-C4 chemistry. Sequencing reads were processed and de novo assembled using the HGAP 3 program of SMRT Analysis v2.3 (Chin et al. 2013) with default parameters. To check the assemblies, the filtered PacBio subreads were mapped to the genome assemblies using BWA-MEM (<http://bio-bwa.sourceforge.net/bwa.shtml>). The assembly was visually inspected and manually verified using Tablet v1.15.09.01 (Milne et al. 2013). Misassembled terminal repeat overlap sequences, known to be an error of the HGAP assembly of circular molecules (Chin et al. 2013; Hunt et al. 2015), were identified and subsequently trimmed manually. Circularization results were verified using Circlator (Hunt et al. 2015), confirming that the manual assembly correction correlated with the automated method. The complete sequences of five plasmids were confirmed: two plasmids for the strain 16Pre36, one plasmid for strain RB151 and two plasmids for strain 06-1619 (table 1). As the sequence start point of assemblies are arbitrary, the position one of each plasmid was shifted according to the *repA* gene (pRB151-NDM and p16Pre36-2), pPrY2001 (p16Pre36-NDM and p06-1619-2) or pNDM-BJ01 (p06-1619-NDM) to facilitate

comparative genomics. The plasmids were annotated using Prokka v1.11 (Seemann 2014) and manual curation of the automated annotation was facilitated using Artemis (Rutherford et al. 2000). Antibiotic resistance genes were identified using ARIBA (<https://github.com/sanger-pathogens/ariba/wiki>) and insertion sequence (IS) elements and transposons (Tn) were identified using ISfinder (Siguier et al. 2012) and BLASTn (Altschul et al. 1990). Presence of class 1, 2 or 3 integrons was determined in silico using the primers reported by Marquez et al. (2008).

Comparative Genomics

We used mapping of consensus data from the MiSeq libraries to explore our set of samples for the presence (or residues) of Colombian and Mexican *bla*_{NDM-1}-positive sequenced plasmids (table 1) and other related *bla*_{NDM-1}-positive (pPrY2001 and pNDM-BJ01) and *bla*_{NDM-1}-negative (p06-1619-2) plasmids. For use in the mapping consensus, a reference database was generated using the concatenated complete sequence of the plasmids p6234-178kb, p16Pre36-NDM, pRB151-NDM, p06-1619-NDM, p06-1619-2, pPrY2001 and pNDM-BJ01, broken in fragments of 300 bp (x axis). This reference database was mapped with SHRIMP2 (David et al. 2011) and Nsoni (<https://github.com/Victorian-Bioinformatics-Consortium/nsoni>) against the total MiSeq reads from each sample (y axis). The presence of $\geq 90\%$ nucleotide identity when comparing each 300-bp window from the reference plasmids against the consensus generated from MiSeq reads was determined and visualized as black blocks using SeqFindR (<http://github.com/mscook/seqfindr>). It was included as internal control MiSeq simulated reads to the reference plasmids, generated with ART read simulator (Huang et al. 2012). Pairwise plasmid comparisons, verification of SeqFindR results and figures were performed by using BLASTn




FIG. 1.—*bla*_{NDM-1} plasmids circulating among *Acinetobacter baumannii* and *Klebsiella pneumoniae* in Colombia. (A) BLASTn comparison of WGS assemblies of *A. baumannii* 19Aba78 from this study and its *Escherichia coli* transconjugant C19-78, the *Providencia rettgeri* NDM-positive isolates from Mexico (06-1617, 06-1619 and 06-1623) and the *E. coli* transconjugant C06-1623 against the plasmid pNDM-BJ01. Also were included the plasmids p6200-47kb and p6411-9kb previously reported in Colombia. (B) BLASTn comparison of WGS assemblies of *K. pneumoniae* isolates from this study and the *E. coli* transconjugant C16-1 against the plasmid p6234-178kb. Regions Tra1 and Tra2 common to IncA/C2 plasmids are highlighted (Fernandez-Alarcon et al. 2011); the Tra2 region is disrupted by a variable region. Black circles correspond to the reference plasmids p6234-178kb (A) and pNDM-BJ01 (B), included as internal control.

(Altschul et al. 1990), ACT (Carver et al. 2005), Easyfig (Sullivan et al. 2011), and BRIG (Alikhan et al. 2011).

Results

The *bla*_{NDM-1} Gene Is in a Conjugative Element in the Colombian and Mexican Isolates

From a surveillance study in Colombia a total of 12 NDM-positive isolates were collected from 12 patients (one NDM-positive strain per patient). Among these 12 isolates, which were mostly from outpatients with wound or urinary tract infections, 11 were *P. rettgeri* (five from Bucaramanga and six from Bogota) and one was *A. baumannii* isolated in Cali (supplementary data set 1, Supplementary Material online). Additionally, six *K. pneumoniae* and three *P. rettgeri* from previously reported (Barrios et al. 2013; Escobar Perez et al. 2013) clinical outbreaks in a neonatal unit in Bogota (Colombia) and an intensive care unit in Monterrey (Mexico), respectively, were also included in this study. The *K. pneumoniae* and *P. rettgeri* from Bogota were isolated in the same hospital (supplementary data set 1, Supplementary Material online). Analyses of PCR products confirmed the *bla*_{NDM-1} variant was present in all isolates. Thus, in total, 21 NDM-1-positive isolates from Colombia and Mexico were available to investigate in this study.

To explore if the *bla*_{NDM-1} gene in the Latin American isolates could be transferred between bacteria, conjugation experiments were performed using six representative NDM-

1-positive isolates as donor strains: one representing the *K. pneumoniae* from Bogota, one the *A. baumannii* from Cali, one the *P. rettgeri* from Bogota, two representing the *P. rettgeri* from Bucaramanga and one the *P. rettgeri* from Mexico, with the sodium azide-resistant *Escherichia coli* strain J53 used as the recipient strain. *Escherichia coli* NDM-1-positive transconjugants were obtained from all six of the donor strains (supplementary data set 1, Supplementary Material online). These results indicate that the *bla*_{NDM-1} gene is located in a conjugative element in the Colombian and Mexican isolates, and that it can be transferred to other strains allowing dissemination within and between genera. To determine the relationship among the conjugative *bla*_{NDM-1}-positive genetic structures and among the NDM-1-positive strains circulating in Latin America WGS was performed on the set of 21 Latin American NDM-1-positive clinical isolates as well as the six NDM-1-positive *E. coli* transconjugants (supplementary data set 1, Supplementary Material online).

Acinetobacter baumannii 19Aba78 Harbors *bla*_{NDM-1} within a pNDM-BJ01-Like Plasmid

The WGS assembly of *A. baumannii* isolate 19Aba78 from Cali, Colombia, had 100% nucleotide identity over 99% of the pNDM-BJ01 length (fig. 1A and supplementary data set 2, Supplementary Material online), suggesting that *bla*_{NDM-1} is located on a pNDM-BJ01-like plasmid, as has been broadly reported in *Acinetobacter* spp. (Hu et al. 2012;

Sun et al. 2013; Espinal et al. 2015). The 19Aba78 pNDM-BJ01-like plasmid assembled into seven contigs, so the location of all these contigs together on a single plasmid cannot be confirmed from the current assembly. Of note however, *bla*_{NDM-1} is located within Tn125 adjacent to plasmid sequences that are identical to pNDM-BJ01, on a single contig in the 19Aba78 assembly (see supplementary fig. S1A, Supplementary Material online), supporting the presence of *bla*_{NDM-1} on a pNDM-BJ01-like plasmid. Additionally, the *E. coli* transconjugant C19-78 allowed confirmation of *bla*_{NDM-1} located on a pNDM-BJ01-like plasmid (fig. 1A and supplementary data set 2, Supplementary Material online). All contigs that mapped to the genome sequence of the recipient strain, *E. coli* J53 (accession AICK00000000), were removed from the WGS assembly of the transconjugant C19-78, and all remaining contigs were found to map to pNDM-BJ01, confirming the pNDM-BJ01-like plasmid harbored by C19-78 had no insertions or additional sequences (see supplementary fig. S1A, Supplementary Material online).

Two other NDM-1-positive strains of *Acinetobacter* spp., isolated from other Colombian cities (Neiva and Pasto, 520 km away from each other, and 320 and 390 km away from Cali, respectively) were recently reported (Rojas et al. 2016). The *bla*_{NDM-1}-positive plasmids from these strains, p6200-47kb and p6411-9kb were also found to be pNDM-BJ01-like, each with 99% nucleotide identity over 100% of the pNDM-BJ01 length (fig. 1A and supplementary data set 2, Supplementary Material online). In contrast to the similarity between the *bla*_{NDM-1}-positive plasmids found in the three Colombian NDM-1-positive strains of *Acinetobacter* spp., the strains themselves were of different, unrelated sequence type (ST). The Neiva *A. baumannii* (harboring p6200-47kb) was ST322 and the Pasto *A. nosocomialis* (harboring p6411-9kb) was ST464 (Rojas et al. 2016), but the Cali *A. baumannii* isolate 19Aba78 belongs to ST239, as determined by the Pasteur MLST scheme for *A. baumannii* (Diancourt et al. 2010). These observations of the 19Aba78 isolate and the *E. coli* transconjugant C19-78 thus contribute further evidence of dissemination of closely related pNDM-BJ01-like plasmids among unrelated *Acinetobacter* spp. isolates, as has been reported in Asia and Latin America (Hu et al. 2012; Sun et al. 2013; Waterman et al. 2013; Zhang et al. 2013; Wang et al. 2014; Brovedan et al. 2015; Espinal et al. 2015; Feng et al. 2015; Jones et al. 2015; Li et al. 2015; Rojas et al. 2016), and also its capability to transfer to *Enterobacteriaceae*.

The *bla*_{NDM-1}-Positive Plasmids Circulating in *K. pneumoniae* from Colombia Are Closely Related IncA/C2 Plasmids

Analysis of the WGS assemblies for the six *K. pneumoniae* isolates showed they all have sequences with high similarity to p6234-178kb (>99% nucleotide identity over >98% of the p6234-178kb length, fig. 1B and supplementary data

set 2, Supplementary Material online), an IncA/C2 plasmid harboring *bla*_{NDM-1} from a recently reported *K. pneumoniae* isolated in Neiva (Rojas et al. 2016), which is 300 km away from Bogota in Colombia. It therefore seems likely that the *bla*_{NDM-1} gene is located on a closely related conjugative plasmid in all seven of these Colombian *K. pneumoniae* strains. Despite the plasmid similarities, the *K. pneumoniae* strains from Bogota (16Kpn1, 16Kpn2, 16Kpn3, 16Kpn5, 16Kpn7 and 16Kpn10) are all ST1043 (Escobar Perez et al. 2013), and not related to the ST392 of the *K. pneumoniae* isolate from Neiva (Rojas et al. 2016). The sequences that mapped to p6234-178kb, assembled into several contigs for each of the six Bogota *K. pneumoniae* strains (supplementary data set 1, Supplementary Material online), so the order of the contigs and their genomic location could not be determined. However, the WGS assembly of the NDM-1-positive *E. coli* transconjugant C16-1, obtained using *K. pneumoniae* strain 16Kpn1 as the donor, also showed 99% nucleotide identity over 98% of the length of p6234-178kb (fig. 1B and supplementary data set 2, Supplementary Material online), reinforcing the evidence that *bla*_{NDM-1} is likely to be located on such a related conjugative plasmid in the Bogota outbreak. To rule out the possibility that any related sequences were located in the recipient strain, all contigs that mapped to the chromosome of *E. coli* J53 were removed from the genome assembly of the transconjugant C16-1, and all remaining contigs were found to map to p6234-178kb, with no additional sequences relative to p6234-178kb (see supplementary fig. 1B), and only one confirmed difference due to the absence of an IS5075 element. Thus, this data demonstrates that highly related NDM-1-positive IncA/C2 conjugative plasmids are circulating among *K. pneumoniae* with different genetic backgrounds (ST392 and ST1043) in two distant cities in Colombia.

Providencia rettgeri from Colombia and Mexico Harbor *bla*_{NDM-1} in Different Not Reported Plasmids

Comparison of the WGS assemblies of the Colombian *P. rettgeri* isolates against the database of complete *bla*_{NDM-1}-positive plasmid sequences (supplementary data set 2, Supplementary Material online), revealed that there were no matches over the full length of any known plasmid. The most closely related plasmid was the Inc group unclassified pPrY2001, from a Canadian *P. rettgeri* strain (Mataseje et al. 2014), with >99% nucleotide identity over 69–77% of the length of pPrY2001 (supplementary data set 2 and fig. S2, Supplementary Material online). By way of exception, 16Pre47 (isolated in Bogota) and RB152 (isolated in Bucaramanga) had significant nucleotide identity over only 22% and 15% of the length of pPrY2001, respectively (supplementary data set 2, Supplementary Material online). All four NDM-1-positive *E. coli* transconjugants derived from *P. rettgeri* donors, regardless of the relationship of the respective donor strain with the pPrY2001 plasmid, had sequences that

matched to only a small section of pPrY2001 (<15% of the pPrY2001 length covered with >99% nucleotide identity). The main region of identity to pPrY2001 for all four transconjugants, as well as for 16Pre47 and RB152, was limited to the Tn125 remnant (see supplementary fig. S2, Supplementary Material online).

The *P. rettgeri* isolates from Colombia also had significant sequence matches with the IncA/C2 plasmid p6234-178kb (supplementary data set 2, Supplementary Material online), and in general to all the IncA/C2 NDM-positive plasmids, with >99% nucleotide identity over ~50% of the length of IncA/C2 plasmids (supplementary data set 2, Supplementary Material online). However, the key characteristics of IncA/C2 plasmids were not found in the *P. rettgeri* genome assemblies, as neither the *repA* gene, which is highly conserved among IncA/C2 plasmids, nor any marker for known incompatibility groups (except 16Pre46 with an IncN match), could be identified (see supplementary fig. S3, Supplementary Material online). Furthermore, the corresponding representative *E. coli* transconjugant(s) of *P. rettgeri* isolates from Bogota and Bucaramanga differed in their sequence coverage of IncA/C2 plasmids. *Providencia rettgeri* isolates from Bogota had >99% nucleotide identity over ~65% of the length of p6234-178kb (except 16Pre45 with identity to only 28% of the length), as did the corresponding representative *E. coli* transconjugant C16-36, suggesting the *bla*_{NDM-1}-positive plasmids circulating in Bogota in *P. rettgeri* and in *K. pneumoniae* may be related. By contrast, although the *P. rettgeri* isolates from Bucaramanga had >99% nucleotide identity over ~60% of the length of p6234-178kb (except RB152 with only 26% coverage), the corresponding representative *E. coli* NDM-1-positive transconjugants CRB151 and CRB152 mapped just to 9% of p6234-178kb (supplementary data set 2, Supplementary Material online), the 9% associated with just the Tn125 remnant. Thus, the high relatedness to the IncA/C2 plasmids observed in the donor strains but not the transconjugants, suggests that the Bucaramanga *P. rettgeri* isolates probably have the *bla*_{NDM-1} gene located in a plasmid that is unrelated to p6234-178kb, as well as a *bla*_{NDM-1}-negative structure that is related to the *bla*_{NDM-1}-positive plasmid circulating among *P. rettgeri* and *K. pneumoniae* in Bogota.

Interestingly, the Mexican *P. rettgeri* isolates (06-1617, 06-1619 and 06-1623) and the corresponding representative *E. coli* transconjugant (C06-1623) showed a high level of similarity to the *Acinetobacter* spp. pNDM-BJ01-like plasmids (>99% of nucleotide identity over 74–80% of the length of pNDM-BJ01; fig. 1A and supplementary data set 2, Supplementary Material online). This suggests that a pNDM-BJ01-like plasmid harboring *bla*_{NDM-1} is circulating among the *P. rettgeri* from Mexico, but with a truncated Tn125 and downstream deletion, relative to pNDM-BJ01 (fig. 1A). Additionally, the Mexican *P. rettgeri* isolates showed close relatedness to the pPrY2001 plasmid (>99% of nucleotide identity over 64–76% of the length of pPrY2001); however,




Fig. 2.—Phylogenetic tree of *Providencia rettgeri* isolates based on core-genome SNPs. A Maximum Likelihood (ML) tree was built based on the SNPs in the core-genome assemblies of the NDM-1-positive *P. rettgeri* strains reported in this study (red, blue and green) with the *P. rettgeri* Dmelt1 included as an outgroup control. Branch lengths are expressed in units of changes/nucleotide position (scale bar).

the C06-1623 *E. coli* transconjugant did not. It mapped to only 11% of pPrY2001, which corresponded to the Tn125 remnant (see supplementary fig. S2, Supplementary Material online), suggesting that there is another *bla*_{NDM-1}-negative genetic structure in the Mexican *P. rettgeri* that is related to pPrY2001. No known Inc group was identified in any of the genomes for the *P. rettgeri* isolates from Mexico (except 06-1623, with a match to the IncT group), as is also the case for pNDM-BJ01 (see supplementary fig. S3, Supplementary Material online), further supporting the evidence that *bla*_{NDM-1} is located on a pNDM-BJ01-like plasmid in these strains.

Phylogenetic Concordance with the Geographic Origin of *P. rettgeri*

To investigate the genetic relationship in the *P. rettgeri*, the major NDM-1-positive pathogen identified in the Colombian clinical surveillance (supplementary data set 1, Supplementary Material online), we used the WGS assemblies to build a phylogenetic tree based on the core genome SNPs among the 14 NDM-1-positive *P. rettgeri* isolates included in this study, and found that all but one of the isolates clustered according to the city of origin (fig. 2). The exceptional isolate, 16Pre45, clustered together with the strains from Bucaramanga, even though it was isolated in Bogota. Despite the genetic relationship, no epidemiological link with the Bucaramanga region was identified for the patient harboring 16Pre45 (supplementary data set 1, Supplementary Material online). These results suggest that *bla*_{NDM-1} dissemination in *P. rettgeri* in Colombia and Mexico is following a clonal behavior according to the geographic origin.

General Features of the Complete *bla*_{NDM-1}-Positive Plasmids from Latin American *P. rettgeri*

The complete sequences of plasmids from one representative isolate for each of the three *P. rettgeri* clusters (16Pre36 from




Fig. 3.—General description of the *bla*_{NDM-1}-positive plasmids from three representative *Providencia rettgeri* isolates sequenced in this study, including regions encoding genes for transposition and for replication, and virulence and resistance regions. Plasmid p16Pre36-NDM has two *tra* regions, one reported only for a *P. rettgeri* isolate (Tra-Pre) and the other reported in different IncA/C2 plasmids (Tra1). Plasmids pRB151-NDM and p06-1619-NDM have putative *repA* genes belonging to the IncFII family. Insertion sequences or transposons are shown as rectangles containing their respective CDS for transposition and accessory genes. Gray bars between pairs of sequences indicates >90% nucleotide identity in a window of 400 bp. The scale bar indicates sequence length.

Bogota, RB151 from Bucaramanga and 06-1619 from Monterrey), were obtained by PacBio sequencing. The number of plasmids in each strain varied: 16Pre36 had two plasmids (p16Pre36-NDM and p16Pre36-2), RB151 had one plasmid (pRB151-NDM) and 06-1619 had two plasmids (p06-1619-2 and p06-1619-NDM) (table 1). Among these representative *P. rettgeri* isolates, the *bla*_{NDM-1} gene was located on unrelated plasmids (fig. 3). None of the plasmids belonged to any reported incompatibility group using Carattoli et al. (2014) schemes (see supplementary fig. S3, Supplementary Material online), however, annotation of the plasmids pRB151-NDM and p06-1619-NDM, showed they have two and one *repA* genes, respectively (fig. 2), which are closely related to each other (>85% nucleotide identity over the full *repA* sequence). The putative plasmid replication proteins encoded by these *repA* genes each had a significant match to the IncFII RepA protein family in Pfam (Finn et al. 2016). However, these three *repA* genes showed a poor relation (<51% nucleotide identity over the full *repA* sequence) to the reported IncFII *repA* genes (Carattoli et al. 2014), and plasmids pRB151-NDM and p06-1619-NDM are not related to any reported IncFII plasmid (not shown). In p06-1619-NDM the *repA* gene is located inside a putative mobile element, which possibly brought *repA* from another plasmid (fig. 3).

Plasmid p16Pre36-NDM was found to encode two different putative conjugative transfer-associated regions (fig. 3). One (described here as Tra-Pre), has so far only been reported in the *bla*_{NDM-1}-positive plasmid pPrY2001; the other, is a common transfer-associated region of IncA/C2 plasmids described as Tra1 by Fernandez-Alarcon et al. (2011), which is frequently found in widely disseminated plasmids in a broad

host range (Sekizuka et al. 2011; Doublet et al. 2012; Diene et al. 2013; Tijet et al. 2015; Wang et al. 2015; Wasyl et al. 2015; Rojas et al. 2016), including the IncA/C2 p6234-178kb-like *bla*_{NDM-1}-positive plasmids in the Colombian *K. pneumoniae* strains (supplementary data set 1, Supplementary Material online). At 244,116 bp, p16Pre36-NDM is the largest and most variable of the *bla*_{NDM-1}-positive plasmids sequenced in this study, and among the largest *bla*_{NDM-1}-positive plasmids ever reported (supplementary data set 2, Supplementary Material online). It has two resistance regions each containing a toxin–antitoxin system and a class 1 integron. Both class 1 integrons have the genes *dfrA1-aadA1-qacEΔ1-sul1* associated with resistance to quaternary ammonium compounds, aminoglycosides, sulphonamides and trimethoprim. This plasmid also carries the additional resistance genes *aph(3′)-Ia*, *sul2* and *tet(B)*. One of the resistance regions contains a Tn125 remnant (with its *bla*_{NDM-1} gene intact) inside a shuffled Tn21 element, with the two Tn21 inverted repeats flanking the Tn125 remnant (fig. 4). This shuffled Tn21 element has its typical components IS1353, IS1326, the *mer* operon and a class 1 integron (Liebert et al. 1999), but they are rearranged and the *mer* operon is separated from the rest of the Tn21 by the Tn125 remnant (fig. 4). Tn21-like elements are implicated in the global dissemination of antibiotic resistance genes among *Enterobacteriaceae* and *Pseudomonas* (Liebert et al. 1999) and have been reported to generate mosaic structures (Yurieva et al. 1997; Noguchi et al. 2000; Partridge et al. 2001; Valverde et al. 2006). Moreover, the ΔTn125 (having just one copy of the ISAb125 element) surrounding *bla*_{NDM-1} in p16Pre36-NDM has suffered a rearrangement that has not been previously reported: the genes




Fig. 4.—Comparison of variable *bla*_{NDM-1}-containing regions from plasmids p16Pre36-NDM, pRB151-NDM, p06-1619-NDM, p6234-178kb, pPrY2001 and pNDM-BJ01. Insertion sequences or transposons are shown as rectangles containing their respective CDS for transposition and accessory genes in different colors. Outside orange and green triangles correspond to the inverted repeats of a putative Tn3-like (Tn6369) and a Tn21, respectively. The prototype sequence of the Tn21 was included (Liebert et al. 1999). Dashed lines indicate the 2,928-bp sequence containing ISVsa3 carried by p6234.178kb (once) and p16Pre36-NDM (twice). Gray and red (inverted matches) shading between pairs of sequences indicates >90% of nucleotide identity in a window of 400 bp. The scale bar indicates sequence length.

tat-dct-groES-groEL-ISCR21-Δpac that are found upstream of the *bla*_{NDM-1} in p16Pre36-NDM, have always been previously reported downstream (fig. 4). These odd rearrangements of the ΔTn125 and the Tn21 in p16Pre36-NDM were confirmed by mapping of the PacBio reads to the assembled plasmid (no regions of low read coverage or quality were found that could suggest an assembly issue); by investigation of the MiSeq assembly (the same rearrangement was identified on a single contig, see supplementary fig. S4A, Supplementary Material online); and by using PCR to confirm the location of the ΔTn125 inside the Tn21 (see supplementary fig. S4B, Supplementary Material online). The entire ΔTn21 region in p16Pre36-NDM is flanked upstream and downstream by copies of IS26 (fig. 4), thus it is possible that this entire region of DNA may be mobilized as a composite transposon. IS26 intramolecular replicative transposition has been previously identified as the source of reorganization of plasmids carrying multidrug-resistant determinants as could have happened here (He et al. 2015).

Compared with p16Pre36-NDM the pRB151-NDM (from Bucaramanga, Colombia) is a less complex plasmid, with a size of 108,417 bp. This is a novel plasmid backbone, unrelated to any previously reported (without any significant match against the NCBI nucleotide database), that encodes a putative conjugative transfer machinery, plasmid replication and partition proteins, a restriction-modification system and a putative fimbrial operon (fig. 3). It possesses only one resistance and virulence region, and that contains the *bla*_{NDM-1} gene as the plasmid’s only antibiotic resistance determinant (table 1). The variable region has a ΔTn125 harboring the *bla*_{NDM-1} with a typical structure except that the two flanking ISAbal125 are both truncated (fig. 4). Downstream of the ΔTn125 in the resistance region is a novel transposon, registered as Tn6369 in the Tn Number Registry (Roberts et al. 2008), whilst upstream there is another putative mobile element (or its remains) encoding two putative transposon resolvases and a ParED toxin–antitoxin system that is also present in the p06-1619-NDM plasmid, plus an upstream ΔISYal1 (fig. 4).




FIG. 5.—BLASTn comparison of (A) pNDM-BJ01 with the related p06-1619-NDM plasmid from *Providencia rettgeri*, and (B) pPrY-like plasmids (p16Pre36-NDM, pPrY2001 and p06-1619-2) from *P. rettgeri* with the IncA/C2 related p6234-178kb plasmid reported in *Klebsiella pneumoniae*. Conserved pPrY-like region is highlighted in purple rectangles with dashed lines. Gray and red (inverted matches) shading between pairs of sequences indicate >90% of nucleotide identity in a window of 400 bp. The scale bar indicates sequence length.

The p06-1619-NDM plasmid (from Mexico) has a size of 54,712 bp, and belongs to the conserved pNDM-BJ01-like family of plasmids (from *Acinetobacter* spp.), with the same backbone (99% nucleotide identity over 74% of the pNDM-BJ01 length, fig. 5A). However, it has a putative mobile genetic element (MGE) flanked by identical copies of a novel transposon with intact transposition genes, registered as Tn6368, that has inserted downstream of the *bla_{NDM-1}* gene. The insertion of this MGE, has truncated the Tn125 after the *tat* gene upstream (fig. 4), as well as deleted the genes *parB* and *virB5* and truncated *virB4* downstream, so truncating the T4SS locus (fig. 5A). These genes are implicated in the mating pair formation and DNA partitioning process (Christie et al. 2005; Schumacher and Funnell 2005; Kusiak et al. 2011). Their loss in this plasmid may not affect its conjugation capability given that in this study we were able to obtain an *E. coli* transconjugant from a Mexican *P. rettgeri* strain harboring the p06-1619-NDM plasmid (fig. 1A and supplementary fig. S1E, Supplementary Material online); although, it is also possible that the other plasmid in this strain, p06-1619-2, which contains the putative *P. rettgeri* conjugation machinery (Tra-Pre), could act as a helper for the conjugation process, as has been reported for other

antibiotic resistance plasmids (Dery et al. 1997; Bennett 2008; Al-Marzooq et al. 2015). As well as the deletions described here, insertion of the putative novel mobile element has given the plasmid two toxin–antitoxin systems (*parED* and *reLEB*), *repA* and the IS elements IS5 and IS26 (fig. 5A).

Pairwise comparisons show there is no relationship between the three different *P. rettgeri* *bla_{NDM-1}*-positive plasmids circulating in Colombia and Mexico, apart from the Tn125 remains and the presence of multiple copies of the IS26 element (fig. 3). Only the p16Pre36-NDM plasmid shows some relationship with p6234-178kb-like plasmids found in Colombian *K. pneumoniae* strains (supplementary data set 2, Supplementary Material online, and fig. 5B). Both p16Pre36-NDM and p6234-178kb share the same Tra1 region, although the Tn125 (harboring *bla_{NDM-1}*) is located within a different genetic context (figs. 4 and 5B). These similarities explain the genetic relationship found among the *bla_{NDM-1}*-positive IncA/C2 plasmids and the *P. rettgeri* from Bogota (supplementary data set 2, Supplementary Material online). However, p6234-178kb does not have the Tra-Pre region found in p16Pre36-NDM, and has a more complex antibiotic resistance gene profile (table 1) due to the presence




Fig. 6.—Presence of bla_{NDM-1} -plasmids in the clinical isolates and transconjugants. Complete sequence of bla_{NDM-1} -plasmids circulating among Colombian and Mexican NDM-1-positive isolates and some related bla_{NDM-1} -positive (pPrY2001 and pNDM-BJ01) and bla_{NDM-1} -negative (p06-1619-2) plasmids are shown along the x axis. Black shading indicates a match of $\geq 90\%$ nucleotide identity in a window of 300 bp, calculated by comparing the query sequence (x axis, reference plasmids) against the consensus from mapped reads for each strain (y axis). Horizontal orange and light green bars represent the Tra1 and pPrY-like regions, respectively; blue rectangles represent the Tn125 (or remnants) region in each plasmid. Vertical bars correspond to: gray for *Klebsiella pneumoniae* from Bogota (Colombia), red for *Providencia rettgeri* from Bogota (Colombia), blue for *P. rettgeri* from Bucaramanga (Colombia), green for *P. rettgeri* from Monterrey (Mexico), purple for *Acinetobacter baumannii* from Cali (Colombia). Simulated reads for the reference plasmids were included as an internal control.

of an additional class 1 integron containing most of those genes.

Surprisingly, despite no previously reported epidemiological connection, a genetic link between *P. rettgeri* strains from Colombia, Mexico and Canada was found through analysis of the plasmids the strains harbor. We found the putative *P. rettgeri* conjugation machinery (Tra-Pre) and some additional regions, first identified in pPrY2001 from the Canadian *P. rettgeri* strain 09ACRGNY2001, were also present in the bla_{NDM-1} -negative p06-1619-2 plasmid from Mexico and in the bla_{NDM-1} positive p16Pre36-NDM plasmid from Bogota, Colombia (fig. 5B).

Genomic Epidemiology of the bla_{NDM-1} -Positive Plasmids Circulating in Colombia and Mexico

To gain a better understanding of the dissemination dynamics of bla_{NDM-1} -positive plasmids in Latin America, the MiSeq sequencing reads for all isolates included in the study were mapped against the fully sequenced bla_{NDM-1} -positive

plasmids from the representative Colombian and Mexican strains, as well as the related bla_{NDM-1} -positive plasmids pNDM-BJ01, pPrY2001 and the bla_{NDM-1} -negative p06-1619-2 (fig. 6).

The reads from four out of six *P. rettgeri* strains from Bogotá were found to map well over the entire sequence of p16Pre36-NDM (fig. 6), so are likely have bla_{NDM-1} in a plasmid closely related to p16Pre36-NDM. Of the remaining two *P. rettgeri* strains from Bogotá, one, 16Pre47, lacked the Tra-Pre locus and the mapped reads covered only one section of the p16Pre36-NDM sequence, suggesting that 16Pre47 carries a much smaller variant of that plasmid; the other, 16Pre45, is positive for the Tra-Pre region, but has no reads mapping to several other regions of p16Pre36-NDM, so it is possible that it has a pPrY-like plasmid or an unrelated plasmid harboring the bla_{NDM-1} gene. Additionally, the reads for the *E. coli* transconjugant strain C16-36, generated from 16Pre36, did not map to the whole of the donor plasmid sequence, covering only a 138,178 bp section of p16Pre36-NDM, from position 67,813–206,190 bp (fig. 6 and supplementary fig.

S1C, Supplementary Material online). This 138-kb region encodes the Tra1 region and Δ Tn125 harboring the *bla*_{NDM-1} and has a 2,928-bp direct repeat sequence at each end (99% nucleotide identity to each other), which encodes ISVsa3 (known as an ISCR2-like element). The mapping profile (fig. 6) showed no C16-36 reads mapped to the Tra-Pre region common to pPrY2001 and its related plasmids (fig. 5B). Using primers specific to the Tra-Pre region a PCR product was generated for the donor strain 16Pre36 but not the transconjugant C16-36 (data not shown), confirming the absence of the Tra-Pre region in the transconjugant was due to transfer of only part of the donor plasmid and not a problem with the WGS data. Taken together, these data indicate that p16Pre36-NDM contains a smaller, self-mobilizing transposable element, 138 kb in size, that can be transferred via conjugation to a new host, possibly via the self-encoded Tra1 conjugative apparatus, common to IncA/C2 plasmids.

The sequencing reads from all five *P. rettgeri* isolates from Bucaramanga and the corresponding representative transconjugants CRB151 and CRB152, mapped to almost the entire pRB151-NDM sequence, suggesting that they harbor *bla*_{NDM-1} in a plasmid closely related to pRB151-NDM (fig. 6). Plasmid pRB151-NDM does not encode the Tra-Pre and Tra1 regions and the strain RB151 does not have additional plasmids, yet four out of five of the related *P. rettgeri* isolates from Bucaramanga (including RB151) had sequences that mapped to the pPrY2001-like and IncA/C2 Tra1 regions (fig. 6). Investigation of the complete RB151 genome sequence (Marquez-Ortiz et al. 2017) revealed the presence of a genomic island with a high degree of similarity to p16Pre36-NDM, including the Tra-Pre and Tra1 regions, which is chromosomally inserted in strain RB151, flanked by 14.6-kb direct repeats (99% nucleotide identity) (see supplementary fig. S5A, Supplementary Material online). This chromosomal insertion was confirmed by PCR using specific primers (see supplementary fig. S5B, Supplementary Material online).

The read mapping data also showed that all three of the *P. rettgeri* Mexican isolates have both p06-1619-NDM and p06-1619-2 plasmids, and the reads from the representative transconjugant C06-1623 showed good coverage over the entire length of p06-1619-NDM, confirming the presence of *bla*_{NDM-1} on this plasmid.

Thus, the conjugative transfer-associated region Tra-Pre was observed to be present in 12 out of 14 NDM-1-positive *P. rettgeri* isolates (fig. 6), from Colombia (Bogota and Bucaramanga) and Mexico. In addition to the first report in the Canadian *P. rettgeri* *bla*_{NDM-1}-positive plasmid pPrY2001, this Tra-Pre region is also found in the partially sequenced genome of a NDM-positive *P. rettgeri* isolated in Israel, strain H1736 (Olaitan et al. 2015) (see supplementary fig. S2, Supplementary Material online). However, the partial assembly of the Israeli isolate prevented determination of whether or not the *bla*_{NDM-1} gene is located in the same plasmid as the Tra-Pre region. Together, these results validate a genetic link

among epidemiologically unrelated isolates of NDM-1-positive *P. rettgeri*.

Discussion

In this study, we used WGS data to provide a high-resolution picture of *bla*_{NDM-1} dissemination in Latin America, which led us to interesting findings about the dissemination route of this gene between *Enterobacteriaceae* and *Acinetobacter* species. *Acinetobacter* spp. harboring *bla*_{NDM-1} in pNDM-BJ01-like plasmids are frequently detected all over the world (Feng et al. 2015; Fu et al. 2015); these (or NDM-positive isolates with *bla*_{NDM-1} genetic surroundings suggesting the presence of pNDM-BJ01-like plasmids) have even been found in Latin America (Waterman et al. 2013; Pasteran et al. 2014; Brovedan et al. 2015; Quinones et al. 2015; Montana et al. 2016; Rojas et al. 2016). Here, we report an *A. baumannii* clone isolated in Colombia that also harbors *bla*_{NDM-1} in a pNDM-BJ01-like plasmid. This finding supports observations of such a *bla*_{NDM-1}-harboring plasmid present in *Acinetobacter* spp. of different genetic backgrounds. However, although the majority of plasmids harboring *bla*_{NDM} in *Acinetobacter* spp. are pNDM-BJ01-like plasmids, and although there is some evidence to suggest that *Acinetobacter* spp. passed *bla*_{NDM} on to the *Enterobacteriaceae* (Toleman et al. 2012), pNDM-BJ01-like plasmids do not seem to have good fitness in non-*Acinetobacter* bacteria, or at least in *Enterobacteriaceae*. To date, only one non-*Acinetobacter* harboring a pNDM-BJ01-like plasmid has been reported (Feng et al. 2015), whereas a plethora of *Enterobacteriaceae* hosting diverse, completely unrelated *bla*_{NDM}-positive plasmids have been found (supplementary data set 2, Supplementary Material online). The mechanisms of *bla*_{NDM} gene transmission from *Acinetobacter* spp. to *Enterobacteriaceae* are not yet understood.

In this study, we identified a second *Enterobacteriaceae* family member harboring a pNDM-BJ01-like plasmid (p06-1619-NDM), a *P. rettgeri* isolated in an outbreak in Mexico (Barrios et al. 2013). However, the p06-1619-NDM plasmid has suffered a major modification in the variable region, with the insertion of a previously unreported mobile element introducing two toxin–antitoxin systems (flanked by two novel Tn6368 transposons). It seems plausible that the toxin–antitoxin systems have generated a strong dependence on that plasmid as has been previously reported (Kamruzzaman et al. 2017), thereby avoiding transposition of *bla*_{NDM-1} to another more compatible plasmid and its subsequent elimination. Adding to the advantages conferred by the two addictive systems is the high selective pressure of the environment—an Intensive Care Unit—from which the Mexican *P. rettgeri* strains were isolated. As the other plasmid hosted by these isolates (p06-1619-2) does not have any resistance genes (table 1), selective pressure could force permanent residence of the pNDM-BJ01-like plasmid due to the conferred

resistance to aminoglycosides and beta-lactams, including carbapenems.

Providencia species are frequently found in environmental settings, but are also opportunistic human pathogens, mainly as the causative agents of urinary tract infections (Wie 2015). They are not among the most significant or prevalent human threats, but recently they have been attracting interest due to increasing reports of *P. rettgeri* NDM-positive isolates found around the world (Barrios et al. 2013; Carvalho-Assef et al. 2013; Mataseje et al. 2014; Pollett et al. 2014; Tada et al. 2014; Carmo Junior et al. 2015; Manageiro et al. 2015; Nachimuthu et al. 2015; Wailan, Paterson, et al. 2016). Here, we also found *P. rettgeri* to be the most frequent bacteria harboring the bla_{NDM-1} gene in three hospitals at distant locations from each other in Colombia. Most of the cases correspond to outpatients, suggesting that NDM-1-positive *P. rettgeri* strains are present in the community in Colombia. In spite of the increase in NDM-positive *P. rettgeri* cases, only two completely sequenced bla_{NDM-1} -positive plasmids are in the NCBI nucleotide database for *P. rettgeri*, both of which were isolated in Canada (supplementary data set 2, Supplementary Material online). Here, we report three additional unrelated complete plasmids hosted by *P. rettgeri* (two from Colombia and one from Mexico), providing more information to help elucidate bla_{NDM-1} dissemination in the *Enterobacteriaceae*.

Interestingly, despite the geographic distances between the sites of isolation of NDM-1-positive *P. rettgeri* strains and despite the very different structures of their bla_{NDM-1} -positive plasmids, we found a common feature—a putative conjugative transfer region named here as Tra-Pre—that appears to be stable among *P. rettgeri* from different regions. Supporting this hypothesis of a common, stable feature for bla_{NDM-1} -positive *P. rettgeri*, is the fact that this Tra-Pre region is also found in the partially sequenced genome of the NDM-1-positive *P. rettgeri* H1736, reported in Israel in 2011 (Olaitan et al. 2015).

The Tra-Pre-family plasmids harboring bla_{NDM-1} that are hosted by *P. rettgeri* (pPrY2001 and p16Pre36-NDM) are unrelated to the pNDM-BJ01-like plasmids from *Acinetobacter* species. Nevertheless, it is possible the bla_{NDM-1} -positive Tra-Pre-encoding plasmids emerged in *P. rettgeri* through transposition of bla_{NDM-1} from a pNDM-BJ01-like plasmid (prior to its loss) to a more stable plasmid, as suggested by the coexistence of a bla_{NDM-1} -negative plasmid containing the Tra-Pre region (p06-1619-2) and a pNDM-BJ01-like plasmid in the *P. rettgeri* from Mexico. This proposed mechanism is further supported by the presence of the Tra-Pre region in almost all *P. rettgeri* isolates in this study (12 out of 14), even isolates in which bla_{NDM-1} is found on an unrelated plasmid, for example in the *P. rettgeri* isolates from Bucaramanga, Colombia. The simultaneous detection in the Bucaramanga *P. rettgeri* isolates of a new plasmid harboring the bla_{NDM-1} , and of a region in the bacterial chromosome encoding both

the IncA/C2-related and Tra-Pre regions similar to those found in the bla_{NDM-1} -positive plasmid circulating in Bogota, indicates how a possible transposition of the bla_{NDM-1} region to a new, different backbone may have occurred in this strain. Thus, our study supports the role of gene module transposition in the spread of bla_{NDM-1} among *P. rettgeri* clinical isolates, a role that has been identified as relevant to the evolution of bla_{NDM-1} -positive plasmids (Khong et al. 2016).

We found a further interesting genetic link in the relationship between bla_{NDM-1} -positive plasmids present in *P. rettgeri* (p16Pre36-NDM) and those found in *K. pneumoniae* (p6234-178kb). The isolates harboring p16Pre36-NDM and p6234-178kb were detected in the same Colombian hospital (supplementary data set 1, Supplementary Material online). Although p16Pre36-NDM and p6234-178kb cannot be classified in the same Inc group, they share a large region, commonly found in the IncA/C2 bla_{NDM-1} -positive and negative plasmids from diverse *Enterobacteriaceae*. This region was also found to be inserted in the chromosome of the *P. rettgeri* isolates from Bucaramanga. These results suggest that the complex p16Pre36-NDM plasmid originated in *P. rettgeri*, through the co-integration of a pPrY2001-like plasmid with an acquired IncA/C2 broad host range plasmid from a different *Enterobacteriaceae*. This IncA/C2 plasmid may then have been transferred to (or from) a non-*Providencia* *Enterobacteriaceae*, such as the *K. pneumoniae* in this study. Conjugation of p16Pre36-NDM to *E. coli* J53 resulted in transfer of only part of the plasmid, a putative self-mobilizable ISVsa3 (an ISCR2-like element) composite transposon encoding the bla_{NDM-1} and Tra1 regions, but not the Tra-Pre region. The mapping data indicates that the clinical isolate 16Pre47 also only contains this ISVsa3 composite transposon, and is missing the remainder of p16Pre36-NDM, suggesting this Tra-Pre-negative strain 16Pre47 may have received the ISVsa3 composite transposon from other *P. rettgeri* strain (through partial conjugation) or from a *K. pneumoniae*. This putative bla_{NDM-1} -positive conjugative transposon may have derived from the p6234.178kb circulating in *K. pneumoniae* in Colombia prior to the isolation of the *P. rettgeri*, given their similarity and the presence of a closely related sequence containing ISVsa3 (2,928 bp) in both p6234.178kb and p16Pre36-NDM (figs. 4 and 5). Interestingly, this transposable element was found to be conserved (99% nucleotide identity over the 2,928 bp) in the genomes of a very wide range of bacteria, in a search against the NCBI database, and also ISVsa3-like elements have been recognized as key players in IncA/C plasmids evolution (Toleman and Walsh 2010). Therefore, the novel bla_{NDM-1} -positive putative conjugative transposon identified in this study could facilitate broad dissemination of bla_{NDM-1} through transposition, conjugation and integration of the transferred circular intermediate into the host genome or to other plasmids via homologous recombination.

The low levels of similarity in the vicinity of Tn125 (or its remnants) between p16Pre36-NDM and p6234-178kb could




Fig. 7.—Possible roles of *Providencia rettgeri* in bla_{NDM-1} -plasmids evolution in Latin America. In an initial stage pNDM-BJ01-like plasmids are acquired from *Acinetobacter* spp. (–1). Shortly after, bla_{NDM-1} is transposed to pPrY-like plasmids (from *P. rettgeri* circulation; –2) or IncA/C2 plasmids (from *Klebsiella pneumoniae*, *Escherichia coli* or other *Enterobacteriaceae*; –3) via Tn125 transposition or by mean of other mobile genetic elements surrounding the Tn125 (or its remnants). IncA/C2 bla_{NDM-1} -plasmids could be transferred to a broad bacteria host range (–4). It is also possible that a non-*Providencia Enterobacteriaceae* could capture a pNDM-BJ01-like plasmid and transposes Tn125 to a broad host range IncA/C2 plasmid (–5); later this IncA/C2 bla_{NDM-1} -plasmid could be conjugated to *P. rettgeri* (–6). An interesting finding of the present study is the generation in *P. rettgeri* of new plasmids by mean of co-integration of pPrY-like plasmids and IncA/C2 plasmids (–7). These chimeric structures can also be transposed to the *P. rettgeri* chromosome (–8). The Tn125 (or its remnants) could be transposed to new plasmid backbones with possible implications upon its dissemination (–9). Additionally, by mean of partial conjugation could be disseminated IncA/C2-related (*repA* negative) bla_{NDM-1} -plasmids (–10).

be due to the later isolation of the *P. rettgeri* strain harboring p16Pre36-NDM. Although both the *P. rettgeri* and the *K. pneumoniae* strains were isolated at the same hospital (supplementary data set 1, Supplementary Material online), *P. rettgeri* harboring p16Pre36-NDM was isolated more than a year after the *K. pneumoniae* outbreak. The length of time the *P. rettgeri* were present in the hospital under selective pressure, could promote the genetic rearrangements observed in the p16Pre36-NDM plasmid. In an earlier isolate of *K. pneumoniae*, the bla_{NDM-1} surroundings on the p6234-178-kb plasmid are more closely related to those in *Acinetobacter* species.

Dissemination of the IncA/C2 p6234-178kb plasmid in different cities and in strains of different genetic backgrounds, is consistent with the globally observed trend for IncA/C2 (recently designated as IncC; Harmer et al. 2016) group plasmids, which are known to be associated with spread of multidrug resistance genes to different countries and different bacterial hosts (Fricke et al. 2009; Roy Chowdhury et al. 2011; Carattoli et al. 2012), meaning they post a significant risk to human health, particularly as they are capable of disseminating bla_{NDM-1} . The importance of improving existing infection control measures, such as isolation of patients harboring

resistant pathogens and hand hygiene, is further substantiated by our findings suggesting that some bla_{NDM-1} -positive plasmids are likely to have originated by co-integration of less stable bla_{NDM-1} -positive plasmids with more stable and disseminative plasmids in environmental bacteria. A case in point is found in the *P. rettgeri* isolates, in which the Tn125 (or its remains) may have transposed from pNDM-BJ01-like plasmids to pPrY2001-like or IncA/C2-related plasmids that could subsequently be transferred to other bacterial species, including more problematic non-*Providencia Enterobacteriaceae*. An important scenario where all these factors can be found simultaneously is in the mammalian gut where *Enterobacteriaceae* can thrive, aiding the inter-species and inter-genera dissemination of the NDM-1 antibiotic resistance gene among the bacterial community of the gut. Host-specific conditions, such as the inflammatory host response, can also boost horizontal gene transfer and hence microbiota evolution (Stecher et al. 2012), that may have led to the plasmid rearrangements observed here, probably under the selective pressure of the hospital environment. However, direct transfer from *Acinetobacter* spp. to non-*Providencia Enterobacteriaceae* cannot be ruled out, due to conjugation

of pNDM-BJ01-like plasmids from *Acinetobacter* spp. to non-*Providencia* *Enterobacteriaceae* (mainly *E. coli*) has been demonstrated in this and other studies (Hu et al. 2012; Huang et al. 2015). However, more work needs to be done to better understand the genetic basis of the dissemination of *bla*_{NDM-1}-positive plasmids in Latin America, by evaluating the stability, fitness cost and conjugation capability of pNDM-BJ01-like and pPrY2001-like plasmids to other *Enterobacteriaceae*.

In our analysis of NDM-1-positive clinical isolates in Latin America, the high variability of *bla*_{NDM-1}-positive plasmids present in different species, highlights that *bla*_{NDM-1}-dissemination has not only followed a predominantly clonal evolution, but rather a Russian doll model (Sheppard et al. 2016). In this Russian doll model, a resistance gene such as *bla*_{NDM-1} resides on nested transmissible units and therefore can move through the environment at multiple different levels, that can be both coincident and independent of one another. For example, bacterial cell hosting resistance gene; plasmid within bacterial cell harboring resistance gene; mobile element within plasmid harboring resistance gene; and mobile element within mobile element harboring resistance gene, with the consequence that the resistance gene may move between plasmids within a bacterial cell via multiple mechanisms (Sheppard et al. 2016). In line with this model, different clones have acquired different *bla*_{NDM-1}-positive plasmids and related strains have disseminated locally, as for example in the NDM-1 outbreaks in Colombia and Mexico (Barrios et al. 2013; Escobar Perez et al. 2013) and also the cases of the *P. rettgeri* isolated in Bogota and Bucaramanga (Colombia) in the recent surveillance study. These related strains acquired the *bla*_{NDM-1} from a variety of plasmids, such as IncA/C2-related or pPrY2001-like plasmids, that in turn received *bla*_{NDM-1} from plasmid co-integration or transposition from another plasmid, or from *Acinetobacter* spp. pNDM-BJ01-like plasmids in an initial dissemination stage. In our Russian doll model, *P. rettgeri* plays an important role as a reservoir of *bla*_{NDM-1} available for transmission into highly disseminative plasmids due to its high recombination capability supported by the high plasmid variability found in this species (fig. 7). In this study, the presence of pNDM-BJ01-like, Tra-Pre-encoding and IncA/C2-related plasmids or genetic structures in *P. rettgeri*, and their relationship with the plasmids present in *K. pneumoniae* and *Acinetobacter* species, illustrates the evolution route of *bla*_{NDM-1}-positive plasmids in Latin America, where *P. rettgeri* appears to be crucial for *bla*_{NDM-1} transmission from *Acinetobacter* spp. to *Enterobacteriaceae*.

Taken together, these findings expose the role of microorganisms such as *P. rettgeri*, that generally are not the target of public health surveillance systems, in the dissemination and storage of resistance genes, highlighting the importance of more comprehensive studies, which do not merely focus on the most frequently occurring pathogens but also encompass the resistance determinants and their mobilization machinery.

Supplementary Material

Supplementary data are available at *Genome Biology and Evolution* online.

Acknowledgments

We gratefully acknowledge financial support from the Departamento Administrativo de Ciencia, Tecnología e Innovación, Colciencias (grant number 130871250819 and Fellowship 567-2012 to R.A.M.); Vicerrectoría de Investigaciones, Universidad El Bosque (grant number PCI-2012-330) and University of Technology Sydney. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author Contributions

R.A.M., J.E. and N.K.P. designed research; R.A.M., L.H. and N.K.P. performed research; N.O., C.D., U.G.R., J.S.S., B.E.C., E.M.S., M.B., M.V.M., J.E.C. and A.V. contributed new reagents/analytic tools; R.A.M., J.E. and N.K.P. analyzed data; R.A.M., I.G.C., N.V., J.E. and N.K.P. conceived the study; and R.A.M. and N.K.P. wrote the paper.

Literature Cited

- Al-Marzooq F, Mohd Yusof MY, Tay ST. 2015. Molecular analysis of antibiotic resistance determinants and plasmids in Malaysian isolates of multidrug resistant *Klebsiella pneumoniae*. *PLoS One* 10:e0133654.
- Alikhan NF, Petty NK, Ben Zakour NL, Beatson SA. 2011. BLAST Ring Image Generator (BRIG): simple prokaryote genome comparisons. *BMC Genomics* 12:402.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol.* 215:403–410.
- Bankevich A, et al. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol.* 19:455–477.
- Barrios H, et al. 2013. Isolation of carbapenem-resistant NDM-1-positive *Providencia rettgeri* in Mexico. *J Antimicrob Chemother.* 68:1934–1936.
- Bennett PM. 2008. Plasmid encoded antibiotic resistance: acquisition and transfer of antibiotic resistance genes in bacteria. *Br J Pharmacol.* 153(Suppl 1):S347–S357.
- Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27:578–579.
- Boetzer M, Pirovano W. 2012. Toward almost closed genomes with GapFiller. *Genome Biol.* 13:1.
- Brovedan M, et al. 2015. Complete sequence of a *bla*_{NDM-1}-harboring plasmid in an *Acinetobacter bereziniae* clinical strain isolated in Argentina. *Antimicrob Agents Chemother.* 59:6667–6669.
- Carattoli A, Villa L, Poirel L, Bonnin RA, Nordmann P. 2012. Evolution of IncA/C *bla*_{CMY-2}-carrying plasmids by acquisition of the *bla*_{NDM-1} carbapenemase gene. *Antimicrob Agents Chemother.* 56:783–786.
- Carattoli A, et al. 2014. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother.* 58:3895–3903.
- Carmo Junior NV, et al. 2015. First report of a NDM-producing *Providencia rettgeri* strain in the state of Sao Paulo. *Braz J Infect Dis.* 19:675–676.
- Carvalho-Assef AP, et al. 2013. Isolation of NDM-producing *Providencia rettgeri* in Brazil. *J Antimicrob Chemother.* 68:2956–2957.

- Carver TJ, et al. 2005. ACT: the Artemis Comparison Tool. *Bioinformatics* 21:3422–3423.
- CDC 2013. Vital signs: carbapenem-resistant *Enterobacteriaceae*. *MMWR Morb Mortal Wkly Rep.* 62:165–170.
- Chain PS, et al. 2009. Genomics. Genome project standards in a new era of sequencing. *Science* 326:236–237.
- Chen Z, et al. 2015. NDM-1 encoded by a pNDM-BJ01-like plasmid p3SP-NDM in clinical *Enterobacter aerogenes*. *Front Microbiol.* 6:294.
- Chin CS, et al. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569.
- Christie PJ, Atmakuri K, Krishnamoorthy V, Jakubowski S, Cascales E. 2005. Biogenesis, architecture, and function of bacterial type IV secretion systems. *Annu Rev Microbiol.* 59:451–485.
- David M, Dzamba M, Lister D, Ilie L, Brudno M. 2011. SHRIMP2: sensitive yet practical SHort Read Mapping. *Bioinformatics* 27:1011–1012.
- Dery KJ, et al. 1997. Characterization of the replication and mobilization regions of the multiresistance *Klebsiella pneumoniae* plasmid pJHCMW1. *Plasmid* 38:97–105.
- Diancourt L, Passet V, Nemeč A, Dijkshoorn L, Brisse S. 2010. The population structure of *Acinetobacter baumannii*: expanding multiresistant clones from an ancestral susceptible genetic pool. *PLoS One* 5:e10034.
- Diene SM, et al. 2013. The rhizome of the multidrug-resistant *Enterobacter aerogenes* genome reveals how new "killer bugs" are created because of a sympatric lifestyle. *Mol Biol Evol.* 30:369–383.
- Doublet B, et al. 2012. Complete nucleotide sequence of the multidrug resistance IncA/C plasmid pR55 from *Klebsiella pneumoniae* isolated in 1969. *J Antimicrob Chemother.* 67:2354–2360.
- Escobar Perez JA, et al. 2013. Outbreak of NDM-1-producing *Klebsiella pneumoniae* in a neonatal unit in Colombia. *Antimicrob Agents Chemother.* 57:1957–1960.
- Espinal P, et al. 2011. Dissemination of an NDM-2-producing *Acinetobacter baumannii* clone in an Israeli rehabilitation center. *Antimicrob Agents Chemother.* 55:5396–5398.
- Espinal P, et al. 2015. Identification of NDM-1 in a putatively novel *Acinetobacter* species ("NB14") closely related to *Acinetobacter pittii*. *Antimicrob Agents Chemother.* 59:6657–6660.
- Feng J, et al. 2015. Coexistence of a novel KPC-2-encoding MDR plasmid and an NDM-1-encoding pNDM-HN380-like plasmid in a clinical isolate of *Citrobacter freundii*. *J Antimicrob Chemother.* 70:2987–2991.
- Fernandez-Alarcon C, Singer RS, Johnson TJ. 2011. Comparative genomics of multidrug resistance-encoding IncA/C plasmids from commensal and pathogenic *Escherichia coli* from multiple animal sources. *PLoS One* 6:e23415.
- Finn RD, et al. 2016. The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Res.* 44:D279–D285.
- Fricke WF, et al. 2009. Comparative genomics of the IncA/C multidrug resistance plasmid family. *J Bacteriol.* 191:4750–4757.
- Fu Y, et al. 2015. Spread of a common *bla*_{NDM-1}-carrying plasmid among diverse *Acinetobacter* species. *Infect Genet Evol.* 32:30–33.
- Harmer CJ, Hamidian M, Ambrose SJ, Hall RM. 2016. Destabilization of IncA and IncC plasmids by SGI1 and SGI2 type Salmonella genomic islands. *Plasmid* 87–88:51–57.
- He S, et al. 2015. Insertion sequence IS26 reorganizes plasmids in clinically isolated multidrug-resistant bacteria by replicative transposition. *MBio* 6:e00762.
- Hu H, et al. 2012. Novel plasmid and its variant harboring both a *bla*_{NDM-1} gene and type IV secretion system in clinical isolates of *Acinetobacter lwoffii*. *Antimicrob Agents Chemother.* 56:1698–1702.
- Huang TW, et al. 2015. Effective transfer of a 47 kb NDM-1-positive plasmid among *Acinetobacter* species. *J Antimicrob Chemother.* 70:2734–2738.
- Huang W, Li L, Myers JR, Marth GT. 2012. ART: a next-generation sequencing read simulator. *Bioinformatics* 28:593–594.
- Hunt M, et al. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol.* 16:294.
- Huson DH, Scornavacca C. 2012. Dendroscope 3: an interactive tool for rooted phylogenetic trees and networks. *Syst Biol.* 61:1061–1067.
- Johnson AP, Woodford N. 2013. Global spread of antibiotic resistance: the example of New Delhi metallo-beta-lactamase (NDM)-mediated carbapenem resistance. *J Med Microbiol.* 62:499–513.
- Jones LS, et al. 2015. Characterization of plasmids in extensively drug-resistant *Acinetobacter* strains isolated in India and Pakistan. *Antimicrob Agents Chemother.* 59:923–929.
- Kamruzzaman M, Shoma S, Thomas CM, Partridge SR, Iredell JR. 2017. Plasmid interference for curing antibiotic resistance plasmids in vivo. *PLoS One* 12:e0172913.
- Khong WX, et al. 2016. Local transmission and global dissemination of New Delhi Metallo-Beta-Lactamase (NDM): a whole genome analysis. *BMC Genomics* 17:452.
- Kusiak M, Gapczynska A, Plochocka D, Thomas CM, Jagura-Burdzy G. 2011. Binding and spreading of ParB on DNA determine its biological function in *Pseudomonas aeruginosa*. *J Bacteriol.* 193:3342–3355.
- Lemos EV, et al. 2014. Impact of carbapenem resistance on clinical and economic outcomes among patients with *Acinetobacter baumannii* infection in Colombia. *Clin Microbiol Infect.* 20:174–180.
- Li P, et al. 2015. *Acinetobacter calcoaceticus* from a fatal case of pneumonia harboring *bla*_{NDM-1} on a widely distributed plasmid. *BMC Infect Dis.* 15:131.
- Liebert CA, Hall RM, Summers AO. 1999. Transposon Tn21, flagship of the floating genome. *Microbiol Mol Biol Rev.* 63:507–522.
- Loytynoja A. 2014. Phylogeny-aware alignment with PRANK. *Methods Mol Biol.* 1079:155–170.
- Manageiro V, et al. 2015. Draft genome sequence of the first NDM-1-producing *Providencia stuartii* strain isolated in Portugal. *Genome Announc.* 3:e01077–15.
- Marquez C, et al. 2008. Urinary tract infections in a South American population: dynamic spread of class 1 integrons and multidrug resistance by homologous and site-specific recombination. *J Clin Microbiol.* 46:3417–3425.
- Marquez-Ortiz RA, et al. 2017. First complete *Providencia rettgeri* genome sequence, the NDM-1-producing clinical strain RB151. *Genome Announc.* 5:e01472–16.
- Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* 17:10–12.
- Mataseje LF, et al. 2014. Complete sequences of a novel *bla*_{NDM-1}-harbouring plasmid from *Providencia rettgeri* and an FII-type plasmid from *Klebsiella pneumoniae* identified in Canada. *J Antimicrob Chemother.* 69:637–642.
- Milne I, et al. 2013. Using Tablet for visual exploration of second-generation sequencing data. *Brief Bioinform.* 14:193–202.
- Montana S, et al. 2016. Presence of New Delhi metallo-beta-lactamase gene (NDM-1) in a clinical isolate of *Acinetobacter junii* in Argentina. *New Microbes New Infect.* 11:43–44.
- Nachimuthu R, et al. 2015. Characterization of carbapenem-resistant Gram-negative bacteria from Tamil Nadu. *J Chemother.* 28:371–374.
- Noguchi N, Katayama J, Sasatsu M. 2000. A transposon carrying the gene *mphB* for macrolide 2'-phosphotransferase II. *FEMS Microbiol Lett.* 192:175–178.
- Olaitan AO, Diene SM, Assous MV, Rolain JM. 2015. Genomic plasticity of multidrug-resistant NDM-1 positive clinical isolate of *Providencia rettgeri*. *Genome Biol Evol.* 8:723–728.
- Page AJ, et al. 2015. Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics* 31:3691–3693.
- Palzkill T. 2013. Metallo-beta-lactamase structure and function. *Ann N Y Acad Sci.* 1277:91–104.
- Partridge SR, Iredell JR. 2012. Genetic contexts of *bla*_{NDM-1}. *Antimicrob Agents Chemother.* 56:6065–6067. author reply 6071.

- Partridge SR, Recchia GD, Stokes HW, Hall RM. 2001. Family of class 1 integrons related to In4 from Tn1696. *Antimicrob Agents Chemother.* 45:3014–3020.
- Pasteran F, et al. 2014. Emergence of genetically unrelated NDM-1-producing *Acinetobacter pittii* strains in Paraguay. *J Antimicrob Chemother.* 69:2575–2578.
- Poirel L, Dortet L, Bernabeu S, Nordmann P. 2011. Genetic features of bla_{NDM-1}-positive *Enterobacteriaceae*. *Antimicrob Agents Chemother.* 55:5403–5407.
- Pollett S, et al. 2014. Phenotypic and molecular characteristics of carbapenem-resistant *Enterobacteriaceae* in a health care system in Los Angeles, California, from 2011 to 2013. *J Clin Microbiol.* 52:4003–4009.
- Prabaker K, Weinstein RA. 2011. Trends in antimicrobial resistance in intensive care units in the United States. *Curr Opin Crit Care* 17:472–479.
- Quinones D, et al. 2015. High prevalence of bla_{OXA-23} in *Acinetobacter* spp. and detection of bla_{NDM-1} in *A. soli* in Cuba: report from National Surveillance Program (2010–2012). *New Microbes New Infect.* 7:52–56.
- Rhomberg PR, Jones RN. 2009. Summary trends for the Meropenem Yearly Susceptibility Test Information Collection Program: a 10-year experience in the United States (1999–2008). *Diagn Microbiol Infect Dis.* 65:414–426.
- Roberts AP, et al. 2008. Revised nomenclature for transposable genetic elements. *Plasmid* 60:167–173.
- Rojas LJ, et al. 2016. Initial assessment of the molecular epidemiology of bla_{NDM-1} in Colombia. *Antimicrob Agents Chemother.* 60:4346–4350.
- Roy Chowdhury P, et al. 2011. Dissemination of multiple drug resistance genes by class 1 integrons in *Klebsiella pneumoniae* isolates from four countries: a comparative study. *Antimicrob Agents Chemother.* 55:3140–3149.
- Rutherford K, et al. 2000. Artemis: sequence visualization and annotation. *Bioinformatics* 16:944–945.
- Schmieder R, Edwards R. 2011. Quality control and preprocessing of metagenomic datasets. *Bioinformatics* 27:863–864.
- Schumacher MA, Funnell BE. 2005. Structures of ParB bound to DNA reveal mechanism of partition complex formation. *Nature* 438:516–519.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069.
- Sekizuka T, et al. 2011. Complete sequencing of the bla_{NDM-1}-positive IncA/C plasmid from *Escherichia coli* ST38 isolate suggests a possible origin from plant pathogens. *PLoS One* 6:e25334.
- Sheppard AE, et al. 2016. Nested Russian doll-like genetic mobility drives rapid dissemination of the carbapenem resistance gene blaKPC. *Antimicrob Agents Chemother.* 60:3767–3778.
- Siguier P, Varani A, Perochon J, Chandler M. 2012. Exploring bacterial insertion sequences with ISfinder: objectives, uses, and future developments. *Methods Mol Biol.* 859:91–103.
- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30:1312–1313.
- Stecher B, et al. 2012. Gut inflammation can boost horizontal gene transfer between pathogenic and commensal *Enterobacteriaceae*. *Proc Natl Acad Sci U S A.* 109:1269–1274.
- Sullivan MJ, Petty NK, Beatson SA. 2011. Easyfig: a genome comparison visualizer. *Bioinformatics* 27:1009–1010.
- Sun Y, et al. 2013. Complete genome sequence of an *Acinetobacter* strain harboring the NDM-1 Gene. *Genome Announc.* 1:e0002312.
- Tada T, et al. 2014. NDM-1 Metallo-beta-Lactamase and ArmA 16S rRNA methylase producing *Providencia rettgeri* clinical isolates in Nepal. *BMC Infect Dis.* 14:56.
- Talavera G, Castresana J. 2007. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Syst Biol.* 56:564–577.
- Tijet N, Richardson D, MacMullin G, Patel SN, Melano RG. 2015. Characterization of multiple NDM-1-producing *Enterobacteriaceae* isolates from the same patient. *Antimicrob Agents Chemother.* 59:3648–3651.
- Toleman MA, Spencer J, Jones L, Walsh TR. 2012. bla_{NDM-1} is a chimera likely constructed in *Acinetobacter baumannii*. *Antimicrob Agents Chemother.* 56:2773–2776.
- Toleman MA, Walsh TR. 2010. ISCR elements are key players in IncA/C plasmid evolution. *Antimicrob Agents Chemother.* 54:3534; author reply 3534.
- Valverde A, et al. 2006. In117, an unusual In0-like class 1 integron containing CR1 and bla_{CTX-M-2} and associated with a Tn21-like element. *Antimicrob Agents Chemother.* 50:799–802.
- van Duijn PJ, Dautzenberg MJ, Oostdijk EA. 2011. Recent trends in antibiotic resistance in European ICUs. *Curr Opin Crit Care* 17:658–665.
- Wailan AM, Paterson DL, et al. 2016. Genomic characteristics of NDM-producing *Enterobacteriaceae* isolates in Australia and their bla_{NDM} genetic contexts. *Antimicrob Agents Chemother.* 60:136–141.
- Wailan AM, Sidjabat HE, et al. 2016. Mechanisms involved in acquisition of bla_{NDM} genes by IncA/C2 and IncFIY Plasmids. *Antimicrob Agents Chemother.* 60:4082–4088.
- Wang R, et al. 2015. IncA/C plasmids harboured in serious multidrug-resistant *Vibrio cholerae* serogroup O139 strains in China. *Int J Antimicrob Agents* 45:249–254.
- Wang X, et al. 2014. Complete genome sequence of *Acinetobacter baumannii* ZW85-1. *Genome Announc.* 2:e01083–13.
- Wasyl D, Kern-Zdanowicz I, Domanska-Blicharz K, Zajac M, Hoszowski A. 2015. High-level fluoroquinolone resistant *Salmonella enterica* serovar Kentucky ST198 epidemic clone with IncA/C conjugative plasmid carrying bla_{CTX-M-25} gene. *Vet Microbiol.* 175:85–91.
- Waterman PE, et al. 2013. Bacterial peritonitis due to *Acinetobacter baumannii* sequence type 25 with plasmid-borne new delhi metallo-beta-lactamase in Honduras. *Antimicrob Agents Chemother.* 57:4584–4586.
- Wie SH. 2015. Clinical significance of *Providencia* bacteremia or bacteriuria. *Korean J Intern Med.* 30:167–169.
- Woodford N, Turton JF, Livermore DM. 2011. Multiresistant Gram-negative bacteria: the role of high-risk clones in the dissemination of antibiotic resistance. *FEMS Microbiol Rev.* 35:736–755.
- World Health Organization. 2014. Antimicrobial resistance: global report on surveillance. Geneva: World Health Organization.
- Yong D, et al. 2009. Characterization of a new metallo-beta-lactamase gene, bla_{NDM-1}, and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother.* 53:5046–5054.
- Yurieva O, et al. 1997. Intercontinental spread of promiscuous mercury-resistance transposons in environmental bacteria. *Mol Microbiol.* 24:321–329.
- Zhang WJ, et al. 2013. Complete sequence of the bla_{NDM-1}-carrying plasmid pNDM-AB from *Acinetobacter baumannii* of food animal origin. *J Antimicrob Chemother.* 68:1681–1682.

Associate editor: Bill F. Martin