

# Predation Response of *Vibrio fischeri* Biofilms to Bacterivorous Protists

Alba Chavez-Dozal,<sup>a</sup> Clayton Gorman,<sup>a</sup> Martina Erken,<sup>b</sup> Peter D. Steinberg,<sup>c,d</sup> Diane McDougald,<sup>b,e</sup> Michele K. Nishiguchi<sup>a</sup>

Department of Biology, New Mexico State University, Las Cruces, New Mexico, USA<sup>a</sup>; Centre for Marine Biofouling and Bio-Innovation and School of Biotechnology and Biomolecular Sciences, University of South Wales, Sydney, New South Wales, Australia<sup>b</sup>; Centre for Marine Biofouling and Bio-Innovation and School of Biological, Earth and Environmental Science, University of South Wales, Sydney, New South Wales, Australia<sup>c</sup>; Sydney Institute of Marine Science, Mosman, NSW, Australia<sup>d</sup>; Advanced Environmental Biotechnology Centre, Nanyang Technological University, Singapore<sup>e</sup>

***Vibrio fischeri* proliferates in a sessile, stable community known as a biofilm, which is one alternative survival strategy of its life cycle. Although this survival strategy provides adequate protection from abiotic factors, marine biofilms are still susceptible to grazing by bacteria-consuming protozoa. Subsequently, grazing pressure can be controlled by certain defense mechanisms that confer higher biofilm antipredator fitness. In the present work, we hypothesized that *V. fischeri* exhibits an antipredator fitness behavior while forming biofilms. Different predators representing commonly found species in aquatic populations were examined, including the flagellates *Rhynchomonas nasuta* and *Neobodo designis* (early biofilm feeders) and the ciliate *Tetrahymena pyriformis* (late biofilm grazer). *V. fischeri* biofilms included isolates from both seawater and squid hosts (*Euprymna* and *Sepiolo* species). Our results demonstrate inhibition of predation by biofilms, specifically, isolates from seawater. Additionally, antiprotozoan behavior was observed to be higher in late biofilms, particularly toward the ciliate *T. pyriformis*; however, inhibitory effects were found to be widespread among all isolates tested. These results provide an alternative explanation for the adaptive advantage and persistence of *V. fischeri* biofilms and provide an important contribution to the understanding of defensive mechanisms that exist in the out-of-host environment.**

Biofilms form at almost every surface that is in contact with water, and they are a natural part of aquatic ecosystems (1, 2). Moreover, biofilm communities offer a refuge toward diverse stresses, such as antibiotics (3–5), dehydration and osmotic stress (5, 6), UV light exposure (6, 7), and starvation (3). The pronounced stress resistance of biofilms has been observed to be prevalent in marine communities, particularly ones comprised of environmental *Vibrio* biofilms (2, 8, 9).

Biofilms formed by *Vibrio* species are ubiquitous in aquatic ecosystems, although no study has specified the prevalence of *Vibrio fischeri* biofilms. *V. fischeri* (and other marine bacteria) in its planktonic state are found over broad geographical ranges, and their biofilms are subjected to multiple physiological stresses that lead to alterations in bacterial physiology (promoting bacterial fitness and bacterial speciation) (10). Thus, the survival of planktonic cells and biofilms in the environment is not only defined by the capacity to overcome abiotic pressures but also by the ability to serve as a protective niche against natural protozoan consumers (11, 12). Thus, an alternative function for the prevalence and relative fitness of *V. fischeri* biofilms is that they serve as refuges to combat a range of predators, including protozoan grazers. Grazing is one of the most common mortality factors of bacterial populations (13–15) and causes rapid changes in the morphology and species composition of microbial communities (13, 16–18). Interactions between bacteria and protozoa within biofilm communities remain largely unexplored; however, recent studies have revealed the impact of grazing on the dynamics of natural biofilm communities. Quorum sensing is an important factor for anti-predatory activity in many bacterial species (19–22), and observations have suggested that bacterial genetic diversity enhances grazing resistance (23). Grazing resistance was observed in biofilms of *Pseudomonas aeruginosa* when early and late biofilm communities exhibited antipredatory behavior against two flagellates (*Bodo saltans* and *Rhynchomonas nasuta*) and the ciliate *Tetrahymena* sp. (18). Similar results were also observed when communities of

*Vibrio cholerae* prevented predation through an antiprotozoal factor regulated by the response regulator HapR (14, 20). Moreover, *V. cholerae* biofilms exhibit widespread grazing resistance among toxigenic and nontoxigenic isolates, which has an impact on strain distribution and cholera epidemics (20). Previous studies also reported that some bacterial communities synthesize chemical compounds (e.g., violacein) that inhibit protozoan feeding by inducing cell lysis. These specific chemical defenses are prevalent in the tropical aquatic and soil bacterium *Chromobacterium violaceum* and other marine bacteria, such as *Janthinobacterium lividum* and *Pseudoalteromonas luteoviolacea* (24).

The protozoan community in marine ecosystems is cosmopolitan and follows a succession pattern depending on the nature of the grazer. For example, early biofilm colonizers (or generalists, including flagellates and ciliates) are highly motile, allowing fast surface feeding, while intermediate late colonizers (some ciliates and amoebas) are classified as specialists and are abundant in mature biofilms (25, 26). Therefore, the aim of our study was to investigate whether *V. fischeri* biofilms are resistant to protozoan grazing, and if any differences exist in predator avoidance between various free-living and symbiotic strains. We tested symbiotic strains isolated from the light organ of two different squid genera, *Euprymna* (Indo-West Pacific) and *Sepiolo* (Mediterranean), along with free-living strains isolated directly from seawater. Three different protozoan predators among the 20 most commonly reported species of predators were chosen for these studies (20, 25) and included two early-feeding flagellates, *Neobodo designis* and

Received 4 September 2012 Accepted 26 October 2012

Published ahead of print 9 November 2012

Address correspondence to Michele K. Nishiguchi, nish@nmsu.edu.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.02710-12

TABLE 1 *Vibrio fischeri* strains used in this study

Strain	Host	Location
WH1	Free living	USA (Woods Hole, MA)
MDR7	Free living	USA (Marina del Rey, CA)
CB37	Free living	Australia (Coogee Bay, Sydney, NSW)
CB21	Free living	Australia (Coogee Bay, Sydney, NSW)
CHB8	Free living	Australia (Chowder Bay, Sydney, NSW)
CHB12	Free living	Australia (Chowder Bay, Sydney, NSW)
CHB30	Free living	Australia (Chowder Bay, Sydney, NSW)
BSM40	Free living	France (Banyuls sur mer)
BSM46	Free living	France (Banyuls sur mer)
BSM50	Free living	France (Banyuls sur mer)
PP3	Free living	USA (Kaneohe Bay, O'ahu, HI)
PP42	Free living	USA (Kaneohe Bay, O'ahu, HI)
VLS2	Free living	USA (Kaneohe Bay, O'ahu, HI)
SR5	<i>Sepiolo robusta</i>	France (Banyuls sur mer)
SL518	<i>Sepiolo lingulata</i>	France (Banyuls sur mer)
SA1G	<i>Sepiolo affinis</i>	France (Banyuls sur mer)
SA18	<i>Sepiolo affinis</i>	France (Banyuls sur mer)
SA25	<i>Sepiolo affinis</i>	France (Banyuls sur mer)
SA70	<i>Sepiolo affinis</i>	France (Banyuls sur mer)
SI66	<i>Sepiolo intermedia</i>	Italy (Bari)
SI1D	<i>Sepiolo intermedia</i>	France (Banyuls sur mer)
EM17	<i>Euprymna morsei</i>	Japan (Tokyo Bay)
ET101	<i>Euprymna tasmanica</i>	Australia (Townsville, QLD)
ETWW	<i>Euprymna tasmanica</i>	Australia (Woy Woy, NSW)
ETBB20	<i>Euprymna tasmanica</i>	Australia (Botany Bay, Sydney, NSW)
ETBB45	<i>Euprymna tasmanica</i>	Australia (Botany Bay, Sydney, NSW)
ETBB67	<i>Euprymna tasmanica</i>	Australia (Botany Bay, Sydney, NSW)
ETSB1	<i>Euprymna tasmanica</i>	Australia (Shark Bay, WA)
EB12	<i>Euprymna berryi</i>	Japan (Tosa Bay)
ES114	<i>Euprymna scolopes</i>	USA (Kaneohe Bay, O'ahu, HI)
ESP915	<i>Euprymna scolopes</i>	USA (Paiko, O'ahu, HI)
ESL5	<i>Euprymna scolopes</i>	USA (Kaneohe Bay, O'ahu, HI)
ESC9	<i>Euprymna scolopes</i>	USA (Kaneohe Bay, O'ahu, HI)

*Rhynchomonas nasuta*, and the late ciliate colonizer *Tetrahymena pyriformis*. This experimental setting allowed us to identify biofilm survival depending on (i) strain type, (ii) age of biofilm, (iii) protozoan colonizer, and (iv) protozoan feeding type, and these factors have implications for understanding how *V. fischeri* survives, persists, and diversifies in the presence of grazers.

## MATERIALS AND METHODS

**Bacterial and protozoan strains used in this study.** Bacterial strains used in this study are listed in Table 1. *V. fischeri* strains were grown on Luria-Bertani high-salt agar (LBS; with a per liter composition of 10 g tryptone, 5 g yeast extract, 20 g sodium chloride, 50 ml 1 M Tris [pH 7.5], 3.75 ml 80% glycerol, 15 g agar, and 950 ml distilled water) and incubated for 24 h at 28°C. *V. fischeri* strains were isolated from different squid light organs captured live (*Euprymna* or *Sepiolo*) or directly from seawater (27). All strains were subsequently subcultured in LBS liquid medium (no agar) and incubated with moderate shaking (200 rpm) for 18 h. The benthic flagellate grazers *R. nasuta* and *N. designis* were isolated from Chowder Bay at the Sydney Institute for Marine Science (SIMS), New South Wales, Australia, treated with an antibiotic cocktail (kanamycin, gentamicin, streptomycin, ampicillin, and trombamycin at 150 µg/ml each), and diluted through many generations (~15) to remove the natural bacterial community (14). *R. nasuta* and *N. designis* were maintained axenically in 0.5× nine salts solution medium (NSS; with a per liter composition of 8.8 g NaCl, 0.735 g Na<sub>2</sub>SO<sub>4</sub>, 0.04 g NaHCO<sub>3</sub>, 0.125 g KCl, 0.02 g KBr, 0.935 g MgCl<sub>2</sub> · 6H<sub>2</sub>O, 0.205 g CaCl<sub>2</sub> · 2H<sub>2</sub>O, 0.004 g SrCl<sub>2</sub> · 6H<sub>2</sub>O, 0.004 g H<sub>3</sub>BO<sub>3</sub>). Additional cultures were supplemented with heat-killed *P.*

*aeruginosa*, which served as prey. The ciliate *T. pyriformis* (CCAP 1630/1W; Culture Collection of Algae and Protozoa, Windmere, United Kingdom) was maintained in PPY medium (with a per liter composition of 20 g proteose peptone and 2.5 g yeast extract). Cultures were incubated at room temperature (20 to 23°C) for 2 weeks.

**Grazing assays.** Grazing experiments were completed in 24-well microtiter plates as previously described (20) for both early and late biofilms. For the early setup, overnight cultures of all bacterial strains were incubated at a dilution of 10<sup>6</sup> cells per ml in a total of 1 ml of fresh Vääänen-NSS medium (VNSS; with a composition of 1 g peptone, 0.5 g yeast extract, 0.5 g dextrose, 0.01 g FeSO<sub>4</sub> · 7H<sub>2</sub>O, 0.01 g Na<sub>2</sub>HPO<sub>4</sub>, mixed with 1 liter of NSS medium) (14), which allowed growth of both bacteria and protozoa. As a negative control, sterile uninoculated VNSS medium was used. Strains were grown for 6 h (to let the early biofilm to become established) at room temperature (28°C) in the 24-well microtiter plates, and subsequently the planktonic population was removed and replaced with 1 ml of fresh VNSS medium that contained overnight cultures of either *R. nasuta* or *N. designis* (early grazers) at an abundance of 10<sup>2</sup> cells per ml. In the late biofilm setup, overnight cultures of all bacterial strains were inoculated as described for the early biofilm setup, but with the bacterial biofilms forming for 24 h prior to addition of the late biofilm grazer *T. pyriformis*. After formation of mature (late) biofilms, the planktonic suspension was removed and *Tetrahymena* was added at a concentration of 10<sup>2</sup> cells per ml. Plates were incubated for 24 h at 20°C with shaking at 50 rpm. Additionally, as a positive control, bacterial cultures were inoculated in parallel (10<sup>6</sup> cells per ml in 1 ml of VNSS medium) and incubated without protozoan grazers for 6 h (early biofilms) or 24 h (late biofilms) with shaking (50 rpm) at 20°C. For each experiment, all samples were inoculated in triplicate (3 wells), and assays were repeated three times (3 independent studies), for a total of 9 wells per bacterial strain.

**Enumeration of protozoans and quantification of biofilm formation.** Numbers of grazers and grazer growth rates were calculated from direct cell counts using light microscopy. The number of grazers was calculated when 5 µl of supernatant was fixed with acid Lugol's solution (5%, final concentration) and enumerated using direct microscopy. Bacterial biofilms were measured using a colorimetric assay, where the supernatant from each plate was removed and wells were washed three times with 1 ml of sterile VNSS medium. One milliliter of 0.2% crystal violet solution was added to each well, and the mixture was incubated for 30 min at room temperature. After incubation, the crystal violet solution was removed and plates were washed five times with distilled water and dried. One milliliter of ethanol (95%) was added, and plates were incubated for 30 min to allow the dye to solubilize. The contents of each well (including negative controls) were transferred to a new plate, and the optical density was measured at 562 nm using a plate reader (Bio-Tek FLX 800; MTX Lab systems Inc., VA). The optical density observed is directly proportional to the amount of biofilm formed, and values were corrected by blank readings (uninoculated VNSS wells). For biofilm biomass quantifications, pairwise comparisons were performed using two-factor analysis of variance and Tukey's *post hoc* comparisons in order to test for significant differences between treatments (grazing versus nongrazing). Three plates were used for evaluation of statistical significance (3 independent studies).

**Scanning electron microscopy.** To observe toxic effects or morphological changes in the protozoan cultures, light microscopy and scanning electron microscopy (SEM) were performed. Overnight cultures of all strains were reinoculated in triplicate in glass test tubes containing 5 ml of VNSS with an immersed sterile coverslip. Early (2 sets of tubes) or late (1 set of tubes) biofilms were allowed to form. After 6 or 24 h of incubation, grazers were added (either *Rhynchomonas* or *Neobodo* for the early biofilms and *Tetrahymena* for the late biofilms) and incubated at room temperature for 24 h. After incubation, coverslips were washed with sterile VNSS, fixed with a 0.5% solution of glutaraldehyde, and gold coated for SEM with a Hitachi S34000 SEM apparatus (Hitachi, Schaumburg, IL) as previously described (19).




FIG 1 Quantification of biofilm biomass of free-living and symbiotic (*Euprymna* and *Sepiola*) *V. fischeri* strains before grazing (gray bars) and after grazing (white bars) on early (*R. nautae* and *N. designis*) and late (*T. pyriformis*) biofilms. The optical densities were measured after solubilization of crystal violet, with the abundance of biofilm considered directly proportional to the optical density reading. Error bars represent the standard deviations. \*,  $P < 0.05$  for the difference between grazed and nongrazed bacteria; \*\*,  $P < 0.01$ . The composite data are from 13 free-living strains, 8 *Sepiola* strains, and 12 *Euprymna* strains (see Table 1).

## RESULTS

**Resistance to grazing of early and late biofilms.** To determine potential inhibitory effects of biofilms (early and late) against three different protozoans with contrasting feeding modes, we compared the quantity of bacterial biofilm that was formed with and without exposure to each predator (Fig. 1). For early symbiotic strains, a significant reduction ( $P < 0.001$ ) in biofilm biomass was observed after addition of the ciliated predator *T. pyriformis*, whereas biofilms formed by free-living strains (Table 1; Fig. 1) did not show any significant differences from nongrazed biofilms. In late biofilms, the raptorial feeder *R. nasuta* had no effect on the biomass of any of the strains examined; however, the flagellate *N. designis* was able to reduce the biofilm biomass of all strains by more than 30% ( $P < 0.05$ ).

**Biofilms cause a decrease in predator number.** In this study, we hypothesized that a reduction in the number of grazers would occur if the biofilm biomass did not decrease compared to cells that were consumed. Our experiments demonstrated that anti-protozoal effects occurred in both early and late biofilms and that variations in toxicity exist between the two early biofilm flagellates, consistent with other protists that have similar feeding modes (28, 29).

Protist abundance decreased in every case that the biofilms persisted (Fig. 2). Growth rates of *T. pyriformis* (ciliate, late grazer) were significantly reduced when exposed to biofilms formed by free-living strains (isolated from France, Australia, and the United States) (Table 1; Fig. 1), whereas high growth rates were observed when symbiotic biofilms were grazed. The same pattern was observed with *N. designis* (flagellate, early grazer), where numbers increased after grazing of early biofilms formed by both symbiotic and free-living strains. Interestingly, *R. nasuta* (flagellate, early grazer) numbers decreased when grazing on both symbiotic and free-living early biofilms.

**Microscopy studies.** Biofilms that were preformed on a glass coverslip were exposed to all grazers for 24 h and then examined

via SEM in order to visualize possible toxic effects after grazing. *Vibrio* strains that were successfully grazed by the free-swimming filter feeder *T. pyriformis* (including most of the symbiotic strains) were found on glass coverslips with *T. pyriformis* (Fig. 3). The small amount of glass-associated biofilm was distinguishable from the larger ciliated *T. pyriformis* (late grazer). Additionally, evidence of grazing activity was present and included signs of active feeding by a large number of ciliated protozoa (Fig. 3A and B). Discernible changes in grazer morphology were observed in *T. pyriformis* after grazing on free-living strains (Fig. 4). Two different phenotypes in protozoan structure were observed, including grazing morphology (Fig. 4A and B) and cell lysis (Fig. 4C and D). Late biofilms (24 h) were formed at the air-liquid interface of




FIG 2 Protozoan growth rates after grazing on early and late biofilms. Negative numbers represent mortality, and positive numbers represent growth. Different bar patterns represent different groups of *V. fischeri* isolates (free-living or symbiotic with *Euprymna* or *Sepiola*). Error bars are the standard deviations for each treatment. Each protozoan group was significantly different from the others ( $P < 0.05$ ), and within the *Tetrahymena* group there was a significant difference in growth rate when free-living strains were grazed ( $P < 0.05$ ).




FIG 3 SEM images of *T. pyriformis* grazing dynamics. (A) *T. pyriformis* cells on a mature *V. fischeri* biofilm. (B) Magnification of *T. pyriformis* and the biofilm. (C) *T. pyriformis* grazing through a filter feeding strategy.

coverslips. Immediately after biofilm formation, *Tetrahymena* was added to the cultures, and after 24 h coverslips were analyzed using SEM. We observed two different coverslips per strain (for the protozoa *Tetrahymena*), and 60% of them exhibited a phenotype change for *Tetrahymena*, with protozoan lysis being the most common phenotype (after grazing biofilms formed by free-living strains).

## DISCUSSION

Vibrios are a ubiquitous diverse group of heterotrophic bacteria that are found in oceans, estuaries, and marine sediments worldwide. The diversity and dynamics of cooccurring populations has frequently

been linked to environmental changes, including fluctuations in temperature, salinity, ocean hydrodynamics, and nutrient composition (10, 30). Therefore, environmental fluctuations lead to considerable bacterial microdiversity and evolution, including evolution of a wide variety of pathogenic and facultative mutualistic strains. The latter include the bioluminescent bacterium *V. fischeri*, which exists in a free-living stage (seawater) or as a mutualist of sepiolid squids and monocentrid fishes (31–33).

Understanding the influences of abiotic factors on *V. fischeri* biofilm populations has become a central theme of our research (10, 27); however, success in the environment is also dictated by the ability of the biofilm to tolerate natural protozoan consumers.




FIG 4 SEM images of the antiprotozoan effects on *T. pyriformis*. (A and B) Morphological changes in the grazer and the bacterial growth around the predator. (C and D) Cell lysis of the grazer. Arrows in panels A and C point to the corresponding grazers that are shown in panels B and D.

Recent studies have emphasized the effects of protozoan grazers (17, 23, 34, 35), particularly in *Vibrio* communities, such as *V. cholerae* (12, 14), but not in communities of other species, including mutualists, such as *V. fischeri* and *V. logei*. In this study, we investigated the effects of common bacterivores *T. pyriformis* (ciliate, late grazer), *N. designis* (flagellate, early grazer), and *R. nasuta* (flagellate, early grazer) on 33 *V. fischeri* isolates from different origins, including seawater and two squid genera: *Euprymna*, found in Indo-West Pacific waters, and *Sepiolo*, found in the Mediterranean Sea. Symbiotic strains were chosen based on their similar growth and infection capabilities (all symbiotic strains could infect juvenile sepiolid squids) despite being from different host squids and geographical locations, whereas free-living strains were chosen based on their inability to infect juvenile squids yet their being found ubiquitously in the ocean.

Our results indicated that biofilms may protect *V. fischeri* from predation. For example, when the generalist ciliate *T. pyriformis* (late grazer) was added to late biofilms, a selective resistance to grazing was exhibited by free-living strains only and not symbiotic strains. Earlier work showed that free-living strains of *V. fischeri* are able to grow under a wider variety of conditions than their symbiotic relatives (10). This demonstrated that free-living strains have better adaptive mechanisms, which may be the result of genetic changes responsible for shifts in bacterial phenotypes (formation of biofilms) (2, 3).

It has been reported that biofilms of different species of environmental isolates secrete antiprotozoan factors, such as violacein (24) and others of an unknown nature (15, 20). These components may be present in *V. fischeri* biofilms, but they have not been identified thus far. *T. pyriformis* is a ciliated protozoa that grazes mature biofilms, similar to the ones found in the environment. Conversely, biofilm-like structures formed in the squid's light organ are expelled daily due to the diurnal cycle of the squid. This cycle allows exponential growth of *V. fischeri* followed by an expulsion of 99% of the symbionts into seawater every day at dawn (33, 36, 37). Thus, diel expulsion of a large number ( $10^{14}$ ) of bacteria possibly disrupts the preformed biofilm and therefore does not select for production of antiprotozoan compounds. Moreover, these strains may be exposed less frequently to predation and therefore are less capable of responding to predators that are only found in the aquatic environment, as in the case of the protozoan grazers used in this study.

Early biofilms were exposed to the flagellates *N. designis* and *R. nasuta*, which have different feeding modes and are considerably smaller in size than *T. pyriformis* (200 times bigger than *V. fischeri*). Our results revealed that early biofilms exposed to *N. designis* were consumed, resulting in a pronounced increase in flagellate numbers, which was opposite to observations for the flagellate *R. nasuta*. These contrasting results indicate that flagellate size is not necessarily correlated to grazing dynamics. For example, *N. designis* is a direct interception feeder, creating currents that carry suspended or loosely attached bacteria toward the mouth (28). *R. nasuta* is a raptorial feeder that grasps its prey with a proboscis-like structure (26). Another major difference between these two flagellates is that *R. nasuta* has a lower grazing (ingestion) success in sparsely populated biofilms due to its slower moving speed, and it is more successful in waters with higher prey densities (38). These different feeding modes may be crucial to the grazing success regarding *V. fischeri* biofilms. Symbiotic *V. fischeri* strains also exhibit resistance to *R. nasuta* predation (Fig. 1), indi-

cating that some defensive mechanism still exists within biofilms formed by mutualistic strains. These defenses may be coopted to avoid the squid host innate immune response, which is mostly comprised of hemocytes and macrophages (39, 40). Future studies will focus on whether receptor-mediated phagocytosis in protozoa affects immune response evasion through biofilm formation.

In this study, protozoan abundance was reduced after grazing, and antiprotozoan effects on *T. pyriformis* (ciliate, late grazer) were observed (Fig. 2). Morphological changes of *T. pyriformis* occurred with bacteria possibly colonizing the protozoan predator, resulting in subsequent lysis of *T. pyriformis* cells. These two effects may be the result of inactivation of the predator or synthesis of lytic compounds that are released before or after ingestion of *V. fischeri* cells.

Quorum sensing controls biofilm formation (41) and has been described as one possible mechanism for antiprotozoal activity (20). In the case of *V. fischeri*, quorum sensing is under the control of the transcriptional regulator LuxR-LuxI (8). In addition to genes involved in light production, the LuxR regulon activates another number of genes involved in synthesis of efflux proteins, transporters, permeases, and proteases (8). The role of these quorum-sensing proteins in antipredator activity is unknown; however, they may contribute to the competitive fitness of biofilms under grazing pressure. Future studies will help determine the genetic factors responsible for protozoan death and may provide important clues of how survival of *Vibrio* biofilms is linked to quorum-sensing mechanisms.

The nature of the protozoan predator and its feeding characteristics may influence the impact of grazing on *V. fischeri* biofilms. Late/early and generalist/specialist protists have distinct grazing preferences, which are related to the type of *V. fischeri* strain examined (free living or symbiotic). Here we showed that *V. fischeri* biofilms have the ability to quickly adapt to grazing pressure, possibly by releasing antigrazing compounds and products that negatively influence protist survival. The observation that free-living strains are considerably more resistant to grazing pressure than symbiotic strains suggests that host selection may compromise the fitness of *V. fischeri* strains that are more amenable to a stable (and predatorless) environment than the external environment. Future research will address various adaptive mechanisms of bacteria-protist interactions that share fundamental processes with host immune responses.

## ACKNOWLEDGMENTS

We thank P. Cooke for helping with the SEM images and two anonymous reviewers who provided helpful suggestions, ideas, and future studies.

This project was partially supported by The Centre for Marine Bio-Innovation and the Australian Research Council, grant DPDP1096481, and NSF IOS-0744498, NIH NIAID 1SC1AI081659-01, and NIH ARRA-3SC1AI081659-02S1 to M.K.N. A.C.-D. was supported by RISE (NIH NIGMS R25GM061222) and NASA (NMSGC).

We have no conflicts of interests to declare.

## REFERENCES

1. Watnick P, Kotler R. 2000. Biofilm, city of microbes. *J. Bacteriol.* 182: 2675–2679.
2. Yildiz FH, Visick KL. 2008. *Vibrio* biofilms: so much the same yet so different. *Trends Microbiol.* 17:109–118.
3. Chavez-Dozal A, Nishiguchi MK. 2011. Variation in biofilm formation among symbiotic and free-living strains of *Vibrio fischeri*. *J. Basic Microbiol.* 51:452S–458S.

4. Costerton JW, Stewart PS, Greenberg EP. 1999. Bacterial biofilms: a common cause of persistent infections. *Science* 284:1318–1322.
5. Mah TF, O'Toole GA. 2001. Mechanisms of biofilm resistance to antimicrobial agents. *Trends Microbiol.* 9:34–39.
6. Espeland EM, Wetzel RG. 2001. Complexation, stabilization, and UV photolysis of extracellular and surface-bound glucosidase and alkaline phosphatase: implications for biofilms microbiota. *Microb. Ecol.* 42:572–585.
7. Dalesman S, Karnik V, Lukowiak K. 2011. Sensory mediation of memory blocking stressors in the pond snail *Lymnaea stagnalis*. *J. Exp. Biol.* 214:2528–2533.
8. Antunes LCM, Schaefer AL, Ferreira RBR, Qin N, Stevens AM, Ruby EG, Greenberg EP. 2007. Transcriptome analysis of the *Vibrio fischeri* LuxR-LuxI regulon. *J. Bacteriol.* 189:8387–8391.
9. Ariyakumar DS, Nishiguchi MK. 2009. Characterization of two host specific genes, mannose sensitive hemagglutinin (*mshA*) and uridyl phosphate dehydrogenase (UDPDH) that are involved in the *Vibrio fischeri*-*Euprymna tasmanica* mutualism. *FEMS Microbiol. Lett.* 299:65–73.
10. Soto W, Gutierrez J, Remmaga MD, Nishiguchi MK. 2009. Salinity and temperature effects on physiological responses of *Vibrio fischeri* from diverse ecological niches. *Microb. Ecol.* 57:140–150.
11. Kadouri D, O'Toole GA. 2005. Susceptibility of biofilms to *Bdellovibrio bacteriovorus* attack. *Appl. Environ. Microbiol.* 71:4044–4051.
12. Matz C, Kjelleberg S. 2005. Off the hook: how bacteria survive protozoan grazing. *Trends Microbiol.* 13:302–307.
13. Adiba S, Nizak C, Van Baalen M, Denamur E, Depaulis F. 2010. From grazing resistance to pathogenesis: the coincidental evolution of virulence factors. *PLoS One* 5:e11882. doi:10.1371/journal.pone.0011882.
14. Erken M, Weitere K, Kjelleberg S, McDougald SD. 2011. In situ grazing resistance of *Vibrio cholerae* in the marine environment. *FEMS Microbiol. Ecol.* 76:504–512.
15. Matz C, Jurgens K. 2001. Effects of hydrophobic and electrostatic cell surface properties of bacteria on feeding of heterotrophic nanoflagellates. *Appl. Environ. Microbiol.* 67:814–820.
16. Hazlett BA. 2003. Predator recognition and learned irrelevance in the crayfish *Orconectes virilis*. *Ethology* 109:765–780.
17. Miki T, Jacquet S. 2008. Complex interactions in the microbial world: underexplored key links between viruses, bacteria and protozoan grazers in aquatic environments. *Aquat. Microb. Ecol.* 51:195–208.
18. Weitere M, Bergfeld T, Rice SA, Matz C, Kjelleberg S. 2005. Grazing resistance of *Pseudomonas aeruginosa* biofilms depends on type of protective mechanism, developmental stage and protozoan feeding mode. *Environ. Microbiol.* 7:1593–1601.
19. Greiner LL, Edwards JL, Shao J, Rabinak C, Entz D, Apicella MA. 2005. Biofilm formation by *Neisseria gonorrhoeae*. *Infect. Immun.* 73:1964–1970.
20. Matz C, McDougald D, Moreno AM, Yung PY, Yildiz FH, Kjelleberg S. 2005. Biofilm formation and phenotype variation enhance predation-driven persistence of *Vibrio cholerae*. *Proc. Natl. Acad. Sci. U. S. A.* 102:16819–16824.
21. Queck SY, Weitere M, Moreno AM, Rice SA, Kjelleberg S. 2006. The role of quorum sensing mediated developmental traits in the resistance of *Serratia marcescens* biofilms against protozoan grazing. *Environ. Microbiol.* 8:1017–1025.
22. Turovskiy Y, Rosenberg L, Chikindas M. 2007. Autoinducer-2-mediated quorum sensing is not involved in *Listeria monocytogenes* adaptive responses to the food preservatives lactic acid and nisin. *J. Food Saf.* 27:386–399.
23. Koh KS, Matz C, Tan CH, Le HL, Rice SA, Marshall DJ, Steinberg PD, Kjelleberg S. 2012. Minimal increase in genetic diversity enhances predation resistance. *Mol. Ecol.* 21:1741–1749.
24. Matz C, Webb JS, Schupp PJ, Phang SY, Penesyan A, Egan S, Steinberg P, Kjelleberg S. 2008. Marine biofilm bacteria evade eukaryotic predation by targeted chemical defense. *PLoS One* 3:e2744. doi:10.1371/journal.pone.0002744.
25. Arndt H, Schmidt-Denter K, Auer B, Weitere M. 2003. Protozoans and biofilms, p 173–189. In Krumbein WE, Paterson DM, Zavarzin GA (ed), *Fossil and recent biofilms*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
26. Bohme A, Risse-Buhl U, Kusel K. 2009. Protists with different feeding modes change biofilm morphology. *FEMS Microbiol. Ecol.* 69:158–169.
27. Zamborsky DJ, Nishiguchi MK. 2011. Phylogeographical patterns among sympatric populations of sepioid squids and their *Vibrio* symbionts in the Mediterranean Sea. *Appl. Environ. Microbiol.* 77:642–649.
28. Artolozaga I, Ayo B, Latatu A, Azua I, Unanue M, Iriberrri J. 2000. Spatial distribution of protists in the presence of macroaggregates in a marine system. *FEMS Microbiol. Ecol.* 33:191–196.
29. Boenigk J, Arndt H. 2002. Bacterivory by heterotrophic flagellates: community structure and feeding strategies. *Antonie Van Leeuwenhoek* 81:465–480.
30. Thompson JR, Randa MA, Marcelino LA, Tomita-Mitchell A, Lim E, Polz MF. 2004. Diversity and dynamics of a north atlantic coastal *Vibrio* community. *Appl. Environ. Microbiol.* 70:4103–4110.
31. Nishiguchi MK, Lopez JE, Boletzky SV. 2004. Enlightenment of old ideas from new investigations: the evolution of bacteriogenic light organs in squids. *Evol. Dev.* 6:41–49.
32. Nyholm SV, Nishiguchi MK. 2008. The evolutionary ecology of a sepioid squid-*Vibrio* association: from cell to environment. *Vie Milieu Paris* 58:175–184.
33. Ruby EG, McFall-Ngai MJ. 1999. Oxygen-utilizing reactions and symbiotic colonization of the squid light organ by *Vibrio fischeri*. *Trends Microbiol.* 7:414–420.
34. Jurgens K, Matz C. 2002. Predation as a shaping force for the phenotypic and genotypic composition of planktonic bacteria. *Antonie Van Leeuwenhoek* 81:413–434.
35. Kicklighter CE, Hay ME. 2006. Integrating prey defensive traits: contrasts of marine worm from temperate and tropical habitats. *Ecol. Monogr.* 76:195–215.
36. Ruby EG. 1996. Lessons from a cooperative, bacterial-animal association: the *Vibrio fischeri*-*Euprymna scolopes* light organ symbiosis. *Annu. Rev. Microbiol.* 50:591–624.
37. Ruby EG, Lee K. 1998. The *Vibrio fischeri*-*Euprymna scolopes* light organ association: current ecological paradigms. *Appl. Environ. Microbiol.* 64:805–812.
38. Erken M, Farrenschon N, Speckmann S, Arndt H, Weitere M. 2012. Quantification of individual flagellate-bacteria interactions within semi-natural biofilms. *Protist* 163:632–642.
39. Castillo MG, Goodson MS, McFall-Ngai M. 2009. Identification and molecular characterization of a complement C3 molecule in lophotrochozoan, the Hawaiian bobtail squid *Euprymna scolopes*. *Dev. Comp. Immunol.* 33:69–76.
40. McFall-Ngai MJ, Nyholm S, Castillo M. 2010. The role of the immune system in the initiation and persistence of the *Euprymna scolopes*-*Vibrio fischeri* symbiosis. *Semin. Immunol.* 22:48–53.
41. Hammer BK, Bassler BL. 2003. Quorum sensing controls biofilm formation in *Vibrio cholerae*. *Mol. Microbiol.* 50:101–104.