

Erratum: “Enhanced broadband near-IR luminescence and gain spectra of bismuth/erbium co-doped fiber by 830 and 980 nm dual pumping” [AIP Advances 7, 045012 (2017)]

Qiancheng Zhao, Yanhua Luo, Wenyu Wang, John Canning, and Gang-Ding Peng

Citation: [AIP Advances](#) **7**, 059901 (2017); doi: 10.1063/1.4983016

View online: <https://doi.org/10.1063/1.4983016>

View Table of Contents: <http://aip.scitation.org/toc/adv/7/5>

Published by the [American Institute of Physics](#)

Articles you may be interested in

[Enhanced broadband near-IR luminescence and gain spectra of bismuth/erbium co-doped fiber by 830 and 980 nm dual pumping](#)

[AIP Advances](#) **7**, 045012 (2017); 10.1063/1.4981903

[Erratum: “Effects of the partial substitution of Ni by Cr on the transport, magnetic, and magnetocaloric properties of \$\text{Ni}_{50}\text{Mn}_{37}\text{In}_{13}\$ ” \[AIP Advances 7, 056433 \(2017\)\]](#)

[AIP Advances](#) **7**, 059902 (2017); 10.1063/1.4983194

[Publisher’s Note: “Continuous tuning of two-section, single-mode terahertz quantum-cascade lasers by fiber-coupled, near-infrared illumination” \[AIP Advances 7, 055201 \(2017\)\]](#)

[AIP Advances](#) **7**, 059903 (2017); 10.1063/1.4984032

[Optimized radiation-hardened erbium doped fiber amplifiers for long space missions](#)

[Journal of Applied Physics](#) **121**, 163104 (2017); 10.1063/1.4981532

[Magnetic scanning gate microscopy of CoFeB lateral spin valve](#)

[AIP Advances](#) **7**, 056808 (2017); 10.1063/1.4977891

[Spatial resolution and maximum compensation factor of two-dimensional selective excitation pulses for MRI of objects containing conductive implants](#)

[AIP Advances](#) **7**, 056726 (2017); 10.1063/1.4977498

PHYSICS TODAY

WHITEPAPERS

MANAGER’S GUIDE

Accelerate R&D with
Multiphysics Simulation

READ NOW

PRESENTED BY

 **COMSOL**

Erratum: “Enhanced broadband near-IR luminescence and gain spectra of bismuth/erbium co-doped fiber by 830 and 980 nm dual pumping” [AIP Advances 7, 045012 (2017)]

Qiancheng Zhao,^{1,a} Yanhua Luo,^{1,2} Wenyu Wang,¹ John Canning,³ and Gang-Ding Peng¹

¹*Photonics and Optical Communication, School of Electrical Engineering and Telecommunications, University of New South Wales, Kensington, NSW 2052, Australia*

²*Key Lab of In-fiber Integrated Optics, Ministry Education of China, Harbin Engineering University, Harbin 150001, China*

³*Interdisciplinary Photonics Laboratories, School of Chemistry, University of Sydney, NSW 2006, Australia*

(Received 20 April 2017; accepted 20 April 2017; published online 1 May 2017)

[<http://dx.doi.org/10.1063/1.4983016>]

ACKNOWLEDGMENTS

This work was supported by NSFC (61520106014, 61405014, and 61377096), Fund of Key Lab of In-fiber Integrated Optics, Ministry Education of China, STCSM (SKLSFO2015-01 and 15220721500), Open Fund of SKLFPM (DHU) (LK1502), and Fund of IPOC (BUPT) (IPOC2016ZT07).

^aCorresponding author: Qiancheng.zhao@unsw.edu.au