



Genome Sequence of *Dehalobacter* sp. Strain TeCB1, Able To Respire Chlorinated Benzenes

Ricardo Alfán-Guzmán,^a Haluk Ertan,^{a,b} Mike Manefield,^a Matthew Lee^a

School of Biotechnology and Biomolecular Sciences, University of New South Wales, Sydney, Australia^a;
Department of Molecular Biology and Genetics, Istanbul University, Vezneciler, Turkey^b

ABSTRACT *Dehalobacter* sp. strain TeCB1 was isolated from groundwater contaminated with a mixture of organohalides and is able to respire 1,2,4,5-tetrachlorobenzene and 1,2,4-trichlorobenzene. Here, we report its 3.13-Mb draft genome sequence.

Chlorinated benzenes are a group of isomeric aromatic compounds extensively used in the production of pesticides, insecticides, heat transfer agents, dyes, and as additives for rubber products (1, 2). This group of compounds are common groundwater pollutants and probable carcinogens (3). Release of these compounds to the environment often occurs due to inadequate handling or disposal; e.g., 1,2,4,5-tetrachlorobenzene (TeCB) and 1,2,4-trichlorobenzene (TCB) have been found in crops, fish, groundwater, and air in the United States, Europe, and China. Both substances are not only probable carcinogens, but it has been shown that acute exposure may lead to kidney or liver damage in humans (4–6).

Detoxification of chlorinated benzenes can occur when they are used as terminal electron acceptors in microbial respiration. *Dehalococcoides mccartyi* strain CBDB1 is able to reductively dechlorinate hexachlorobenzene (HCB) to 1,3,5-TCB, 1,3-, and 1,4-dichlorobenzene (DCB) (7). In 2014, Nelson et al. (8) isolated two *Dehalobacter* strains and a highly enriched culture able to respire a variety of chlorinated benzenes. In this study, we present the genome sequence of *Dehalobacter* sp. strain TeCB1, which is able to respire 1,2,4,5-TeCB and 1,2,4-TCB, yielding 1,3- and 1,4-DCB. Strain TeCB1 was isolated from organohalide-contaminated groundwater in Sydney, Australia.

Genomic DNA from strain TeCB1 was extracted according to Murray et al. (9) and then sequenced using an Illumina HiSeq 2500 sequencer at Novogene Bioinformatics Technology (Beijing, China). A total of 2,341,965 high-quality 100-bp paired-end reads were generated (coverage 100×). *De novo* assembly was performed using SPAdes assembler version 3.6.1 standard pipeline (10), generating 67 contigs. Annotation of the assembled genome was conducted via the NCBI Prokaryotic Genome Annotation Pipeline (version 3.3). The *Dehalobacter* sp. strain TeCB1 complete genome is 3.13 Mb long, with N_{50} value of 105,922 bp, and comprises a G+C content of 44%; a total of 3,081 genes were discovered, in which 2,962 are protein-coding genes, seven were rRNA genes (i.e., three 5S, one 16S, and three 23S), and 50 were tRNA genes. A genome quality assessment tool (CheckM) was used to assess the quality and purity of the draft genome (11). It reported a completeness of 99.94% based on the finding of 418/420 lineage-specific marker genes (marker lineage *Clostridia*), contamination of 0.17%, and strain heterogeneity tested by the amino acid identity between multicopy genes of zero.

Although other strains able to respire chlorinated benzenes belonging to the genus *Dehalobacter* have been isolated, to date, only the draft genome sequence of strain TeCB1 is publicly available (<https://www.ncbi.nlm.nih.gov/nucleotide/1055168950>).

Received 12 December 2016 Accepted 16 December 2016 Published 23 February 2017

Citation Alfán-Guzmán R, Ertan H, Manefield M, Lee M. 2017. Genome sequence of *Dehalobacter* sp. strain TeCB1, able to respire chlorinated benzenes. *Genome Announc* 5:e01681-16. <https://doi.org/10.1128/genomeA.01681-16>.

Copyright © 2017 Alfán-Guzmán et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Matthew Lee, mattlee@unsw.edu.au.

Strain TeCB1's genome was compared to that of its closest relative, *Dehalobacter restrictus* PER-K23; among the many differences found, TeCB1 possesses a single 16S rRNA gene, while the PER-K23 genome contains four (12). Strain TeCB1 encodes 24 reductive dehalogenase (RdhA) homologs, one of them N-terminally truncated and five unique to this strain, together with a complete set of genes for *de novo* cobalamin and menaquinone biosynthesis and the Wood-Ljungdahl pathway. The genome also comprises the genes encoding various kinds of [NiFe]- and [FeFe]-hydrogenases, including Hup-type and bifurcating hydrogenases.

Accession number(s). This whole-genome shotgun project has been deposited at DDB/ENA/GenBank under the version number [MCHF00000000](https://www.ncbi.nlm.nih.gov/genbank/mchf/00000000).

ACKNOWLEDGMENTS

This research was supported by Consejo Nacional de Ciencia y Tecnología (CONACYT-Mexico) and the Australian Research Council (grant LP130100454).

REFERENCES

1. Fishbein LU. 2011. Potential industrial carcinogens and mutagens. Amsterdam: Elsevier Science.
2. Woo YT, Lai DY, Arcos JCU. 2013. Aliphatic and polyhalogenated carcinogens: structural bases and biological mechanisms. Chicago: Elsevier Science.
3. Field JA, Sierra-Alvarez R. 2008. Microbial degradation of chlorinated benzenes. *Biodegradation* 19:463–480. <https://doi.org/10.1007/s10532-007-9155-1>.
4. Zhang H, Wang Y, Sun C, Yu M, Gao Y, Wang T, Liu J, Jiang G. 2014. Levels and distributions of hexachlorobutadiene and three chlorobenzenes in biosolids from wastewater treatment plants and in soils within and surrounding a chemical plant in China. *Environ Sci Technol* 48:1525–1531. <https://doi.org/10.1021/es405171t>.
5. Wang MJ, McGrath SP, Jones KC. 1995. Chlorobenzenes in field soil with a history of multiple sewage sludge applications. *Environ Sci Technol* 29:356–362. <https://doi.org/10.1021/es00002a011>.
6. Dong WH, Zhang P, Lin XY, Zhang Y, Tabouré A. 2015. Natural attenuation of 1,2,4-trichlorobenzene in shallow aquifer at the Luhuaqiang's landfill site, Kaifeng, China. *Sci Total Environ* 505:216–222. <https://doi.org/10.1016/j.scitotenv.2014.10.002>.
7. Jayachandran G, Görisch H, Adrian L. 2003. Dehalorespiration with hexachlorobenzene and pentachlorobenzene by *Dehalococcoides* sp. strain CBDB1. *Arch Microbiol* 180:411–416. <https://doi.org/10.1007/s00203-003-0607-7>.
8. Nelson JL, Jiang J, Zinder SH. 2014. Dehalogenation of chlorobenzenes, dichlorotoluenes, and tetrachloroethene by three *Dehalobacter* spp. *Environ Sci Technol* 48:3776–3782. <https://doi.org/10.1021/es4044769>.
9. Murray AE, Preston CM, Massana R, Taylor LT, Blakis A, Wu K, DeLong EF. 1998. Seasonal and spatial variability of bacterial and archaeal assemblages in the coastal waters near Anvers Island, Antarctica. *Appl Environ Microbiol* 64:2585–2595.
10. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
11. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
12. Kruse T, Maillard J, Goodwin L, Woyke T, Teshima H, Bruce D, Detter C, Tapia R, Han C, Huntemann M, Wei CL, Han J, Chen A, Kyrpides N, Szeto E, Markowitz V, Ivanova N, Pagani I, Pati A, Pitluck S, Nolan M, Holliger C, Smidt H. 2013. Complete genome sequence of *Dehalobacter restrictus* PER-K23(T). *Stand Genomic Sci* 8:375–388. <https://doi.org/10.4056/sigs.3787426>.