



# Complete Sequences of Multiple-Drug Resistant IncHI2 ST3 Plasmids in *Escherichia coli* of Porcine Origin in Australia

Ethan R. Wyrsch<sup>1</sup>, Cameron J. Reid<sup>1</sup>, Matthew Z. DeMaere<sup>1</sup>, Michael Y. Liu<sup>1</sup>, Toni A. Chapman<sup>2</sup>, Piklu Roy Chowdhury<sup>1,2</sup> and Steven P. Djordjevic<sup>1\*</sup>

<sup>1</sup> The ithree institute, University of Technology Sydney, Sydney, NSW, Australia, <sup>2</sup> Elizabeth Macarthur Agricultural Institute, NSW Department of Primary Industries, Sydney, NSW, Australia

## OPEN ACCESS

### Edited by:

Ömer Akineden,  
University of Giessen, Germany

### Reviewed by:

Aurora García-Fernández,  
Istituto Superiore di Sanità (ISS), Italy  
Costas C. Papagiannitsis,  
University of Thessaly, Greece

### \*Correspondence:

Steven P. Djordjevic  
steven.djordjevic@uts.edu.au

### Specialty section:

This article was submitted to  
Agro-Food Safety,  
a section of the journal  
Frontiers in Sustainable Food Systems

Received: 14 January 2019

Accepted: 06 March 2019

Published: 26 March 2019

### Citation:

Wyrsch ER, Reid CJ, DeMaere MZ,  
Liu MY, Chapman TA, Roy  
Chowdhury P and Djordjevic SP  
(2019) Complete Sequences of  
Multiple-Drug Resistant IncHI2 ST3  
Plasmids in *Escherichia coli* of Porcine  
Origin in Australia.  
Front. Sustain. Food Syst. 3:18.  
doi: 10.3389/fsufs.2019.00018

IncHI2 ST3 plasmids are known carriers of multiple antimicrobial resistance genes. Complete plasmid sequences from multiple drug resistant *Escherichia coli* circulating in Australian swine is however limited. Here we sequenced two related IncHI2 ST3 plasmids, pSDE-SvHI2, and pSDC-F2\_12BHI2, from phylogenetically unrelated multiple-drug resistant *Escherichia coli* strains SvETEC (CC23:O157:H19) and F2\_12B (ST93:O7:H4) from geographically disparate pig production operations in New South Wales, Australia. Unicycler was used to co-assemble short read (Illumina) and long read (PacBio SMRT) nucleotide sequence data. The plasmids encoded three drug-resistance loci, two of which carried class 1 integrons. One integron, hosting *drfA12-orfF-aadA2*, was within a hybrid Tn1721/Tn21, with the second residing within a copper/silver resistance transposon, comprising part of an atypical *sul3*-associated structure. The third resistance locus was flanked by IS15DI and encoded neomycin resistance (*neoR*). An *oqx*-encoding transposon (quinolone resistance), similar in structure to Tn6010, was identified only in pSDC-F2\_12BHI2. Both plasmids showed high sequence identity to plasmid pSTM6-275, recently described in *Salmonella enterica* serotype 1,4,[5],12:i:- that has risen to prominence and become endemic in Australia. IncHI2 ST3 plasmids circulating in commensal and pathogenic *E. coli* from Australian swine belong to a lineage of plasmids often in association with *sul3* and host multiple complex antibiotic and metal resistance structures, formed in part by IS26.

**Keywords:** *Escherichia coli*, antimicrobial resistance (AMR), plasmid, genomics, epidemiology

## INTRODUCTION

In Australia, restrictions on live animal imports, geographic isolation, and sound antibiotic stewardship have limited the incorporation and spread of genes encoding resistance to antibiotics used to treat serious human infections among *Enterobacteriaceae* circulating in food animals (Turner, 2011; Abraham et al., 2015; Reid et al., 2017; Kidsley et al., 2018). Australian porcine *Escherichia coli* are, albeit at low frequency, known to carry *bla*<sub>CMY-2</sub>, *bla*<sub>CTX-M-14</sub>, and *bla*<sub>CTX-M-9</sub>, possibly because of the use of ceftiofur as an off-label, last line antibiotic to treat serious disease (Abraham et al., 2015). First generation antibiotics; commonly tetracyclines,

penicillins, and sulphonamides are otherwise used for the therapeutic treatment of bacterial infections in swine (Jordan et al., 2009). Both pathogenic and commensal *E. coli* sourced from the feces of pig carry class 1 integrons and are often multiple drug resistant (MDR) (Abraham et al., 2015; Wyrsh et al., 2015; Reid et al., 2017). Notably, the insertion element IS26 has infiltrated the genomes of commensal *E. coli* of porcine origin, where it has played a role in altering the genetic context of clinical class 1 integrons and facilitated the acquisition of further resistances (Reid et al., 2017). Furthermore, it is important to identify and characterize the genetic vehicles that can carry class 1 integrons and genes encoding resistance to first generation antibiotics which are widespread in commensal *E. coli* populations in Australian pigs (Abraham et al., 2015; Reid et al., 2017; Kidsley et al., 2018) because they are likely to acquire genes encoding resistance to clinically relevant antibiotics when resident in commensal enterobacterial populations in the gut of humans and companion animals.

Plasmids of the incompatibility group HI2 (IncHI2) have been linked with the carriage of tellurium resistance, plus a broad range of antimicrobial resistance genes, including *bla*<sub>IMP</sub>, *bla*<sub>NDM</sub>, and *bla*<sub>VIM</sub> (carbapenemase resistance) (Abraham et al., 2016; Dolejska et al., 2016; Falgenhauer et al., 2017), *bla*<sub>CTX-M</sub> (Garcia Fernandez et al., 2007), *mcr-1* (plasmid-mediated colistin resistance) (Gilrane et al., 2017; Li et al., 2017; Zheng et al., 2017), *oqxAB* and *aac(6)-Ib-cr* (plasmid-mediated quinolone resistance) (Fang et al., 2016), and resistance genes effective against first generation antibiotics and disinfectants (Cain et al., 2010; Cain and Hall, 2012; Abraham et al., 2016; Dolejska et al., 2016; Gilrane et al., 2017). Notably, clinically-important resistance genes have been associated with IncHI2 plasmids that also encode resistance to copper, zinc and, arsenic residues (Fang et al., 2016). In Australia, IncHI2 plasmids carrying a diverse range of antimicrobial resistance genes have been found in MDR *E. coli* (Dolejska et al., 2016; Saputra et al., 2017), *Salmonella enterica* serovar Typhimurium (Cain et al., 2010; Cain and Hall, 2012; Billman-Jacobe et al., 2018), *Enterobacter cloacae* (Sidjabat et al., 2015), and *Enterobacter hormaechei* subsp. *Oharae* (Monahan et al., 2019).

The insertion sequence IS26 of the IS6 family plays a significant role in the capture, assembly and mobilization of drug resistance genes found on plasmids (Dionisi et al., 2009; Cain et al., 2010; Shahid, 2010; Venturini et al., 2010, 2013; Lai et al., 2013; Abbo and Hooton, 2014; Chavda et al., 2015; Reid et al., 2015; Garcia et al., 2016) and in the chromosome (Roy Chowdhury et al., 2015, 2018). IS26 may promote plasmid stability and persistence by mediating deletions of plasmid backbone sequence, the expression of which incurs a burden to the host (Porse et al., 2016). IS26 can also facilitate the generation of hybrid virulence/resistance plasmids, formed by co-integration of plasmids separately carrying resistance and virulence gene cargo (Mangat et al., 2017; Wong et al., 2017). Although IS26 has no target site specificity (Harmer et al., 2014), it is often observed localizing next to or within class 1 integrons in multiple drug resistant *E. coli* recovered from the feces of both healthy pigs, poultry and cattle with gastrointestinal disease (Dawes et al., 2010; Reid et al., 2017). IS26 can shape

the structure of class 1 integrons by facilitating the addition of foreign DNA flanked by IS26, and by generating inversion and deletions of sequence within complex resistance structures. Consequently, many class 1 integrons have lost genes that reside within the often observed 3'-conserved sequence (3'-CS), particularly the sulphonamide resistance gene *sul1* (Reid et al., 2017). However, sulphonamide resistance persists globally (Grape et al., 2003; Bean et al., 2005; Suhartono et al., 2017), despite restrictions on its use (Enne et al., 2001), with *sul2* (Bean et al., 2005) and *sul3* genes (Grape et al., 2003; Perreten and Boerlin, 2003; Zhou et al., 2014; Reid et al., 2017) found frequently in close association with class 1 integrons. A *sul4* gene associated with a chromosomal locus containing the folate synthesis gene *folK* and a copy of ISCR20 has recently been described (Razavi et al., 2017). Emerging trends show that genes encoding resistance to last line drugs, such as *mcr-1*, *bla*<sub>CTX-M</sub>, and carbapenemases are captured on plasmids already carrying genes encoding resistances to first generation antibiotics (Alonso et al., 2017; Botts et al., 2017; Delannoy et al., 2017; Poirel et al., 2017). Any one of a number of selection pressures including heavy metals and biocides may be sufficient to then facilitate the persistence and spread of multiple drug resistance plasmids (Argudín et al., 2019). This is further compounded by reports that agrichemicals can alter selection for drug resistant bacteria (Kurenbach et al., 2015, 2018). The potential for complex resistance structures to persist under multiple different selective pressures and across diverse environments underpin the importance of adapting a One Health approach to antimicrobial resistance gene surveillance in humans, food and companion animals, agriculture, effluent (municipal, hospital, and agricultural), and the environments impacted by effluent from diverse sources (Djordjevic et al., 2013; Wyrsh et al., 2016; Huang et al., 2017; Mir et al., 2018).

Here we used a combination of Illumina and PacBio SMRT sequencing to completely close two IncHI2:ST3 MDR plasmids from *E. coli* recovered from the feces of Australian swine. One of these, pSDE-SvHI2 is from a severe ETEC/ExPEC pathogen (*E. coli* O157 SvETEC) (Wyrsh et al., 2015), and the other, pSDC-F2\_12BHI2, is from *Escherichia coli* F2\_12B, a commensal *E. coli* ST93 (Reid et al., 2017). Phylogenetic and genomic comparisons were undertaken between these two plasmid sequences and pSTM6-275, recently isolated from a monophasic variant of *Salmonella enterica* in Australian pigs (Dyall-Smith et al., 2017), as well as all available IncHI2 ST3 plasmids on GenBank.

## MATERIALS AND METHODS

### Strains and DNA Preparation

Plasmid pSDE-SvHI2 was resolved from *Escherichia coli* O157 SvETEC (CC23:O157:H19), a severe pathogen that caused intractable disease within an Australian commercial piggery in 2007 (Wyrsh et al., 2015). Likewise, pSDC-F2\_12BHI2 was resolved from *Escherichia coli* F2\_12B, a commensal ST93:O7:H4 strain isolated from a rectal swab of a commercially raised pig, also sourced in 2007 (Reid et al., 2017). Full descriptions of these strains plus information on Illumina

short read sequencing and assembly have been published previously (Wyrsh et al., 2015; Reid et al., 2017). Un-sheared genomic DNA suitable for SMRT sequencing was prepared from mid-log phase sub-cultures of strains grown overnight in LB broth using a gentle phenol-chloroform extraction protocol. Genomic DNA samples were checked for appropriate DNA concentrations and integrity using a Qubit dsDNA HS (high sensitivity, 0.2 to 100 ng) Assay Kit on Qubit 2.0 fluorometer (Life Technologies) and for shearing by agarose gel electrophoresis.

## Genome Sequencing and Assembly

Long read sequencing was performed by the Ramaciotti Center for Genomics using a Pacific Biosciences RSII sequencer with P6-C4 chemistry. One Single Molecule Real-Time (SMRT) Cell was used for each strain. Plasmid sequences were identified from whole-genome assemblies produced from hybrid read sets (both Illumina and SMRT reads) using the Unicycler pipeline v0.3.1 (Wick et al., 2017), which internally relied upon SPAdes v3.10.1 (Nurk et al., 2013), Bowtie2 v2.3.0 (Langmead and Salzberg, 2012), samtools v1.4.1 (Li et al., 2009), and Pilon v1.22 (Walker et al., 2014).

Annotated sequence for pSDE-SvHI2 and pSDC-F2\_12BHI2 have been deposited in GenBank under accession numbers MH287084 and MH287085, respectively.

## Plasmid Typing

Plasmids, particularly those that spread amongst the *Enterobacteriaceae*, have been typed by their *in vivo* incompatibility and cell exclusion patterns. Many of these incompatibility groups can now be subtyped by allelic variations in select conserved genes, forming plasmid multi-locus sequence typing schemes. Incompatibility typing and plasmid multi-locus sequence typing was performed through the Center of Genomic Epidemiology website (<http://www.genomicepidemiology.org/>) using PlasmidFinder (Carattoli et al., 2014) and the IncHI2 pDLST scheme (Garcia-Fernandez and Carattoli, 2010).

## Phylogeny and Alignment Analyses

Analyses of conserved core single nucleotide polymorphisms (SNPs) was performed using the Harvest suite (Parsnp v1.2, utilizing the Phipack recombination filter, and gngp v1.2) (Treangen et al., 2014). An analysis was run utilizing all available IncHI2 ST3 plasmids from GenBank (**Supplementary Table 1**), plus two IncHI2 ST1 plasmids, reference pR478 (NC\_005211) and pIMP4-SEM1 (KX810825). A second tree was then generated with the ST3 plasmids only, with pSDC\_F2\_12BHI2 as reference. Gene identification and genomic comparisons were performed using a combination of BLASTn (Camacho et al., 2009) and progressiveMauve (Darling et al., 2010) alignments. Figures were generated from sequence data using SnapGene v3.3.4, BRIG v0.95 (Alikhan et al., 2011), and Easyfig v2.2.2 (Sullivan et al., 2011).

Annotations were managed using SnapGene v3.3.4. Automated annotations were generated by RASTtk (Brettin et al., 2015). Insertion sequences were identified and annotated manually with the aid of ISfinder (Siguier et al., 2006).



Remaining annotations were performed manually utilizing BLASTn and publicly available databases, including The Repository of Antibiotic-Resistance Cassettes (Tsafnat et al., 2011) (<http://rac.aihi.mq.edu.au/rac/>), The Comprehensive Antibiotic Resistance Database (Jia et al., 2017) (<https://card.mcmaster.ca/>), and the GenBank nucleotide database (<https://www.ncbi.nlm.nih.gov/nucleotide/>).


## RESULTS

Unicycler hybrid assemblies resolved pSDE-SvHI2 as a 275,402 bp circular sequence and pSDC-F2\_12BHI2 as a 288,288 bp circular sequence with 344 (140 hypotheticals) and 357 (134 hypotheticals) coding sequences (CDS) respectively. Both sequences typed as incompatibility group HI2, sequence type 3 (IncHI2 ST3) plasmids. The full map of each plasmid is presented in **Figure 1**, along with content comparisons to the most related ST3 plasmids, most of which are from Australia and China. The plasmids had a typical IncHI2 structure, including the RepHIA and RepHI2 replication operons, *trh* and *tra* transfer operons and tellurite resistance (*ter* operon) (Gilmour et al., 2004), plus each carried three variants of complex resistance loci, two of which are class 1 integron associated, one encoding *sul1* in a Tn1721/Tn21 hybrid transposon, and one encoding *sul3* from within a Tn7-like copper/silver resistance transposon.

Single nucleotide polymorphism analysis was performed on the set of all ST3 plasmids (**Figure 2**, **Supplementary Table 2**), and on this same set plus IncHI2 ST1 plasmids as reference to confirm tree topology (**Supplementary Figure 1**). Forty-three plasmid sequences available from 2006 to 2017 were included in the ST3-only analysis. Based on the availability of metadata, plasmid sequences were from *Enterobacteriaceae* of different sources in the Pacific region, including Australian porcine production operations and multiple human, agricultural and environmental sources in China. The ST3 sequences formed a single clade with one exception, MH715960 from Taiwan, which separated with 125 core SNPs compared to the Australian reference sequence. Of these 125 core SNPs, 101 are within an ~1.5 kb region of the *ter* operon. The remaining sequences formed four major subclades, with closest relatives to the Australian reference ranging from two to 14 SNPs. One subclade, highlighted red in **Figure 2**, demonstrates relatedness between three Australian plasmids from the feces of pigs, pSDE-SvHI2, pSDC-F2\_12BHI2, and pSTM-275, an apparently separate Australian plasmid lineage (pIncHI2-MU3) also from pig feces, and six other plasmid sequences from diverse sources in China. Content comparisons of this clade can be seen in **Figure 1**. Of the remaining plasmids, the most distant relative had 25 SNPs identified from conserved core sequence, suggesting a close evolutionary relationship between plasmid sequences reported from Australia and China.

The ST3 plasmids carried multiple class 1 integron structures with different *sul* genes including *sul1*, *sul2*, *sul3*, and a *sul3*-associated *mefB*, encoding a macrolide efflux pump (**Figure 2**).





One sequence (KX254341) of Chinese origin was observed with only 260 bp remaining of *mefB* ( $\Delta mefB_{260}$ ). The Australian IncHI2 ST3 plasmids explored here carry  $\Delta mefB_{111}$ . The remaining *sul3*-positive plasmids carry a  $\Delta mefB_{33}$  signature, aside from CP026492 from China which completely lacks *mefB*. This includes both the pseudo-phylogenetically distant MH715960 from Taiwan and the earliest Chinese plasmid sourced from 2006. Interestingly, one single subclade of sequences from China and Hong Kong was universally negative for *intI1* and *sul3*, but not for either *sul1* or *sul2*. Many of these plasmids are also associated with various globally important resistance genes, including *bla*<sub>CTX-M</sub>, *oqxAB* and *mcr-1* (Supplementary Table 3).



pSDE-SvHI2, pSDC-F2\_12BHI2, and pSTM-275 host a class 1 integron located within a Tn1721/Tn21 hybrid tetracycline resistance transposon (Figure 3). This integron structure is host to the only complete *intI1* gene on the plasmids, and has acquired trimethoprim (*dfrA12*), *orfF* and streptomycin/spectinomycin (*aadA2*) resistance gene cassettes. Comparisons to similar Tn1721/Tn21 hybrid transposons show identical sequence across the hybridization point between *tnpR*<sub>1721</sub> and *tnpM*<sub>21</sub>, however these sequences also show a modified transposition module, with a loss of Tn1721 *tnpA* sequence between two homologous 8 bp regions (CCAGGGCG), between the second  $\Delta tnpA$ <sub>1721</sub> and its neighboring predicted relaxase. The In2-like class 1 integron *tniA* gene is truncated to 436 bp by insertion of an IS26 element with no observable associated repeats, suggesting a complex evolutionary path to this final structure. As part of the class 1 integron 3'-CS, the structure encodes sulphamide resistance (*sul1*), however in pSTM6-275 the terminal IS26 truncation lies within the *sul1* gene, giving a distinguishable gene marker.

A large (58,759 bp), mosaic combination of a globally observed Tn7-like transposon (copper and silver resistance, *sil/pco*), and transposons mobilizing antimicrobial resistance genes (Figure 1) was also observed in this Australian plasmid lineage. An IS26-associated, class 1 integron-encoding structure

has inserted near the Tn7-like transposition module (Figure 4). From the pSTM6-275 sequence, it appears this insertion was in a Tn3 associated gene, however in pSDE-SvHI2 and pSDC-F2\_12BHI2 IS26 activity has altered sequences around the insertion site. The integron is a variant of a *sul3*-associated structure that has been described previously in Australian plasmid pCERC3 (KR827684), now including signature IS26-mediated deletions in *intI1* ( $\Delta intI1_{705}$ ) and *mefB* ( $\Delta mefB_{111}$ ). The integron cassette array encodes *estX*, *psp*, *aadA2*, *cmlA*, and *aadA1*. The crossover site described near *qacH* reported in pCERC3 was identical, confirming it is a derivative of the same integron sequence encoding *sul3-qacH-mefB*, previously seen within a Tn21 background (Moran et al., 2016). We also noted an IS1203\* insertion into the *qacH* ORF, which may prove epidemiologically useful. The integron structure is followed by a short intermediate sequence encoding *rop*, then by a Tn5393 variant (encoding streptomycin resistance) with a novel IS903B\* insertion into IS1133.

From these sequences it is difficult to ascertain what originally mobilized the  $\Delta intI1$  module into the Tn7-like transposon. While IS26 flanks the captured sequence ( $\Delta intI1$  to *sul3*), no repeats have been identified to indicate a clean insertion location near the Tn7-like transposon, nor the bordering sequence near *rop*.

Of the 1,030 bp sequence by *rop*, 257 bases nearest the Tn5393 repeat are associated with the tetracycline resistance transposon Tn1721 [AJ634602.1] while the remaining 773 bp nearest the IS26/*sul3* end matched IncN plasmid backbone [HF545433.1]. These remnant sequences may give indications as to the sources of these structures should more pertinent references become available. Further, the structures we have described have undergone individual insertions and inversion events. In pSDE-SvHI2, an IS26 insertion into *tnsA* has led to the generation of new direct repeats (DR $\beta$ ) and has subsequently led to an IS26 mediated inversion. In pSDC-F2\_12BHI2, an IS26 insertion near the *tnsA* ORF has generated DR $\gamma$ , and an inversion event has again occurred, flipping the structure



between  $\Delta intI1$  and *sul3*. A separate inversion event has then occurred between the IS26 elements nearest to *rop*, re-inverting this sequence to match the original orientation. Importantly, the insertion of a complex Tn1721/Tn2/IS26 structure into Tn5393 within pSTM6-275 demonstrates further resistance consolidation occurring in the 7-year gap between pSDE-SvHI2/pSDC-F2\_12BHI2 and pSTM-275 isolations. This insert encoded *tetAB* and *bla*<sub>TEM</sub>. Unfortunately, this insertion has removed most of IS1133, and we cannot determine if pSTM6-275 carried the  $\Delta IS1133$ -IS903B\* insertion described above. The other Australian plasmid, pInCHI2-MU3, encodes the Tn7-like *sil/pco* heavy metal resistance transposon and is positive for Tn3 and *intI1*, but lacks any *sul* gene.

Finally, an IS15DI (IS6-family element; 3 SNPs compared to IS26) mobilized resistance region with a highly recombined and varied structure (Figure 5) has also been identified in pSDE-SvHI2 and pSDC-F2\_12BHI2. The original IS15DI insertion into the plasmid backbone has generated 8 bp direct repeats (AACAGCGT) that remain flanking the structure. This

composite transposon appears to mobilize neomycin resistance (*neoR*), and a truncated bleomycin resistance (*ble*) gene. It has also acquired IncN backbone (*tra* genes) and the *relBE* toxin/antitoxin system, alongside various other whole and truncated IS elements. An internal IS15DI element is present in both pSDE-SvHI2 and pSTM6-275 and is replaced by a Tn6010-like element (similar to KT716391.1, mobilized by IS15DI) carrying *oqxABR* in pSDC-F2\_12BHI2. there is an IS15DI mediated inversion of the *tra* associated region in pSDC-F2\_12BHI2. Plasmid pSDE-SvHI2 has an IS15DI insertion into IS1294, generating DR $\delta$ . This has been followed by a rearrangement event leading to the movement of DR $\delta$  and *neoR* toward the terminal IS element, including an inversion, and a loss of sequence through to the  $\Delta tnpA$ <sub>Tn3</sub>.

## DISCUSSION

Plasmids belonging to the incompatibility group HI2 are carriers of antimicrobial resistance genes globally. With the increased

availability of IncHI2 sequences, a di-locus sequence typing scheme was established (Garcia-Fernandez and Carattoli, 2010) to aid in tracking plasmid lineage development and dissemination. In Australia, most IncHI2 plasmids have been observed mobilizing *bla*<sub>IMP-4</sub>, the predominant carbapenemase-encoding gene within *Enterobacteriaceae* on the eastern seaboard of Australia (Sidjabat et al., 2015). Here we report a comprehensive analysis of two Australian IncHI2 ST3 plasmids from *E. coli*, and make comparisons to plasmid pSTM6-275 from *S. enterica* serotype 1,4,[5],12:i:- (Dyall-Smith et al., 2017). Phylogenetic analyses of the current available IncHI2 ST3 dataset showed these three Australian porcine IncHI2 ST3 plasmids are nested within a subclade alongside another Australian porcine ST3 plasmid sequence, pIncHI2-MU3 (Abraham et al., 2018), which hosts neither of the class 1 integrons identified here, but does host the metal resistance genes as part of the Tn7-like transposon. This suggests various lineages of ST3 plasmid may be circulating within Australian porcine agricultural operations.

The IncHI2 ST3 plasmids sequenced to date have been resolved from multiple *Enterobacteriaceae* including *Escherichia coli*, various *Salmonella enterica* serovars, *Shigella flexneri*, *Klebsiella pneumoniae*, and *Raoultella ornithinolytica*. Additionally, these *Enterobacteriaceae* isolates were taken from different sources, including pathogens and non-pathogens. As these ST3 plasmids carry diverse resistance gene cargo and may be found in numerous members of the *Enterobacteriaceae* family they will likely be important targets for future antimicrobial resistance surveillance (Fang et al., 2018). Our analyses also demonstrate evidence of differential gene acquisition by these plasmids, continuing to expand and alter the antimicrobial resistance gene repertoire which they are associated with internationally. Particularly, *sul3*-associated integrons carrying different  $\Delta$ *mefB* fragments were identified here.

The *sul3* gene was first described in pigs (Perreten and Boerlin, 2003) and has since been reported widely in association with *Enterobacteriaceae* in humans, farm animals and their waste—particularly swine (Antunes et al., 2007; Phuong Hoa et al., 2008; Byrne-Bailey et al., 2009; Curiao et al., 2011; Moran et al., 2016). In Australia and elsewhere, *sul3* is identified frequently in commensal *E. coli* from the feces of swine (Guerra et al., 2004; Bischoff et al., 2005; Reid et al., 2017) but less frequently in avian pathogenic *E. coli* (APEC) from diverse poultry production systems in Australia (Cummins et al., 2019). These observations suggest that swine production, and food animal production more broadly, plays a major role in the evolution of IncHI2 ST3 plasmids that carry *sul3*. Plasmid pSTM6-275 was shown to be thermostable at 44°C, and would readily conjugate at 27°C but not at 37°C, highlighting the IncHI2 plasmid families propensity to transfer under environmental conditions (Garcia-Fernandez and Carattoli, 2010; Billman-Jacobe et al., 2018).

Genetic signatures noted in this study; a combination of the IncHI2 ST3 di-locus typing alleles and *mefB*<sub>111</sub> have been seen in commensal *E. coli* short read assemblies from both the farm were strain F2\_12B was sourced, and a separate Australian

porcine operation circa 2007 (Reid et al., 2017). Combined with the isolation of pSTM6-275 in 2014, our data suggests these plasmids are purveyors of drug resistance in porcine agricultural settings, likely aided by the use of copper feed additives and the widespread use of first generation antibiotics (Jordan et al., 2009). It is notable that the IncHI2 plasmids sequenced in our study carry a unique IS6 family mobilized transposon encoding neomycin resistance, and additional fluoroquinolone resistance in pSDC-F2\_12B. Also, they are host to a globally observed Tn7-like heavy metal resistance transposon (Fang et al., 2016), that has in turn become host to a variant of a *sul3-mefB* class 1 integron, which appears unique to Australian samples at this time. The  $\Delta$ *mefB*<sub>111</sub>-associated class 1 integron was carried only by the Australian plasmids pSDE-SvHI2, pSDC-F2\_12BHI2 and pSTM-275, while  $\Delta$ *mefB*<sub>33</sub> was observed throughout the remainder of the ST3 plasmid clade. Notably,  $\Delta$ *mefB*<sub>33</sub> variants are not associated with the presence of the Tn7-like metal resistance transposon, so characterizing their distribution and methods of mobilization will be critical to monitoring complex resistance locus diversification.

Genomic sequence data has shown that IS26 is playing an important role in shaping the context of MDR islands in MDR fecal *E. coli* in Australian commercial pigs. Recently, we reported high carriage rates (101/103 isolates; 98%) of IS26 among commensal *E. coli* carrying class 1 integrons (Reid et al., 2017), however we were unable to determine the genetic context of some class 1 integrons. Nonetheless, the frequency of carriage of *sul3* and IncHI2 plasmids in our earlier study was significant in two separate commercial swine production facilities. Here we show that IS26 (and the IS15DI variant) has played a pivotal role in the evolution of several porcine IncHI2 ST3 plasmids both as a means of creating deletions and inversions and altering antibiotic resistance gene content.

Finally, the epidemiological analysis of *Enterobacteriaceae* has been heavily influenced by the spread of antimicrobial resistance and the development of MDR pathogens, leading to a focus on the detection of clinically-relevant resistance genes (*bla*<sub>IMP</sub>, *mcr-1*, and *bla*<sub>CTX-M</sub> are examples of this). To further understand the true variety and dissemination of multiple drug resistance, sampling from agricultural and environmental sources impacted by antimicrobials will be important.

## DATA AVAILABILITY

The datasets generated for this study can be found in GenBank, MH287084 and MH287085.

## AUTHOR CONTRIBUTIONS

EW performed genomic analyses, generated figures, and drafted the manuscript. TC provided curated *E. coli* collections for the study. PRC assisted with data interpretation. ML, EW, and CR prepared sequencing samples. MD performed genome assemblies. SD conceived the study and together with EW and PRC wrote the manuscript. All authors provided edits, read, and approved the manuscript.

## FUNDING

This work was in part supported by the Australian Research Council, linkage grants LP150100912 and LE150100031. This project was partly funded by the Australian Center for Genomic Epidemiological Microbiology (Ausgem), a collaborative partnership between the NSW Department of Primary Industries and the ithree institute at the University of Technology Sydney. EW and CR are recipients of Australian Government Research Training Program Scholarships.

## REFERENCES

- Abbo, L. M., and Hooton, T. M. (2014). Antimicrobial stewardship and urinary tract infections. *Antibiotics* 3, 174–192. doi: 10.3390/antibiotics3020174
- Abraham, S., Jordan, D., Wong, H. S., Johnson, J. R., Toleman, M. A., Wakeham, D. L., et al. (2015). First detection of extended-spectrum cephalosporin- and fluoroquinolone-resistant *Escherichia coli* in Australian food-producing animals. *J. Global Antimicrob. Resist.* 3, 273–277. doi: 10.1016/j.jgar.2015.08.002
- Abraham, S., Kirkwood, R. N., Laird, T., Saputra, S., Mitchell, T., Singh, M., et al. (2018). Dissemination and persistence of extended-spectrum cephalosporin-resistance encoding IncI1-blaCTXM-1 plasmid among *Escherichia coli* in pigs. *ISME J.* 12, 2352–2362. doi: 10.1038/s41396-018-0200-3
- Abraham, S., O’dea, M., Trott, D. J., Abraham, R. J., Hughes, D., Pang, S., et al. (2016). Isolation and plasmid characterization of carbapenemase (IMP-4) producing *Salmonella enterica* Typhimurium from cats. *Sci. Rep.* 6:35527. doi: 10.1038/srep35527
- Alikhan, N. F., Petty, N. K., Ben Zakour, N. L., and Beatson, S. A. (2011). BLAST Ring Image Generator (BRIG): simple prokaryote genome comparisons. *BMC Genomics* 12:402. doi: 10.1186/1471-2164-12-402
- Alonso, C. A., Michael, G. B., Li, J., Somalo, S., Simon, C., Wang, Y., et al. (2017). Analysis of blaSHV-12-carrying *Escherichia coli* clones and plasmids from human, animal and food sources. *J. Antimicrob. Chemother.* 72, 1589–1596. doi: 10.1093/jac/dkx024
- Antunes, P., Machado, J., and Peixe, L. (2007). Dissemination of sul3-containing elements linked to class 1 integrons with an unusual 3’ conserved sequence region among *Salmonella* isolates. *Antimicrob. Agents Chemother.* 51, 1545–1548. doi: 10.1128/AAC.01275-06
- Argudin, M. A., Hofer, A., and Butaye, P. (2019). Heavy metal resistance in bacteria from animals. *Res. Vet. Sci.* 122, 132–147. doi: 10.1016/j.rvsc.2018.11.007
- Bean, D. C., Livermore, D. M., Papa, I., and Hall, L. M. (2005). Resistance among *Escherichia coli* to sulphonamides and other antimicrobials now little used in man. *J. Antimicrob. Chemother.* 56, 962–964. doi: 10.1093/jac/dki332
- Billman-Jacobe, H., Liu, Y., Haites, R., Weaver, T., Robinson, L., Marena, M., et al. (2018). pSTM6-275, a conjugative IncHI2 plasmid of *Salmonella* that confers antibiotic and heavy metal resistance under changing physiological conditions. *Antimicrob. Agents Chemother.* 62:e02357-17. doi: 10.1128/AAC.02357-17
- Bischoff, K. M., White, D. G., Hume, M. E., Poole, T. L., and Nisbet, D. J. (2005). The chloramphenicol resistance gene *cmlA* is disseminated on transferable plasmids that confer multiple-drug resistance in swine *Escherichia coli*. *FEMS Microbiol. Lett.* 243, 285–291. doi: 10.1016/j.femsle.2004.12.017
- Botts, R. T., Appfel, B. A., Walters, C. J., Davidson, K. E., Echols, R. S., Geiger, M. R., et al. (2017). Characterization of four multidrug resistance plasmids captured from the sediments of an urban Coastal Wetland. *Front. Microbiol.* 8:1922. doi: 10.3389/fmicb.2017.01922
- Brettin, T., Davis, J. J., Disz, T., Edwards, R. A., Gerdes, S., Olsen, G. J., et al. (2015). RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci. Rep.* 5:8365. doi: 10.1038/srep08365
- Byrne-Bailey, K. G., Gaze, W. H., Kay, P., Boxall, A. B., Hawkey, P. M., and Wellington, E. M. (2009). Prevalence of sulfonamide resistance genes in bacterial isolates from manured agricultural soils and pig slurry in the United Kingdom. *Antimicrob. Agents Chemother.* 53, 696–702. doi: 10.1128/AAC.00652-07
- Cain, A. K., and Hall, R. M. (2012). Evolution of IncHI2 plasmids via acquisition of transposons carrying antibiotic resistance determinants. *J. Antimicrob. Chemother.* 67, 1121–1127. doi: 10.1093/jac/dks004
- Cain, A. K., Liu, X., Djordjevic, S. P., and Hall, R. M. (2010). Transposons related to Tn1696 in IncHI2 plasmids in multiply antibiotic resistant *Salmonella enterica* serovar Typhimurium from Australian animals. *Microb. Drug Resist.* 16, 197–202. doi: 10.1089/mdr.2010.0042
- Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., et al. (2009). BLAST+: architecture and applications. *BMC Bioinformatics* 10:421. doi: 10.1186/1471-2105-10-421
- Carattoli, A., Zankari, E., Garcia-Fernandez, A., Voldby Larsen, M., Lund, O., Villa, L., et al. (2014). *In silico* detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob. Agents Chemother.* 58, 3895–3903. doi: 10.1128/AAC.02412-14
- Chavda, K. D., Chen, L., Jacobs, M. R., Rojtmán, A. D., Bonomo, R. A., and Kreiswirth, B. N. (2015). Complete sequence of a bla(KPC)-harboring cointegrate plasmid isolated from *Escherichia coli*. *Antimicrob. Agents Chemother.* 59, 2956–2959. doi: 10.1128/AAC.00041-15
- Cummins, M. L., Reid, C. J., Roy Chowdhury, P., Bushell, R. N., Esbert, N., Tivendale, K. A., et al. (2019). Whole genome sequence analysis of Australian avian pathogenic *Escherichia coli* that carry the class 1 integrase gene. *Microb. Genomics* 5, 1–13. doi: 10.1099/mgen.0.000250
- Curiao, T., Canton, R., Garcillan-Barcia, M. P., De La Cruz, F., Baquero, F., and Coque, T. M. (2011). Association of composite IS26-sul3 elements with highly transmissible IncI1 plasmids in extended-spectrum-beta-lactamase-producing *Escherichia coli* clones from humans. *Antimicrob. Agents Chemother.* 55, 2451–2457. doi: 10.1128/AAC.01448-10
- Darling, A. E., Mau, B., and Perna, N. T. (2010). progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS ONE* 5:e11147. doi: 10.1371/journal.pone.0011147
- Dawes, F. E., Kuzevski, A., Bettelheim, K. A., Hornitzky, M. A., Djordjevic, S. P., and Walker, M. J. (2010). Distribution of class 1 integrons with IS26-mediated deletions in their 3’-conserved segments in *Escherichia coli* of human and animal origin. *PLoS ONE* 5:e12754. doi: 10.1371/journal.pone.0012754
- Delannoy, S., Le Devendec, L., Jouy, E., Fach, P., Drider, D., and Kempf, I. (2017). Characterization of colistin-resistant *Escherichia coli* isolated from diseased pigs in France. *Front. Microbiol.* 8:2278. doi: 10.3389/fmicb.2017.02278
- Dionisi, A. M., Lucarelli, C., Owczarek, S., Luzzi, I., and Villa, L. (2009). Characterization of the plasmid-borne quinolone resistance gene *qnrB19* in *Salmonella enterica* serovar Typhimurium. *Antimicrob. Agents Chemother.* 53, 4019–4021. doi: 10.1128/AAC.00294-09
- Djordjevic, S. P., Stokes, H. W., and Roy Chowdhury, P. (2013). Mobile elements, zoonotic pathogens and commensal bacteria: conduits for the delivery of resistance genes into humans, production animals and soil microbiota. *Front. Microbiol.* 4:86. doi: 10.3389/fmicb.2013.00086
- Dolejska, M., Masarikova, M., Dobiasova, H., Jamborova, I., Karpiskova, R., Havlicek, M., et al. (2016). High prevalence of *Salmonella* and IMP-4-producing *Enterobacteriaceae* in the silver gull on Five Islands, Australia. *J. Antimicrob. Chemother.* 71, 63–70. doi: 10.1093/jac/dkv306
- Dyall-Smith, M. L., Liu, Y., and Billman-Jacobe, H. (2017). Genome sequence of an Australian monophasic *Salmonella enterica* subsp.

## ACKNOWLEDGMENTS

We are grateful for the helpful discussions with Helen Jacobe-Billman.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fsufs.2019.00018/full#supplementary-material>

- enterica Typhimurium Isolate (TW-Stm6) carrying a large plasmid with multiple antimicrobial resistance genes. *Genome Announc.* 5:e00793-17. doi: 10.1128/genomeA.00793-17
- Enne, V. I., Livermore, D. M., Stephens, P., and Hall, L. M. (2001). Persistence of sulphonamide resistance in *Escherichia coli* in the UK despite national prescribing restriction. *Lancet* 357, 1325–1328. doi: 10.1016/S0140-6736(00)04519-0
- Falgenhauer, L., Ghosh, H., Guerra, B., Yao, Y., Fritzenwanker, M., Fischer, J., et al. (2017). Comparative genome analysis of IncHI2 VIM-1 carbapenemase-encoding plasmids of *Escherichia coli* and *Salmonella enterica* isolated from a livestock farm in Germany. *Vet. Microbiol.* 200, 114–117. doi: 10.1016/j.vetmic.2015.09.001
- Fang, L., Li, X., Li, L., Li, S., Liao, X., Sun, J., et al. (2016). Co-spread of metal and antibiotic resistance within ST3-IncHI2 plasmids from *E. coli* isolates of food-producing animals. *Sci. Rep.* 6:25312. doi: 10.1038/srep25312
- Fang, L.-X., Li, X.-P., Deng, G.-H., Li, S.-M., Yang, R.-S., Wu, Z.-W., et al. (2018). High genetic plasticity in multidrug resistant ST3-IncHI2 plasmids revealed by sequence comparison and phylogenetic analysis. *Antimicrob. Agents Chemother.* 62:e02068-17. doi: 10.1128/AAC.02068-17
- Garcia Fernandez, A., Cloeckaert, A., Bertini, A., Praud, K., Doublet, B., Weill, F. X., et al. (2007). Comparative analysis of IncHI2 plasmids carrying blaCTX-M-2 or blaCTX-M-9 from *Escherichia coli* and *Salmonella enterica* strains isolated from poultry and humans. *Antimicrob. Agents Chemother.* 51, 4177–4180. doi: 10.1128/AAC.00603-07
- Garcia, V., Garcia, P., Rodriguez, I., Rodicio, R., and Rodicio, M. R. (2016). The role of IS26 in evolution of a derivative of the virulence plasmid of *Salmonella enterica* serovar Enteritidis which confers multiple drug resistance. *Infect. Genet. Evol.* 45, 246–249. doi: 10.1016/j.meegid.2016.09.008
- Garcia-Fernandez, A., and Carattoli, A. (2010). Plasmid double locus sequence typing for IncHI2 plasmids, a subtyping scheme for the characterization of IncHI2 plasmids carrying extended-spectrum beta-lactamase and quinolone resistance genes. *J. Antimicrob. Chemother.* 65, 1155–1161. doi: 10.1093/jac/dkq101
- Gilmour, M. W., Thomson, N. R., Sanders, M., Parkhill, J., and Taylor, D. E. (2004). The complete nucleotide sequence of the resistance plasmid R478: defining the backbone components of incompatibility group H conjugative plasmids through comparative genomics. *Plasmid* 52, 182–202. doi: 10.1016/j.plasmid.2004.06.006
- Gilrane, V. L., Lobo, S., Huang, W., Zhuge, J., Yin, C., Chen, D., et al. (2017). Complete genome sequence of a colistin-resistant *Escherichia coli* strain harboring mcr-1 on an IncHI2 plasmid in the United States. *Genome Announc.* 5:e01095-17. doi: 10.1128/genomeA.01095-17
- Grape, M., Sundstrom, L., and Kronvall, G. (2003). Sulphonamide resistance gene sul3 found in *Escherichia coli* isolates from human sources. *J. Antimicrob. Chemother.* 52, 1022–1024. doi: 10.1093/jac/dkg473
- Guerra, B., Junker, E., and Helmuth, R. (2004). Incidence of the recently described sulfonamide resistance gene sul3 among German *Salmonella enterica* strains isolated from livestock and food. *Antimicrob. Agents Chemother.* 48, 2712–2715. doi: 10.1128/AAC.48.7.2712-2715.2004
- Harmer, C. J., Moran, R. A., and Hall, R. M. (2014). Movement of IS26-associated antibiotic resistance genes occurs via a translocatable unit that includes a single IS26 and preferentially inserts adjacent to another IS26. *MBio* 5:e01801–01814. doi: 10.1128/mBio.01801-14
- Huang, X., Yu, L., Chen, X., Zhi, C., Yao, X., Liu, Y., et al. (2017). High prevalence of colistin resistance and mcr-1 gene in *Escherichia coli* isolated from food animals in China. *Front. Microbiol.* 8:562. doi: 10.3389/fmicb.2017.00562
- Jia, B., Raphenya, A. R., Alcock, B., Waglechner, N., Guo, P., Tsang, K. K., et al. (2017). CARD 2017: expansion and model-centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res.* 45, D566–d573. doi: 10.1093/nar/gkw1004
- Jordan, D., Chin, J. J., Fahy, V. A., Barton, M. D., Smith, M. G., and Trott, D. J. (2009). Antimicrobial use in the Australian pig industry: results of a national survey. *Aust. Vet. J.* 87, 222–229. doi: 10.1111/j.1751-0813.2009.00430.x
- Kidsley, A. K., Abraham, S., Bell, J. M., O’dea, M., Laird, T. J., Jordan, D., et al. (2018). Antimicrobial susceptibility of *Escherichia coli* and *Salmonella* spp. isolates from healthy pigs in Australia: results of a Pilot National Survey. *Front. Microbiol.* 9:1207. doi: 10.3389/fmicb.2018.01207
- Kurenbach, B., Hill, A. M., Godsoe, W., Van Hamelsveld, S., and Heinemann, J. A. (2018). Agrichemicals and antibiotics in combination increase antibiotic resistance evolution. *PeerJ* 6:e5801. doi: 10.7717/peerj.5801
- Kurenbach, B., Marjoshi, D., Amabile-Cuevas, C. F., Ferguson, G. C., Godsoe, W., Gibson, P., et al. (2015). Sublethal exposure to commercial formulations of the herbicides dicamba, 2,4-dichlorophenoxyacetic acid, and glyphosate cause changes in antibiotic susceptibility in *Escherichia coli* and *Salmonella enterica* serovar Typhimurium. *MBio* 6:e00009-15. doi: 10.1128/mBio.00009-15
- Lai, J., Wang, Y., Shen, J., Li, R., Han, J., Foley, S. L., et al. (2013). Unique class 1 integron and multiple resistance genes co-located on IncHI2 plasmid is associated with the emerging multidrug resistance of *Salmonella* Indiana isolated from chicken in China. *Foodborne Pathog. Dis.* 10, 581–588. doi: 10.1089/fpd.2012.1455
- Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat. Methods* 9, 357–359. doi: 10.1038/nmeth.1923
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25, 2078–2079. doi: 10.1093/bioinformatics/btp352
- Li, R., Xie, M., Zhang, J., Yang, Z., Liu, L., Liu, X., et al. (2017). Genetic characterization of mcr-1-bearing plasmids to depict molecular mechanisms underlying dissemination of the colistin resistance determinant. *J. Antimicrob. Chemother.* 72, 393–401. doi: 10.1093/jac/dkw411
- Mangat, C. S., Bekal, S., Irwin, R. J., and Mulvey, M. R. (2017). A Novel hybrid plasmid carrying multiple antimicrobial resistance and virulence genes in *Salmonella enterica* Serovar Dublin. *Antimicrob. Agents Chemother.* 61:e02601-16. doi: 10.1128/AAC.02601-16
- Mir, R. A., Weppelmann, T. A., Teng, L., Kirpich, A., Elzo, M. A., Driver, J. D., et al. (2018). Colonization dynamics of cefotaxime resistant bacteria in beef cattle raised without cephalosporin antibiotics. *Front. Microbiol.* 9:500. doi: 10.3389/fmicb.2018.00500
- Monahan, L. G., Demaere, M. Z., Cummins, M. L., Djordjevic, S. P., Roy Chowdhury, P., and Darling, A. E. (2019). High contiguity genome sequence of a multidrug-resistant hospital isolate of *Enterobacter hormaechei*. *Gut Pathog.* 11, 3. doi: 10.1186/s13099-019-0288-7
- Moran, R. A., Holt, K. E., and Hall, R. M. (2016). pCERC3 from a commensal ST95 *Escherichia coli*: a ColV virulence-multiresistance plasmid carrying a sul3-associated class 1 integron. *Plasmid* 84–85, 11–19. doi: 10.1016/j.plasmid.2016.02.002
- Nurk, S., Bankevich, A., Antipov, D., Gurevich, A., Korobeynikov, A., Lapidus, A., et al. (2013). *Assembling Genomes and Mini-metagenomes from Highly Chimeric Reads*. Berlin; Heidelberg: Springer, 158–170.
- Perreten, V., and Boerlin, P. (2003). A new sulfonamide resistance gene (sul3) in *Escherichia coli* is widespread in the pig population of Switzerland. *Antimicrob. Agents Chemother.* 47, 1169–1172. doi: 10.1128/AAC.47.3.1169-1172.2003
- Phuong Hoa, P. T., Nonaka, L., Hung Viet, P., and Suzuki, S. (2008). Detection of the sul1, sul2, and sul3 genes in sulfonamide-resistant bacteria from wastewater and shrimp ponds of north Vietnam. *Sci. Total Environ.* 405, 377–384. doi: 10.1016/j.scitotenv.2008.06.023
- Poirel, L., Jayol, A., and Nordmann, P. (2017). Polymyxins: antibacterial activity, susceptibility testing, and resistance mechanisms encoded by plasmids or chromosomes. *Clin. Microbiol. Rev.* 30, 557–596. doi: 10.1128/CMR.00064-16
- Porse, A., Schonning, K., Munck, C., and Sommer, M. O. (2016). Survival and evolution of a large multidrug resistance plasmid in new clinical bacterial hosts. *Mol. Biol. Evol.* 33, 2860–2873. doi: 10.1093/molbev/msw163
- Razavi, M., Marathe, N. P., Gillings, M. R., Flach, C. F., Kristianson, E., and Joakim Larsson, D. G. (2017). Discovery of the fourth mobile sulfonamide resistance gene. *Microbiome* 5, 160. doi: 10.1186/s40168-017-0379-y
- Reid, C. J., Roy Chowdhury, P., and Djordjevic, S. P. (2015). Tn6026 and Tn6029 are found in complex resistance regions mobilised by diverse plasmids and chromosomal islands in multiple antibiotic resistant *Enterobacteriaceae*. *Plasmid* 80, 127–137. doi: 10.1016/j.plasmid.2015.04.005
- Reid, C. J., Wyrsch, E. R., Roy Chowdhury, P., Zingali, T., Liu, M., Darling, A. E., et al. (2017). Porcine commensal *Escherichia coli*: a reservoir for class 1 integrons associated with IS26. *Microb. Genomics* 3, 1–42. doi: 10.1099/mgen.0.000143
- Roy Chowdhury, P., Charles, I. G., and Djordjevic, S. P. (2015). A role for Tn6029 in the evolution of the complex antibiotic resistance gene loci in genomic

- island 3 in enteroaggregative hemorrhagic *Escherichia coli* O104:H4. *PLoS ONE* 10:e0115781. doi: 10.1371/journal.pone.0115781
- Roy Chowdhury, P., Mckinnon, J. M., Liu, M. Y., and Djordjevic, S. P. (2018). Multidrug resistant uropathogenic *Escherichia coli* ST405 with a novel, composite IS26 transposon in a unique chromosomal location. *Front. Microbiol.* 9:3212. doi: 10.3389/fmicb.2018.03212
- Saputra, S., Jordan, D., Mitchell, T., Wong, H. S., Abraham, R. J., Kidsley, A., et al. (2017). Antimicrobial resistance in clinical *Escherichia coli* isolated from companion animals in Australia. *Vet. Microbiol.* 211, 43–50. doi: 10.1016/j.vetmic.2017.09.014
- Shahid, M. (2010). Citrobacter spp. simultaneously harboring blaCTX-M, blaTEM, blaSHV, blaampC, and insertion sequences IS26 and orf513: an evolutionary phenomenon of recent concern for antibiotic resistance. *J. Clin. Microbiol.* 48, 1833–1838. doi: 10.1128/JCM.01467-09
- Sidjabat, H. E., Townell, N., Nimmo, G. R., George, N. M., Robson, J., Vohra, R., et al. (2015). Dominance of IMP-4-producing *Enterobacter cloacae* among carbapenemase-producing *Enterobacteriaceae* in Australia. *Antimicrob. Agents Chemother.* 59, 4059–4066. doi: 10.1128/AAC.04378-14
- Siguier, P., Perochon, J., Lestrade, L., Mahillon, J., and Chandler, M. (2006). ISfinder: the reference centre for bacterial insertion sequences. *Nucleic Acids Res.* 34, D32–36. doi: 10.1093/nar/gkj014
- Suhartono, S., Savin, M. C., and Gbur, E. E. (2017). Transmissible plasmids and integrons shift *Escherichia coli* population toward larger multiple drug resistance numbers. *Microb Drug Resist.* 24, 244–252. doi: 10.1089/mdr.2016.0329
- Sullivan, M. J., Petty, N. K., and Beatson, S. A. (2011). Easyfig: a genome comparison visualizer. *Bioinformatics* 27, 1009–1010. doi: 10.1093/bioinformatics/btr039
- Treangen, T. J., Ondov, B. D., Koren, S., and Phillippy, A. M. (2014). The Harvest suite for rapid core-genome alignment and visualization of thousands of intraspecific microbial genomes. *Genome Biol.* 15, 524. doi: 10.1186/s13059-014-0524-x
- Tsafnat, G., Copty, J., and Partridge, S. R. (2011). RAC: Repository of Antibiotic resistance Cassettes. *Database* 2011:bar054. doi: 10.1093/database/bar054
- Turner, A. (2011). Quarantine, exports and animal disease in Australia 1901–2010. *Aust. Vet. J.* 89, 366–371. doi: 10.1111/j.1751-0813.2011.00822.x
- Venturini, C., Beatson, S. A., Djordjevic, S. P., and Walker, M. J. (2010). Multiple antibiotic resistance gene recruitment onto the enterohemorrhagic *Escherichia coli* virulence plasmid. *FASEB J.* 24, 1160–1166. doi: 10.1096/fj.09-144972
- Venturini, C., Hassan, K. A., Roy Chowdhury, P., Paulsen, I. T., Walker, M. J., and Djordjevic, S. P. (2013). Sequences of two related multiple antibiotic resistance virulence plasmids sharing a unique IS26-related molecular signature isolated from different *Escherichia coli* pathotypes from different hosts. *PLoS ONE* 8:e78862. doi: 10.1371/journal.pone.0078862
- Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., et al. (2014). Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS ONE* 9:e112963. doi: 10.1371/journal.pone.0112963
- Wick, R. R., Judd, L. M., Gorrie, C. L., and Holt, K. E. (2017). Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput. Biol.* 13:e1005595. doi: 10.1371/journal.pcbi.1005595
- Wong, M. H., Chan, E. W., and Chen, S. (2017). IS26-mediated formation of a virulence and resistance plasmid in *Salmonella* Enteritidis. *J. Antimicrob. Chemother.* 72, 2750–2754. doi: 10.1093/jac/dkx238
- Wyrsh, E., Roy Chowdhury, P., Abraham, S., Santos, J., Darling, A. E., Charles, I. G., et al. (2015). Comparative genomic analysis of a multiple antimicrobial resistant enterotoxigenic *E. coli* O157 lineage from Australian pigs. *BMC Genomics* 16:165. doi: 10.1186/s12864-015-1382-y
- Wyrsh, E. R., Roy Chowdhury, P., Chapman, T. A., Charles, I. G., Hammond, J. M., and Djordjevic, S. P. (2016). Genomic microbial epidemiology is needed to comprehend the global problem of antibiotic resistance and to improve pathogen diagnosis. *Front. Microbiol.* 7:843. doi: 10.3389/fmicb.2016.00843
- Zheng, B., Huang, C., Xu, H., Guo, L., Zhang, J., Wang, X., et al. (2017). Occurrence and genomic characterization of ESBL-producing, MCR-1-harboring *Escherichia coli* in farming soil. *Front. Microbiol.* 8:2510. doi: 10.3389/fmicb.2017.02510
- Zhou, T., Lu, J., Tong, Y., Li, S., and Wang, X. (2014). Distribution of antibiotic resistance genes in Bosten Lake, Xinjiang, China. *Water Sci. Technol.* 70, 925–931. doi: 10.2166/wst.2014.321

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Wyrsh, Reid, DeMaere, Liu, Chapman, Roy Chowdhury and Djordjevic. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.