

THE LU-LC CONJECTURE IS FALSE

ZHENG FENG JI

*State Key Laboratory of Computer Science, Institute of Software, Chinese Academy of Sciences
P.O.Box 8718, Beijing 100190, China*

JIANXIN CHEN, ZHAOHUI WEI, and MINGSHENG YING

*State Key Laboratory of Intelligent Technology and Systems,
Department of Computer Science and Technology, Tsinghua University, Beijing 100084, China*

Received January 26, 2009

Revised September 27, 2009

The LU-LC conjecture is an important open problem concerning the structure of entanglement of states described in the stabilizer formalism. It states that two local unitary equivalent stabilizer states are also local Clifford equivalent. If this conjecture were true, the local equivalence of stabilizer states would be extremely easy to characterize. Unfortunately, however, based on the recent progress made by Gross and Van den Nest, we find that the conjecture is false.

Keywords: Entanglement, Stabilizer States, Graph States, LU-LC Conjecture

Communicated by: R Jozsa & B Terhal

1 Introduction

The stabilizer formalism is a group-theoretic framework originally devised to systematically analyze various quantum error-correcting codes [6] and has found applications in other areas of quantum information processing, such as the one-way computation model [12, 4] and the quantum sharing of classical secrets [1].

One of the key features of stabilizer states defined by the stabilizer formalism is the possible presence of high-degree multipartite entanglement in them. A natural approach to studying the properties of the entanglement in stabilizer states is to investigate their local equivalences. Mainly, three types of local equivalences have been studied with respect to stochastic local operations and classical communication (SLOCC), local unitary (LU), and local Clifford operations (LC). In the following, we will call two states SLOCC (LU, LC) equivalent if they can be transformed to each other by means of SLOCC (LU, LC) operations.

Local Clifford operations are local unitary operations that map the Pauli group to itself under conjugation. Therefore, the action of a local Clifford operation on stabilizer state can be represented as a simple way of translating the stabilizers. Because of this close relationship between the stabilizer formalism and the Clifford group, the LC equivalence of stabilizer states has been studied thoroughly in the literature. For example, a polynomial time algorithm has been found to decide whether two stabilizer states are LC equivalent [16] and the action of local Clifford group on graph states (an important subset of stabilizer states) has been translated into elementary graph transformations characterized by a single rule [17]. Specifically, the classification of LC equivalent stabilizer states has been performed systematically up to 12 qubits [11, 2, 3].

Stabilizer states have extremely symmetric structures that put strong restrictions on the local unitary that can map one stabilizer state to the other. In fact, it has been conjectured for several years that any two LU equivalent stabilizer states must also be LC equivalent (the LU-LC conjecture, listed as the 28th open problem in quantum information [14]). If this conjecture were true, all three local equivalence of stabilizer states would be the same, as it has already been proven that two stabilizer states are SLOCC equivalent if and only if they are LU equivalent [15]. Moreover, deciding whether two states are locally equivalent would be efficient and the local equivalence of graph states could be described as purely graph theoretic terms. The conjecture has been proved for large subclasses of stabilizer states in Refs. [19, 21] and is further supported by the results obtained in Ref. [18].

In this paper, however, we will present a counterexample to the LU-LC conjecture that disproves the conjecture in general. Our result is build upon the recent progress that transforms the conjecture to a simpler problem [9]. In addition to our heuristics for the construction of counterexamples, we also present some of the special cases in which LU and LC equivalence are the same.

The organization of this paper is as follows. In the next section, we introduce the basics of stabilizer formalism and some notations and results used in this paper. Sec. 3 develops new representation of the problem in the first part and then considers some special cases of the LU-LC conjecture. The procedure of generating counterexamples and one explicit example are given in Sec. 4. We conclude in Sec. 5 and discuss some possible future work on this problem.

2 Notations

Stabilizer states are quantum states described by a set of commuting operators. This idea of representing states with operators has been proved to be extremely useful in the theory of quantum information. Let $\{I, X, Y, Z\}$ be the identity and Pauli matrices,

$$X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad Y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad Z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix},$$

\mathcal{G} be the Pauli group generated by them, and I_n be $I^{\otimes n}$. Mathematically, a stabilizer of n qubits is an Abelian subgroup of $\mathcal{G}^{\otimes n}$ that does not include $-I_n$. When the subgroup has cardinality exactly 2^n , there will be a unique quantum state determined by it as the simultaneous fix point of all the operators in the subgroup. For example, the EPR pair $(|00\rangle + |11\rangle)/\sqrt{2}$ is stabilized by the group generated by $\{X \otimes X, Z \otimes Z\}$. For any graph G , the corresponding graph state is a special stabilizer state with

$$X_v \bigotimes_{u \in N(v)} Z_u, \quad v \in V(G)$$

as generators of its stabilizer, where $N(v)$ is the neighbor set of vertex v . It is known that any stabilizer state is LC equivalent to some graph state [17, 13, 7]. Therefore, the LU-LC conjecture for stabilizer states and graph states are the same problem. For more details on the power of stabilizer formalism and graph states, the readers are referred to Refs. [6, 10].

There are special structures in the amplitudes of stabilizer states expanded in the compu-

tational basis. It is proved that any stabilizer state can be written as

$$\frac{1}{\sqrt{|T|}} \sum_{x \in T} i^{l(x)} (-1)^{q(x)} |x\rangle, \quad (1)$$

where T is an affine space of \mathbb{F}_2^n , $l(x)$ is linear in x_j with addition modulo 2, and $q(x)$ is a quadratic function of x_j 's. Conversely, any state having the above form is a stabilizer state [5].

The main topic in this paper is to study local equivalences of stabilizer states, especially the LU equivalence and LC equivalence. Local Clifford operators are 2 by 2 unitary operators U such that $U\mathcal{G}U^\dagger = \mathcal{G}$. Up to a global phase, the local Clifford operators form a finite subgroup of $U(2)$. We say two stabilizers $|\psi_0\rangle$ and $|\psi_1\rangle$ are LC equivalent if there exists local Clifford operators U_j such that

$$\bigotimes_{j=1}^n U_j |\psi_0\rangle = |\psi_1\rangle.$$

In the following, we define several problems, each of which is given an abbreviated name for later references. The first one is the original LU-LC conjecture itself, then a restricted case of it, DLU-LC. The third is the DLU-DLC problem and the last is quadratic form phase problem (QFP). One can see the close relationship between QFP and the other three problems from Eq. (1). The previously known relation of these problems is as follows. DLU-LC is shown to be equivalent to the LU-LC conjecture [9, 22]. DLU-DLC and QFP are equivalent and they imply the LU-LC conjecture according to Ref. [9]. All of these four statements are false as indicated by our counterexample given in Sec. 4.

Problem 1 (LU-LC) *Every two LU equivalent stabilizer states are also LC equivalent.*

Problem 2 (DLU-LC) *Every two stabilizer states that can be mapped onto each other by means of a diagonal local unitary, are LC equivalent.*

Problem 3 (DLU-DLC) *If two stabilizer states can be mapped onto each other by means of a diagonal local unitary, then also by a diagonal local Clifford operation.*

Problem 4 (QFP [9]) *Let S be a linear subspace of \mathbb{F}_2^n , and $Q : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ be a quadratic function. If there exists complex phases $\{c_i\}$ such that*

$$(-1)^{Q(x)} = \prod_j^n c_j^{x_j}, \quad \text{for every } x \in S, \quad (2)$$

then the phases can be chosen from $\{\pm 1, \pm i\}$.

3 Simplification of the QFP Problem

Let us start by simplifying the QFP problem. We will give a linearized version of QFP in the following. It's easy to obtain linear representations of QFP by, for example, enumerating all possible $x \in S$ and resulting in a system of linear congruence equations. But this method does not give us much information on how to tackle the problem. What we present in the following is a more symmetric linear representation that leads to the construction of counterexamples of QFP.

3.1 Symmetric Linear Representation

There are two main mathematical objects in the problem, the quadratic function $Q(x)$ and the subspace S . We will find a representation of $Q(x)$ first and then analyze the structure of S . Based on this analysis, some special cases of QFP will be proved. We will also obtain an approximate restatement of the general QFP problem that leads to the construction of counterexamples.

First, noticing that the linear terms in $Q(x)$ can be moved to the right hand side of Eq. (2) without changing the problem, we can consider quadratic forms only. There is a trivial one-to-one correspondence between quadratic forms over \mathbb{F}_2 and simple graphs. For each $Q(x)$, one can identify it with a graph \mathcal{Q} with vertex set $V = \{1, 2, \dots, n\}$ and edge set

$$E(\mathcal{Q}) = \{(i, j) \mid i, j \in V \text{ and } x_i x_j \text{ is a term in } Q(x)\}.$$

Each subset A of V defines a subgraph $\mathcal{Q}|_A$ whose vertex set and edge set are A and $E(\mathcal{Q}) \cap A^2$ respectively. With the above definitions, one can verify that

$$Q(x) = \left| E(\mathcal{Q}|_{I_x}) \right|, \quad (3)$$

where I_x is the indicator set $\{j \mid x_j = 1\}$ and $\mathcal{Q}|_{I_x}$ is the subgraph of \mathcal{Q} which contains only the vertices for which $x_j = 1$.

Next, to analyze the structure of S , we choose a basis $\{\xi^1, \xi^2, \dots, \xi^d\}$ of S . For each $x \in S$, we can represent it as the linear combination

$$x = \sum_{k=1}^d h_k \xi^k.$$

Let $h \in \mathbb{F}_2^d$ be the vector whose k -th coordinate is h_k and we label the above x with it as x^h . The size of S is therefore $D = 2^d$.

We introduce a new concept called *pattern* of a position in S . The pattern for position j is defined to be the vector $m \in \mathbb{F}_2^d$ whose k -th coordinate m_k equals to ξ_j^k . In other words, it's simply the j -th column of a generator matrix for S . See Fig. 1 for an illustration of this definition. Patterns are basis dependant but the choice of basis does not change the following analysis.

Let A_m be the set of positions having pattern m , that is,

$$A_m = \{j \mid m_k = \xi_j^k \text{ for all } k\}.$$

The collection $\{A_m\}$ is obviously a partition of the vertex set V of graph \mathcal{Q} . For any $j \in A_m$, one can calculate the j -th coordinate of x^h as

$$x_j^h = \sum_{k=1}^d h_k \xi_j^k = \sum_{k=1}^d h_k m_k \stackrel{\text{def}}{=} \langle h, m \rangle, \quad (4)$$

where the summation is taken over \mathbb{F}_2 , or modulo 2. That is, the value of x^h at position j is determined solely by the pattern of the position. If position i and j have the same pattern, variables x_i and x_j will always be the same. We can therefore define variable $x_{[m]}$ to be a representative variable of those having pattern m .




Fig. 1. Illustration of patterns

Consider a special case of QFP where A_m is nonempty for all patterns $m \in \mathbb{F}_2^d - \{0\}$. We will prove that in this case QFP is true and the details of the calculation will be useful even when we deal with the general QFP problem.

As all A_m 's are nonempty, we can rewrite Eq. (2) in the QFP problem by replacing the variables with their representatives in the right hand side,

$$(-1)^{Q(x)} = \prod_{m \neq 0} C_{[m]}^{x[m]}, \quad \text{for every } x \in S,$$

where $C_{[m]}$ is the product of all c_j for $j \in A_m$. For a specific $x^h \in S$, this means that

$$(-1)^{Q(x^h)} = \prod_{m \neq 0} C_{[m]}^{\langle h, m \rangle}. \quad (5)$$

From Eq. (3) and the fact that

$$I_{x^h} = \bigcup_{m: \langle m, h \rangle = 1} A_m, \quad (6)$$

we have

$$\begin{aligned} Q(x^h) &= \sum_{m: \langle m, h \rangle = 1} \left| E(\mathcal{Q}|_{A_m}) \right| + \sum_{m < m': \langle m, h \rangle = 1, \langle m', h \rangle = 1} E_{mm'} \\ &= \sum_m \langle m, h \rangle \left| E(\mathcal{Q}|_{A_m}) \right| + \sum_{m < m'} \langle m, h \rangle \langle m', h \rangle E_{mm'}, \end{aligned}$$

where $E_{mm'}$ counts the number of edges between vertex sets A_m and $A_{m'}$ in graph \mathcal{Q} . The first part of the above summation can be omitted as it can be absorbed into the right hand side of Eq. (5) by changing the values of $C_{[m]}$ appropriately. Equation (5) is therefore reduced into the following form,

$$(-1)^{\sum_{m < m'} \langle m, h \rangle \langle m', h \rangle E_{mm'}} = \prod_{m \neq 0} C_{[m]}^{\langle m, h \rangle}. \quad (7)$$

Let $r_{[m]}$ be the real number satisfying

$$C_{[m]} = i^{r_{[m]}}. \quad (8)$$

By taking logarithm on both side of Eq. (7), one gets

$$\sum_{m \neq 0} \langle m, h \rangle r_{[m]} \equiv 2 \sum_{m < m'} \langle m, h \rangle \langle m', h \rangle E_{mm'} \pmod{4}.$$

This equation holds for all $h \in \mathbb{F}_2^d$, and it actually specifies a system of $D - 1$ equations where $D = 2^d$. By defining $D - 1$ by $D - 1$ matrix $G = (\langle i, j \rangle)$ and matrix T with element

$$T_{i,(j,k)} = \langle i, j \rangle \langle i, k \rangle, \quad i, j, k \in \mathbb{F}_2^d - \{0\}, j < k, \quad (9)$$

we can write the system of equations succinctly as

$$G\vec{r} \equiv 2T\vec{e} \pmod{4} \quad (10)$$

where \vec{r} and \vec{e} are vectors consisting of $r_{[m]}$ and $E_{mm'}$ respectively.

Next, we calculate G^{-1} and the product $2G^{-1}T$ based on the following lemma.

Lemma 1 *The following identities hold with the indexes traversing $\mathbb{F}_2^d - \{0\}$.*

$$\sum_j \langle i, j \rangle = D/2, \quad (11)$$

$$\sum_j \langle i, j \rangle \langle j, k \rangle = \begin{cases} D/2 & i = k \\ D/4 & \text{otherwise} \end{cases} \quad (12)$$

For $k \neq l$, we have

$$\sum_j \langle i, j \rangle \langle j, k \rangle \langle j, l \rangle = \begin{cases} D/4 & i = k \text{ or } i = l \\ 0 & i = k \oplus l \\ D/8 & \text{otherwise} \end{cases} \quad (13)$$

We prove the last identity only; others are simpler and can be dealt with similarly. When $i = k$ or $i = l$ the result follows from the second identity. In the case of $i = k \oplus l$, there does not exist j so that $\langle j, i \rangle, \langle j, k \rangle, \langle j, l \rangle$ are all 1. What remains to show is when $i \neq k, i \neq l, k \neq l$ and $i \neq k \oplus l$. Consider a 3 by d matrix F with i, k, l as its rows, then it is a matrix of rank 3 under the above conditions. j contributes 1 to the sum when it is a solution of $Fj = (1, 1, 1)^T$. The number of solutions is obviously $D/8$.

With the above identities, it's easy to check that G^{-1} is given by

$$G^{-1} = \frac{2}{D}(2\langle i, j \rangle - 1), \quad (14)$$

which is obtained by replacing the zero elements in G with -1 and multiplying a normalization constant. More importantly, the product $2G^{-1}T$ is still an integral matrix with 1 at $(i, (j, k))$ where $i = j$ or $i = k$, -1 where $i = j \oplus k$ and 0's elsewhere. This immediately proves the QFP problem in the special case where A_m 's are all nonempty, as we can always choose \vec{r} to be

$2G^{-1}T\vec{e}$, an integral vector. The proof is completed without even referring to the condition that there exists a real solution of \vec{r} for Eq. (10) as promised by the QFP problem.

However, the general case QFP problem is much more complicated and, in fact, false sometimes. When A_m is empty for some pattern m , we cannot introduce representative variable $x_{[m]}$ as there is no such corresponding variables. In order to make use of the above formalism, however, we keep the variables $x_{[m]}$ for all those missing patterns m and, at the same time, fix the corresponding $C_{[m]}$'s to be 1, or fix $r_{[m]}$'s to be 0 equivalently. In this way, the QFP problem is approximately reduced to the following statement in terms of linear congruence equations (LCE). QFP and LCE are not necessarily equivalent. First of all, the vector \vec{e} obtained in QFP records information of edges in some graph and is generally not arbitrary. Second, we have intentionally simplified the problem by omitting the indication of places forced to be 0 in \vec{r} in LCE. This gives us a stronger statement than QFP, that is, the LCE problem implies QFP and any counterexample of QFP also invalidates LCE. Compared to the trivial linearization of QFP, we have actually used more variables to make the representation more symmetric.

Problem 5 (LCE) *If the following equation (Eq. (10)) of \vec{r} has a real solution,*

$$G\vec{r} \equiv 2T\vec{e} \pmod{4}$$

then it also has an integral solution that preserves all zero entries in the real solution.

3.2 The Low-Rank Case

In this subsection, we discuss the QFP problem with low-rank subspace S . Namely, we want to show that for $d \leq 5$, there are no counterexamples of QFP. As QFP is a sufficient condition for LU-LC, it follows that LU-LC conjecture holds for $d \leq 5$.

We need only to prove that there is no counterexamples for LCE in this case. First, any solution \vec{r} of Eq. (10) can be written as

$$\vec{r} = 2G^{-1}T\vec{e} + 4G^{-1}\vec{s},$$

for some integral vector \vec{s} , and vice versa. Substitute Eq. (14) into the second part,

$$\vec{r} = 2G^{-1}T\vec{e} + \frac{16}{D}G\vec{s} - \frac{8\sigma}{D}\vec{1},$$

where $\vec{1}$ is the vector whose entries are all 1 and σ is the summation of all entries in vector \vec{s} . When $d \leq 3$ all parts in the summation are integral. If $d = 4$, only the last term can be half-integral. However, when this happens, no entry in \vec{r} is integral, and the LCE problem holds by choosing $\vec{s} = 0$.

Consider now the case of $d = 5$. Let $[x]$ be the largest integer not exceeding x , and $\{x\}$ be $x - [x]$. We generalize this operation to vectors \vec{a} , obtaining the truncation $[\vec{a}]$. We claim that if \vec{r} is a solution of Eq. (10), the truncation $[\vec{r}]$ is also a valid solution. It suffices to prove that, for any integral \vec{s} ,

$$G[4G^{-1}\vec{s}] \equiv 0 \pmod{4}. \quad (15)$$

Employing Eq. (14) and a detailed discussion on the four possible cases of $\sigma \pmod{4}$, we can reduce the above equation to

$$G\left\{\frac{1}{2}G\vec{s}\right\} \equiv 0 \pmod{4},$$

which in turn follows easily from the second identity of Lemma 1 by noticing that

$$2 \left\{ \frac{1}{2} G \vec{s} \right\}$$

is a column of G .

We have shown that it is impossible to construct counterexamples of the QFP problem when the rank of subspace S is less than or equal to 5. Naturally, the next step is to investigate the case of $d = 6$, which will lead to counterexamples.

4 Random generation of counterexamples

Notice that the “zero-entry preserving” requirement is essential in LCE. Without this additional requirement, LCE will be true. Therefore, we try greedily to find a real solution of \vec{r} with as many zero entries as possible. As the number of nonzero entries will be the size of our counterexample, this approach also tries to minimize the size of counterexamples.

Recall the key equation of the problem

$$\vec{r} = 2G^{-1}T\vec{e} + 4G^{-1}\vec{s}.$$

As the first part of the right hand side is integral, we will at least need to find an integral vector \vec{s} so that $4G^{-1}\vec{s}$ has many integral entries in order to have an \vec{r} with many 0's. Conversely, once $4G^{-1}\vec{s}$ is chosen, we can try to cancel most of its integral entries by choosing \vec{e} . And after \vec{s} and \vec{e} are fixed, we can calculate \vec{r} and verify whether there is no integral solution that preserves zero entries in it. It is a backward approach that chooses in order, \vec{s} , \vec{e} , and \vec{r} . More details on this approach are given in the following.

We first choose \vec{s}_0 at random. But it generally does not give us many integral entries in $4G^{-1}\vec{s}_0$. We can employ the rounding technique once again previously used in the case of $d = 5$. For $d = 6$, the following equation similar to Eq. (15) can be proved

$$G \left[8G^{-1}\vec{s}_0 \right] \equiv 0 \pmod{8}.$$

Therefore, there exists an integral \vec{s}_1 such that

$$\frac{1}{2} \left[8G^{-1}\vec{s}_0 \right] \equiv 4G^{-1}\vec{s}_1 \pmod{4}.$$

There will be generally more integers in $4G^{-1}\vec{s}_1$ than in $4G^{-1}\vec{s}_0$ and we can use \vec{s}_1 instead of \vec{s}_0 .

In the next step, vector \vec{e} is chosen to cancel the integral entries in $4G^{-1}\vec{s}_1$. Thanks to the simple form of $2G^{-1}T$, it is flexible to find \vec{e} that fulfills the requirement and, in addition, satisfies that $E_{mm'} = 0$ when pattern m corresponds to an integral entry in $4G^{-1}\vec{s}_1$.

Vector \vec{e} gives most of the useful information on the quadratic form $Q(x)$, and we construct subspace S by deleting columns of G corresponding to zero entry positions in \vec{r} . This is how a candidate counterexample is generated.

Finally, we need to verify whether the $Q(x)$ and S obtained here constitute a valid counterexample of QFP, that is, to verify that there do not exist $c_j \in \{\pm 1, \pm i\}$ satisfying Eq. (2). This can be done by a modified Gaussian elimination method described below.

Let A be the matrix that contains all vectors in S as its rows. The problem of whether c_i can all be chosen from $\{\pm 1, \pm i\}$ is equivalent to whether an equation of the following form has an integral solution

$$Ax \equiv b \pmod{4}. \quad (16)$$

The main difficulty in performing Gaussian elimination is that in congruence equations we cannot perform division to normalize the pivot. Fortunately, for equations of modular 4, division operations can be avoided. When performing the elimination procedure, try first to find a row with odd pivot and if the pivot is congruent to 3, multiply 3 on it. Otherwise, if we cannot find a row having odd pivot, the division is also unnecessary as one can do elimination simply by subtraction. Using this modified elimination method, we can derive from Eq. (16) a contradicting equation as

$$0 \equiv 2 \pmod{4}, \quad (17)$$

under the assumption of the existence of an integral solution. It is worth noting that after this work was done, the authors are aware that there is a more straightforward and general method for solving linear congruence equations based on the Smith normal form and the elementary divisors [20] of integral matrices. The Smith normal form method may be helpful in further understanding the structure of the counterexamples of LU-LC.

The above procedure of randomly generating different counterexamples is implemented and is available at <http://arxiv.org/e-print/0709.1266> as a gzipped tar (.tar.gz) file. Here is one of the counterexamples it generates.

In the example, $n = 27$ and the subspace S of \mathbb{F}_2^{27} is given by a set of basis as

$$\begin{aligned} \xi^1 &= 1000100010101010100011110 \\ \xi^2 &= 101010111001100000001010101 \\ \xi^3 &= 011001100111100111100110011 \\ \xi^4 &= 000111100000011001100001111 \\ \xi^5 &= 000000011111111000011111111 \\ \xi^6 &= 000000000000000111111111111. \end{aligned} \quad (18)$$

One can see that every two columns (patterns) are different and the columns are organized in increasing lexicographic order. S has a rank of 6 and consists 64 elements.

The quadratic form $Q(x)$ is the summation of 11 terms:

$$x_1x_2 + x_1x_3 + x_1x_8 + x_2x_4 + x_2x_8 + x_2x_{16} + x_3x_4 + x_3x_8 + x_3x_{16} + x_4x_8 + x_8x_{16}. \quad (19)$$

We note that this is not directly generated by the random procedure. In fact, we have found a Q with smaller number of terms that has the same value $Q(x)$ as the generated quadratic form for all $x \in S$. Originally, the quadratic form contains more than 60 terms. This simplification is suggested by one of the referees of QIP 2008.

The QFP problem defined by the above Q and S has a solution where the phases are powers of $e^{\pi i/4}$, but not in $\{\pm 1, \pm i\}$. We give the exponents sequentially:

$$e = [3, 5, 7, 5, 1, 3, 5, 7, 1, 3, 5, 5, 3, 7, 3, 3, 7, 1, 7, 3, 1, 5, 5, 5, 3, 5, 3]. \quad (20)$$

Our final aim is to disprove the LU-LC conjecture and what we have already shown is that QFP is false generally. As proved in Ref. [9], QFP implies LU-LC. Therefore, QFP is false does not logically guarantee that LU-LC is false. Fortunately, however, the above counterexample of QFP can be transformed to a counterexample of LU-LC. Although not proved, this transformation seems to work all the time.

Here is the way we perform the transformation. Define two states

$$\begin{aligned} |S\rangle &= \sum_{x \in S} |x\rangle \\ |Q, S\rangle &= \sum_{x \in S} (-1)^{Q(x)} |x\rangle. \end{aligned} \tag{21}$$

From Eq. (1), we know that they are stabilizer states. The corresponding QFP problem of the same Q and S indicates that they are LU equivalent, in fact, even DLU equivalent. We need to show that they are not LC equivalent. This can be done efficiently by using the LC equivalence decision algorithm [16] for graph states after efficiently finding graph states $|G_S\rangle$ and $|G_{Q,S}\rangle$ which are LC equivalent to $|S\rangle$ and $|Q, S\rangle$ respectively. If the algorithm tells that $|G_S\rangle$ and $|G_{Q,S}\rangle$ are not LC equivalent, nor are the states $|S\rangle$ and $|Q, S\rangle$ (see Fig. 2). The four states in the diagram are in the same equivalence class under LU criterion. But when LC equivalence is considered, the upper two and the lower two belong to different classes. To implement the above procedure, we have used a computer program and the code is also available in <http://arxiv.org/e-print/0709.1266>.

$$\begin{array}{ccc} |Q, S\rangle = \frac{1}{\sqrt{|S|}} \sum_{x \in S} (-1)^{Q(x)} |x\rangle & \xleftrightarrow{\text{LC}} & |G_{Q,S}\rangle \\ & \updownarrow \text{LU} & \updownarrow \text{NOT LC} \\ |S\rangle = \frac{1}{\sqrt{|S|}} \sum_{x \in S} |x\rangle & \xleftrightarrow{\text{LC}} & |G_S\rangle \end{array}$$

Fig. 2. State Transformation

A better solution to show the non-LC equivalence of $|Q, S\rangle$ and $|S\rangle$ would be a proof that DLU and LC imply DLC. This is open generally, but it is pointed out by B. Zeng that when local unitaries are all non-Clifford, just as in the above case, a proof can be found using results on minimal support of stabilizers developed in Ref. [19].

The two graph states $|G_S\rangle$ and $|G_{Q,S}\rangle$ have corresponding graphs shown in Fig. 3. The graphs with and without the dotted edge are graph $G_{Q,S}$ and G_S respectively. Interestingly, two LU equivalent but not LC equivalent graph states can differ only in one edge. Note that the simplification of $Q(x)$ discussed above does not simplify or change the form of the four stabilizer states in Fig. 2.

5 Conclusion

In summary, we have shown that the LU-LC conjecture is false by giving an explicit counterexample of it. It is also clear that in order to disprove the conjecture, we have to consider stabilizer states with the rank of the support no less than 6. This result leads us to rethink




Fig. 3. Corresponding Graph States

about the local equivalence problem of stabilizers as most of the previous work focus on proving the conjecture to be true.

The random procedure generates counterexamples of size 27 and 35. Though not proved, we believe that 27 is the smallest possible size of counterexamples of LU-LC. Although the LU-LC conjecture is now disproved, we still know little about the relation of LU and LC equivalence and do not have an explicit understanding of why or when LU equivalence differs from LC equivalence. When $d = 6$, the random generation procedure can find a counterexample in seconds, but it is in fact not an efficient algorithm when d is large or even when $d = 7$. That fact is we have never obtained a valid counterexample of $d = 7$ using the random procedure. Fortunately, larger scale counterexamples, including those of $d = 7$, have been found [8] motivated by the randomly generated counterexamples.

As LU and LC equivalences are now known to be different, it is challenging to ask whether there is an efficient algorithm deciding LU equivalence for stabilizers or whether there is a graph theoretical interpretation of LU equivalent graph states. This seems to be difficult as local unitary operations are much less linked up with the stabilizer formalism.

Acknowledgement

We are thankful to David Gross, Maarten van den Nest, Bei Zeng, Markus Grassl and the colleagues in the Quantum Computation and Information Research Group of Tsinghua for helpful discussions. We note that a gap in an earlier version of this paper was found by Andre Ahlbrecht and communicated to the authors by David Gross. This work was partly supported by the National Natural Science Foundation of China (Grant Nos. 60736011, 60721061 and 60621062).

References

1. K. Chen and H.-K. Lo. Multi-partite quantum cryptographic protocols with noisy GHZ states, 2004. arXiv:quant-ph/0404133.
2. L. E. Danielsen. On self-dual quantum codes, graphs, and boolean functions, 2005. arXiv:quant-ph/0503236.
3. L. E. Danielsen and M. G. Parker. On the classification of all self-dual additive codes over $\text{GF}(4)$ of length up to 12. *J. Comb. Theory Ser. A*, 113(7):1351–1367, 2006.
4. V. Danos, E. Kashefi, and P. Panangaden. The measurement calculus. *J. ACM*, 54(2):8, 2007.
5. J. Dehaene and B. D. Moor. Clifford group, stabilizer states, and linear and quadratic operations over $\text{GF}(2)$. *Physical Review A (Atomic, Molecular, and Optical Physics)*, 68(4):042318, 2003.
6. D. Gottesman. *Stabilizer Codes and Quantum Error Correction*. PhD thesis, Caltech, 1997. arXiv:quant-ph/9705052.
7. M. Grassl, A. Klappenecker, and M. Roetteler. Graphs, quadratic forms, and quantum codes. In *Proceedings 2002 IEEE International Symposium on Information Theory (ISIT 2002)*, 2002. arXiv:quant-ph/0703112.
8. M. Grassl and B. Zeng. Personal communication.
9. D. Gross and M. van den Nest. The LU-LC conjecture, diagonal local operations and quadratic forms over $\text{GF}(2)$. *Quantum Information and Computation*, 8:263–281, 2008. arXiv:0707.4000.
10. M. Hein, W. Dur, J. Eisert, R. Raussendorf, M. van den Nest, and H. J. Briegel. Entanglement in graph states and its applications, 2006. arXiv:quant-ph/0602096.
11. M. Hein, J. Eisert, and H. J. Briegel. Multiparty entanglement in graph states. *Physical Review A (Atomic, Molecular, and Optical Physics)*, 69(6):062311, 2004.
12. R. Raussendorf and H. J. Briegel. A one-way quantum computer. *Physical Review Letters*, 86(22):5188–5191, 2001.
13. D. Schlingemann. Stabilizer codes can be realized as graph codes, 2001. arXiv:quant-ph/0111080.
14. D. Schlingemann. Local equivalence of graph states, 2005. in O. Krueger and R. F. Werner, Some Open Problems in Quantum Information Theory, arXiv:quant-ph/0504166. See also <http://www.imaph.tu-bs.de/qi/problems/>.
15. M. van den Nest, J. Dehaene, and B. de Moor. Local equivalence of stabilizer states. In *Sixteenth International Symposium on Mathematical Theory of Networks and Systems (MTNS2004)*, Katholieke Universiteit Leuven, Belgium, 2004.
16. M. van den Nest, J. Dehaene, and B. D. Moor. Efficient algorithm to recognize the local clifford equivalence of graph states. *Physical Review A (Atomic, Molecular, and Optical Physics)*, 70(3):034302, 2004.
17. M. van den Nest, J. Dehaene, and B. D. Moor. Graphical description of the action of local clifford transformations on graph states. *Physical Review A (Atomic, Molecular, and Optical Physics)*, 69(2):022316, 2004.
18. M. van den Nest, J. Dehaene, and B. D. Moor. Invariants of the local clifford group. *Physical Review A (Atomic, Molecular, and Optical Physics)*, 71(2):022310, 2005.
19. M. van den Nest, J. Dehaene, and B. D. Moor. Local unitary versus local clifford equivalence of stabilizer states. *Physical Review A (Atomic, Molecular, and Optical Physics)*, 71(6):062323, 2005.
20. B. L. van der Waerden. *Algebra, Volume 2*. Frederick Ungar Publishing Co., Inc., New York, 1970.
21. B. Zeng, H. Chung, A. W. Cross, and I. L. Chuang. Local unitary versus local clifford equivalence of stabilizer and graph states. *Physical Review A (Atomic, Molecular, and Optical Physics)*, 75(3):032325, 2007.
22. B. Zeng, A. Cross, and I. L. Chuang. Transversality versus universality for additive quantum codes, 2007. arXiv:0706.1382.