

# **Arsenic Accumulation in Rice (*Oryza sativa* L.); Human Exposure through Food Chain**

**M. Azizur Rahman\*<sup>1</sup>; H. Hasegawa<sup>1</sup>; M. Mahfuzur Rahman<sup>2</sup> and M. A. Mazid Miah<sup>3</sup>;  
A. Tasmen<sup>4</sup>**

<sup>1</sup>Graduate School of Natural Science & Technology, Kanazawa University, Kakuma,  
Kanazawa 920-1192, Japan; <sup>2</sup>Faculty of Biological Sciences, Department of Botany,  
Jahangirnagar University, Savar, Dhaka-1342, Bangladesh ; <sup>3</sup> Division of soil Science,  
Bangladesh Rice Research Institute (BRRI), Gazipur, Bangladesh. <sup>4</sup>Institute of Research  
and Education (IER), University of Dhaka, Dhaka-1000, Bangladesh

\*Corresponding author

Tel: 76-265-3132

E-mail: arahmsn@stu.kanazawa-u.ac.jp

aziz\_ju@yahoo.com

**Abstract**

Although human exposure to arsenic is sought to be caused mainly through arsenic contaminated underground drinking water, the use of this water for irrigation enhances the possibility of arsenic uptake into crop plants. Rice is the staple food grain all over Bangladesh. As such arsenic content in straw, grain and husk of rice is especially important since paddy fields are extensively irrigated with underground water having high level of arsenic concentration. On the contrary, straw and husk are widely used as cattle feed. Arsenic concentration in rice grain was  $0.5 \pm 0.02 \text{ mg kg}^{-1}$  with the highest concentrations being in grains grown on soil treated with  $40 \text{ mg As kg}^{-1}$  soil. With the average rice consumption between 400 and 650 g/day by typical adults of the arsenic affected areas of Bangladesh, the intake of arsenic through rice stood at 0.20 to 0.35 mg/day when with a daily consumption of 4 L, arsenic intake through drinking water was 0.2 mg/day. Moreover, when the rice plant was grown in  $60 \text{ mg of As kg}^{-1}$  soil, arsenic concentrations in rice straw were  $20.6 \pm 0.52$  at panicle initiation stage and  $23.7 \pm 0.44$  at maturity stage while it was  $1.6 \pm 0.20 \text{ mg kg}^{-1}$  in husk. Cattle drink a large amount of water. So alike human beings, arsenic gets deposited into cattle body through rice straw and husk as well as from drinking water which in turn finds route into human body. Arsenic intake in human body from rice and cattle could be potential in addition to that from drinking water. Therefore, a hypothesis has been put forward elucidating the possible food chain pathways through which arsenic may enter into human body.

**Key words:** Arsenic, Rice, Toxicity, Food chain, Human exposure

## **Funding Sources:**

The research was supported by the NSICT fellowship of the Ministry of Science, Information and Communication Technology, Government of the People's Republic of Bangladesh for the young scientists. The facilities for glasshouse experiments and chemical analysis were provided by the Soil Science Division, Bangladesh Rice Research Institute (BRRI), Gazipur, Bangladesh.

## **Introduction**

Arsenic contamination in ground water has turned into the gravest natural disaster with spatial extent encompassing Bangladesh, India (West Bengal), China, Taiwan, Vietnam, United States of America, Argentina, Chile, Mexico etc. In Bangladesh, arsenic concentration in ground water has exceeded the safe level ( $0.05 \text{ mg As L}^{-1}$  of water is the Bangladesh standard) in 59 districts out of 64 districts and about 80 million people are exposed to arsenic poisoning. The natural contamination of shallow hand tube wells in Bangladesh with arsenic has caused widespread human exposure to this toxic element through drinking water (Karim, 2000; Paul et al., 2000). Use of arsenic-contaminated shallow tube-well water for irrigation of crops has put forward the question - is arsenic contaminated drinking water the only pathway of human exposure to arsenic? If not, what are the other pathways through which such exposure is taking place? With this question in mind, we conducted glasshouse and field level experiments to investigate the concentrations of arsenic in rice, the main food stuff of Bangladeshis, and straw and husk of rice, the main fodder for cattle in the country.

The impact of arsenic contaminated irrigation water on the arsenic content in rice is especially important as rice is the staple food for the population of arsenic epidemic areas and it is grown in flooded (reduced) condition where arsenic availability is high (Duxbury et al., 2003). Different consumers of natural ecosystem, such as primary, secondary or tertiary,

are taking arsenic contaminated food and water and as manifested by reports - arsenic is getting deposited into their bodies (Bruce et al., 2003; Shariatpanahi and Anderson, 1984; Thornton and Webb, 1979).

Another important aspect of the present study is the extent and severity of arsenic poisoning in human body through these crop plants, directly or indirectly. We tried to trace food chain pathways of natural ecosystem through which arsenic may enter into human body so that we can assess the potentiality of these pathways in exposing human to arsenic. It is quite difficult to investigate all the arsenic transferring food chain pathways of natural ecosystem even in small scale. So in this paper, we focused mainly on the extent and severity of arsenic poisoning in human body through “Plant (rice)-Animal (cattle)-Man” food chain pathway.

## **Materials and Methods**

### **Soil Preparation**

Pot experiments were conducted in a glasshouse at Bangladesh Rice Research Institute (BRRI). Soil, collected from BRRI farm at a depth of 0-15cm, were sun dried for 7 days and then the massive aggregates were broken down by gentle crushing with hammer. The unwanted materials viz. dry roots, grasses, stones were removed from the bulk soil. Then the soil was mixed thoroughly, crushed and sieved with 2 mm sieve. Sample from this initial soil was collected into a plastic bottle for physico-chemical analysis.

### **Pot Preparation**

Five kilogram soil was taken in six liter plastic pots which were used to avoid leaching and to protect absorption of water soluble arsenic from the soil. Before taking the soils into them, all plastic pots were washed by tap water and sun dried. There were altogether 30 pots

comprising ten arsenic treatments with three replications for each. The pots were arranged following the factorial Randomized Complete Block Design (RCBD).

### **Arsenic Treatment**

The arsenic concentration in agricultural soil of arsenic affected areas of Bangladesh is between 20 and 90 mg kg<sup>-1</sup> (Ullah, 1998). Therefore, arsenic was mixed thoroughly with the soil at the rate of 0 (control), 10, 20, 30, 40, 50, 60, 70, 80 and 90 mg As kg<sup>-1</sup> soil. After application of arsenic as aqueous solution of sodium arsenate, the spiked soils were left for two days without irrigation. Before transplantation each pot was irrigated by 4.5 L of water having an arsenic concentration of 0.01 mg l<sup>-1</sup>.

### **Intercultural Operation and Fertilizer Application**

The pots were placed on a plane cemented table. The overall temperature in the glasshouse ranged from 22.4 to 33.9°C, relative humidity from 59.9 to 83.7%, average evaporation from 3.8 to 6.0 mm, sunshine from 3.4 to 7.8 h/day. BRRI dhan26 (Rice) was used as test crop. Four 35-days old seedlings were transplanted in each pot at equal spacing. After transplantation, the rice plants were grown under flooded condition. Pots were irrigated regularly, maintaining a water depth of 3 cm, throughout the post-transplantation period until harvesting. Urea, Triple Super Phosphate (TSP) and Muriate of Potash (MP) were applied at the rate of 30, 40 and 20 kg per hectare for nitrogen, phosphorous and potassium, respectively in the spiked soil. One-third of the amount of urea and full amount of other two fertilizers were applied as basal in the individual pot before transplantation. The fertilizers were incorporated with the soil by hand. The second and third splits of urea were applied after 30 (maximum tillering stage) and 60 days (panicle initiation stage) of transplantation.

### **Physico-chemical Properties of Initial Soil**

The physico-chemical properties of initial soil were measured to determine its fertility and behavior since the toxicity limit and mobility of arsenic are function of the physico-chemical properties of soil such as particle size, texture, soil reaction, mineral nutrient content etc. In sandy soil, arsenic is more mobile and bio-available than that of clayey soil. The availability of the arsenic in soils is affected by changes in pH. In general there is an increase in arsenic toxicity effects on plants, as the soil become acidic, particularly at pH below 5. However, the uptake of arsenic by plants may be increased on higher pH soil (Campbell et al., 1985). Phosphate has been reported to displace arsenic from soil (Peryea, 1991). Heavy additions of P to arsenic polluted soils have been reported to displace approximately 77% of total arsenic in the soil. Therefore, it is relevant to know about these soil properties to evaluate the influence of arsenic on its accumulation in rice as well as on soil properties. The results of the physico-chemical properties of initial soil have been presented in Table 1 and 2.

**Physical Properties:** Physical properties of soil such as distribution of particle size, textural classes, moisture content were determined and are presented in Table 1. The soil was Silty-clay-loam (Sand 12.30%, Silt 53.00% and Clay 34.70%) and blackish in color. The moisture content of the soil was 16.04%.

**Chemical Properties:** Chemical properties of soil such as pH, organic carbon, organic matter, total nitrogen, total phosphorus, total potassium, total iron, total arsenic, available phosphorus and available iron were determined and the results are presented in Table 2. The initial soil was acidic (pH 5.27) in nature. The background arsenic of the experimental soil was 3.25 mg kg<sup>-1</sup>. The soil was rich in iron with available iron of 48.02 mg kg<sup>-1</sup>. Organic carbon and organic matter was about 0.77 and 1.32%, respectively. Total nutrients such as nitrogen, phosphorus and potassium in soils were not sufficient (0.25, 0.02 and 0.12%,

respectively). Fertilizers of these nutrients elements were applied to reduce their deficiency. Available phosphorus was about 6.15% in the soil.

### **Collection of Field Samples**

Rice (*Oryza sativa* L.) samples (1 kg) of two varieties (BRRI dhan28 and BRRI hybrid dhan1) were collected from a severely arsenic affected area of Bangladesh with three replications. Soil samples (1 kg) were also collected from 2m<sup>2</sup> areas and 10-15 cm depth of the selected plots using soil auger. Samples were collected during harvest and sun dried immediately after collection, tagged properly, air tied in polyethylene bags and kept in room temperature for farther laboratory analysis.

### **Chemical Analysis for Arsenic**

The plant samples (straw, grain and husk) were digested with concentrated nitric acid and perchloric acid. 0.5 g of the sample was taken into a dry clean digestion tube and 5 ml of 65% nitric acid was added. The mixture was allowed to stand over night under fume shade. In the following day, the digestion tubes were placed on a heating block and the temperature was raised to 60°C. After heating for about 1 h, the tubes were allowed to cool and 2 ml of concentrated perchloric acid was added. Again, the tubes were heated at 160°C. Heating was stopped when the dense white fumes of perchloric acid occurred. The digests were cooled, diluted in 25 ml distilled deionized water and filtered into plastic bottles through filter paper (Whatman, No.1). Total arsenic was determined by hydride generation atomic absorption spectrophotometer (HG-AAS) using matrix-matched standards ([Welsch et al., 1990](#)). All glassware and plastic bottles were previously washed by distilled deionized water and dried.

All instruments were calibrated using matrix-matched standards. In each analytical batch at least two reagent blanks, one spike and three duplicate samples were included in the acid

digests to assess the accuracy of the chemical analysis. Accuracy of the method, according to the spike, was  $92.3 \pm 1.5\%$ .

### **Chemicals**

Nitric acid ( $\text{HNO}_3$ ), Sulfuric acid ( $\text{H}_2\text{SO}_4$ ), Perchloric acid ( $\text{HClO}_4$ ) and Sodium arsenate ( $\text{Na}_2\text{HAsO}_4 \cdot 7\text{H}_2\text{O}$ ) were purchased from Merck. Other chemicals were from AnalaR. All the reagents were of analytical grade.

### **Statistical Analysis**

Duncan's Multiple Range Test (DMRT) was computed at 5% level to see the significant differences among the treatments and Pearson correlation was estimated by SPSS 10.0 for windows.

## **Results and Discussion**

To investigate the potential of "plant-human" food chain pathway in arsenic poisoning of human body, we determined the arsenic concentration in tissues of rice. A hypothesis may also be demonstrated from it reflecting the possibility of arsenic poisoning of human body through different food chain pathways, especially the "Plant-Animal-Man", on the basis of data of "plant-human" food chain pathway. In the first phase of this experiment, rice was cultivated in artificially spiked soil with different levels of arsenic in a glasshouse and the results were compared with that of field data.

### **Arsenic Concentration in Rice Tissues**

#### **Glasshouse study**

Rice plants were grown in arsenic-spiked soils to determine the arsenic concentration in tissues of these plants. Pearson correlation analysis revealed that arsenic concentrations in



rice straw increased significantly ( $r = 0.781$  and  $0.852$ ;  $p = 0.013$  and  $0.002$  for straw of PI stage and maturity stage, respectively) with the increase of soil arsenic concentrations (Figure 2). In  $60 \text{ mg of As kg}^{-1}$  soil treatment, arsenic contents in straw were 34.33- and 26.33- fold higher than that of control. However, correlation analysis also showed that arsenic contents in husk and grain for different soil arsenic treatments did not differ significantly ( $r = 0.358$  and  $0.014$ ;  $p = 0.344$  and  $0.970$  for husk and grain, respectively) from each other (Figure 3). In  $60 \text{ mg of As kg}^{-1}$  soil treatment, husk arsenic content was 8-fold higher than that of control and it was 2.5-fold higher for grain in  $40 \text{ mg of As kg}^{-1}$  soil treatment. In 70, 80 and 90 mg of  $\text{As kg}^{-1}$  soil treatments, arsenic contents in husk and grain were less (but not significant) than those of other arsenic treatments. The results indicate that at higher levels of soil arsenic concentrations, the toxic element causes severe toxicity to rice plant resulting reduced growth rate and lowered translocation of arsenic as well as other nutrients from soil solution into the rice grain. Regardless of soil arsenic concentrations, arsenic concentration in rice tissues followed the trend: straw > husk > grain.

Both the fresh and dried rice straw has been used widely as fodder for cattle in arsenic affected areas like Bangladesh and west Bengal, India. Therefore, arsenic concentrations were measured in rice straw at both panicle initiation (PI) stage and maturity stage (after harvest). Results imply that soil arsenic concentrations influenced its contents in straw of both stages. In straw of PI stage, the highest arsenic content was  $20.6 \pm 0.52 \text{ mg kg}^{-1}$  dry weight at  $60 \text{ mg of As kg}^{-1}$  soil treatment. In 70 and 80 mg of  $\text{As kg}^{-1}$  soil treatments, arsenic contents were less than that of in  $60 \text{ mg of As kg}^{-1}$  soil treatment, possibly, because of reduced translocation of arsenic as well as other nutrients resulted from severe toxicity of this metal to rice plant (Table 3).

In straw of mature stage (after harvest), arsenic content followed the same order of magnitude as in straw of PI stage (Figure 2). However, the highest straw arsenic content

was  $23.7 \pm 0.44 \text{ mg kg}^{-1}$  dry weight in  $60 \text{ mg of As kg}^{-1}$  soil treatment. [Abedin et al. \(2002a\)](#) also reported significant increase of arsenic concentration in rice root, straw and husk with the increase of arsenate concentration in irrigation water. He found  $3.9 \text{ mg kg}^{-1}$  arsenic in straw at the lowest arsenate treatment ( $0.2 \text{ mg l}^{-1}$ ), which increased progressively with increasing arsenate application and reached to  $91.8 \text{ mg kg}^{-1}$  in the highest arsenate treatment ( $8.0 \text{ mg l}^{-1}$ ). Arsenic uptake by plants is a function of plant species ([Liebig, 1966](#)), arsenic concentration in soil ([NRC, 2001](#)), pH and clay content ([Johnson and Hiltbold, 1969](#)), other ions ([Woolson et al., 1973](#), [Khattak et al., 1991](#)) and the chemical form of arsenic ([Marin et al., 2003](#)).

Arsenic contents in rice grain were not significantly ( $p > 0.05$ ) influenced by the soil arsenic concentrations. The highest grain arsenic content was  $0.5 \pm 0.02 \text{ mg kg}^{-1}$  dry weight in  $40 \text{ mg of As kg}^{-1}$  soil treatment and the lowest was  $0.2 \pm 0.01 \text{ mg kg}^{-1}$  dry weight in control and  $90 \text{ mg As kg}^{-1}$  soil treatment (Figure 3). [Abedin et al. \(2002a\)](#) also reported that arsenic concentration in grain remained statistically indifferent with increasing arsenate concentration in irrigation water. He found  $0.15 \text{ mg As kg}^{-1}$  dry weight in grain at control treatment and  $0.24 \text{ mg As kg}^{-1}$  dry weight at  $4.0 \text{ mg of As L}^{-1}$  water treatment. [Williams et al. \(2005\)](#) also reported mean arsenic concentration of  $0.26 \mu\text{g g}^{-1}$  in US long grain rice and  $0.40 \mu\text{g g}^{-1}$  as the highest grain arsenic concentration. They also found mean arsenic concentration of  $0.13 \mu\text{g g}^{-1}$  in Bangladesh rice grain.

### **Field Study**

To get real scenario of arsenic concentrations in rice tissues, we also did a field level investigation. Rice samples were collected directly from the field and measured the concentrations of arsenic in rice tissues. The results of field investigation are presented in figure 4. Arsenic concentration in field soil was  $14.51 \pm 0.21 \text{ mg kg}^{-1}$  and  $0.07 \pm 0.02 \text{ mg L}^{-1}$  in water. Arsenic concentrations in straw, husk and grain of two rice strains (BRRI dhan28

and BRR1 hybrid dhan1) did not differ significantly from each other ( $p>0.05$ ). Arsenic concentration in husk of BRR1 hybrid dhan1 contains 3.8-fold higher than that of rice grain while it was 3.33-fold for BRR1 dhan28.

In glasshouse experiment, arsenic concentrations in rice straw, husk grain were  $2.09\pm 0.09$ ,  $0.27\pm 0.05$  and  $0.25\pm 0.06$  mg kg<sup>-1</sup> dry weight, respectively when the soil arsenic concentration was 13.25 mg kg<sup>-1</sup> (10 mg kg<sup>-1</sup> was spiked arsenic and 3.25 mg kg<sup>-1</sup> was background arsenic concentrations in the soil). In contrary, when the arsenic concentration in the field soil was  $14.51\pm 0.21$  mg kg<sup>-1</sup>, its concentrations in rice straw, husk and grain were  $1.78\pm 0.11$ ,  $1.36\pm 0.01$  and  $0.41\pm 0.01$  mg kg<sup>-1</sup> dry weight. The results revealed that arsenic concentrations in husk and grain of field samples were higher than that of glasshouse samples at the almost same soil arsenic concentration. This may be because the phosphate concentration in glasshouse soil was higher than that of field soil (Table 2) and the phosphate suppresses arsenic uptake in rice plant (Abedin et al., 2002b).

### **Human Exposure to Arsenic through “Plant-Animal-Man” Food Chain Pathway**

It is clear from the present experiment and some other previous reports that arsenic deposits in tissues of crop plants grown in arsenic rich soil, irrigated with arsenic contaminated water. Arsenic accumulation has been reported in maize (Sadiq, 1986), barley and ryegrass (Jiang and Singh, 1994), rice (Duxbury et al., 2003; Abedin et al., 2002a; Marin et al., 1992; Bae et al., 2002; Onken and Hosner, 1995; Rahman et al., 2004; D’lilio et al., 2002), *Spartina alterniflora* (Carbonell et al., 1998) too. The accumulation of arsenic in plants occurs primarily through the root system and the highest arsenic concentrations have been reported in plant roots and tubers (Anastasia and Kender, 1973; Marin et al., 2003). Therefore, tuber crops are expected to have higher arsenic contents than that of other crops when those are grown in arsenic contaminated soil. The concentration of arsenic in edible parts of most plants is generally low (Vaughan, 1993; O’Neil, 1995). Plants seldom

accumulate arsenic at concentrations hazardous to human and animal health because, phytotoxicity usually occurs before such concentrations are reached (Walsh and Keeney, 1975).

Although human may be exposed to arsenic from a variety of environmental sources, food constitutes the largest source of arsenic intake with smaller contribution from air and drinking water (Chen and Lin, 1994). In a tropical country like Bangladesh, water consumption is normally very high. Most of the arsenic affected areas are villages where people are involved in agrarian manual labor. Daily water consumption by an adult ranged between 4 and 6 liters (Farmer and Johnson, 1990) and when the arsenic concentration in drinking water is  $0.05 \text{ mg L}^{-1}$ , the acceptable limit for drinking water in Bangladesh (though in many areas, arsenic concentrations in drinking water has been found to be more than this), an adult is expected to intake 0.2 to 0.3 mg of As/day from drinking water. In contrary, the average daily rice consumption by an adult of this area is between 400 and 650 g raw rice grain (Duxbury et al., 2003). In the present study, arsenic concentrations in rice grain were  $0.5 \pm 0.02$  and  $0.41 \pm 0.01 \text{ mg kg}^{-1}$  dry weight for glasshouse and field sample, respectively when the soil arsenic concentrations were 40 and  $14.51 \pm 0.21 \text{ mg kg}^{-1}$  soil, respectively. In Bangladesh, the soil arsenic concentration has been found to be between 20 and  $90 \text{ mg kg}^{-1}$  (Ullah, 1998).

The daily intake of arsenic from rice grain containing  $0.5 \pm 0.02 \text{ mg kg}^{-1}$  dry weight would be between 0.20 and 0.35 mg (according to the glasshouse data) and between 0.164 and 0.266 mg (according to the field data). Bae et al. (2003) reported that the concentration of arsenic in cooked rice was higher than that of raw rice. Rahman et al. (2006) reported elevated concentrations of arsenic in cooked rice when the rice was cooked with arsenic contaminated water and the gruel was not discarded after cooking. This was because the arsenic in water was absorbed by cooked rice. Ackerman et al. (2005) found 89 - 105% absorption of arsenic by rice from total volume of water [1:1 to 4:1 (water: rice)] used in

cooking for two different contaminated drinking water. Moreover, most of the arsenic in drinking water is dissolved as toxic inorganic forms, while the species of arsenic in raw and cooked rice are poorly characterized (Duxbury et al., 2003). Schoof et al. (1999) reported that between 30 and 85% of arsenic in rice is inorganic. These reports suggest that intake of arsenic from rice and its potential to human exposure should not be ignored.

The highest arsenic concentration in straw is  $23.7 \pm 0.44$  mg kg<sup>-1</sup> dry weight at 60 mg of As kg<sup>-1</sup> soil treatment while it was  $12.3 \pm 0.03$  mg kg<sup>-1</sup> at 40 mg of As kg<sup>-1</sup> soil treatment. Tsutsumi et al. (1980) reported 149 mg of As kg<sup>-1</sup> dry weight in rice straw when soil arsenic concentration was 313 mg kg<sup>-1</sup>. Abedin et al. (2002a) found 25 mg of As kg<sup>-1</sup> dry weight in rice straw when the plant was irrigated by 2 mg of As l<sup>-1</sup> water. Cattle are one of the primary consumers of terrestrial ecosystem. They feed on rice straw and husk and drink water as well. Though there is no direct report of arsenic accumulation in cattle body from rice straw or husk, the consequence of exposure to this toxic element in organs such as the liver and kidneys of this animal is well reported (WHO, 2001). Bruce et al., (2003) reported arsenic accumulation in liver and other tissues of tailing paddock animals though the accumulation was insignificant to cause chronic toxicity or any immediate perceivable contamination. Because Bruce et al., (2003) conducted their experiment for a short time (240 days); they expected more accumulation of arsenic in cattle if the experiment were for longer time. However, the objective of the present experiment was not to calculate the acute toxicity or sub chronic dose rates for cattle rather to justify the accumulation and transfer of arsenic from cattle to human through food chain.

Straw given to cattle in U.K. contained less than 0.20 mg As kg<sup>-1</sup> (Nicholson et al., 1999), though arsenic metabolized by the cattle is dependent on the arsenic species in the straw and on the metabolism of cattle (Abedin et al., 2002b). In another experiment, Shariatpanahi et al. (1984) reported that, sheep those were feed on methylarsonate showed a significant increase of arsenic accumulation in their tissues and milk. Although there have not been

found adequate data on the presence of arsenic in milk and meat of the cattle of Bangladesh and those imported from west Bengal, India (another arsenic epidemic area, where arsenic contamination in ground water is alarming), there is an ample scope of arsenic deposition in cattle body, especially from high arsenic-containing rice straw and husk. Thus, a hypothesis has been put forward elucidating the possible deposition of arsenic in human body not only be from drinking water but also from beef and mutton through “Plant-Animal-Man” and some other food chain pathways (Figure 1). All studies suggest that the possible health risk of human being from arsenic toxicity through “Plant-Animal-Man” food chain pathway should not be ignored. Moreover, when the arsenic contaminated straw is burned as fuel, arsenic may pollute the air as arsenic oxides and inhaled by man.

### **Arsenic Transfer through Food Chain**

The pattern of arsenic accumulation and its transfer from one trophic level to another is important. In the present article, we discussed this aspect by some previous data and the results were incorporated with our proposed hypothesis. [Mason et al. \(2000\)](#) reported a decrease of arsenic levels with the increase of higher trophic level. He also suggested that the subsequent transfer of arsenic to higher trophic levels is related to both the ability of the organisms to depurate and the mode of accumulation, either directly from water or from foodstuffs. Total arsenic concentrations in organisms after accumulation from foodstuffs decreased one order of magnitude per elevation of the trophic level.

[Klose and Braun \(1997\)](#) studied the arsenic content in soil and uptake by crops including fodder plants, spring barley, potatoes, maize, winter rape, pasture grass and clover. In maize, rape, barley and potatoes, arsenic content ranged from 0.04 to 1.31 mg kg<sup>-1</sup> dry matter when grown on 60 – 362 mg of As kg<sup>-1</sup> soil. In experiment with pasture grasses, plant arsenic content ranged from 0.18 to 6.7 mg kg<sup>-1</sup> dry matter when the soil arsenic content ranged from 90 to 1050 mg kg<sup>-1</sup> soil. Limited reports are available on bioaccumulation of

arsenic in different consumers of trophic levels such as animals, insects, birds and also the men.

Because of low concentrations in terrestrial plants, arsenic accumulation in animals from this source is also low. Direct ingestion of arsenic from soil could be a major source of dietary arsenic for grazing livestock (Thornton et al., 1979). Bruce et al., (2003) also reported direct ingestion of arsenic from soil. It is estimated that about 1% of the arsenic in the soil was actually absorbed by the cattle, while the remaining being excreted directly. There have been different possible food chain pathways of natural ecosystem through which, human being (when considered as the topmost consumer of terrestrial ecosystem) may be exposed to arsenic toxicity (Figure 1).

## Conclusion

Many previous reports demonstrated that foodstuffs collected from arsenic epidemic areas contain significant concentrations of arsenic. Roychowdhury et al. (2002) reported the arsenic concentrations in individual composites of cooked items, collected from an arsenic epidemic area of West Bengal, India, as rice (between 374.17 and 666.57  $\mu\text{g kg}^{-1}$ ), freshwater fish (between 830 and 900  $\mu\text{g kg}^{-1}$ ), potato curry (186  $\mu\text{g kg}^{-1}$ ), potato skin fried in oil (617  $\mu\text{g kg}^{-1}$ ), leaf of vegetables (578  $\mu\text{g kg}^{-1}$ ), mixed vegetable (277.33  $\mu\text{g kg}^{-1}$ ), pulses (143  $\mu\text{g kg}^{-1}$ ). Das et al. (2004) reported arsenic concentrations exceeding the food safety limits in *Calocasia antiquorum* (between 0.09 and 3.99  $\text{mg kg}^{-1}$ ), potato (between 0.07 and 1.36  $\text{mg kg}^{-1}$ ), *Ipomoea reptoms* (between 0.1 and 1.53  $\text{mg kg}^{-1}$ ) collected from an arsenic epidemic area of Bangladesh. Arsenic deposition in cattle body (Bruce et al., 2003; Thornton et al., 1979) and tissues and milk of sheep (Shariatpanahi et al., 1984) has also been reported. Liao and Ling (2003) conducted an experiment on arsenic bioaccumulation in tilapia fish (*Oreochromis mossambicus*) and found that the highest ninety-fifth percentile of potential health risk for inorganic arsenic ranged from  $7.36 \times 10^{-4}$  to  $1.12 \times 10^{-3}$  for the

subsistence fishers of Blackfoot disease area of Taiwan. Thus, it is evident that not only “soil-water-human” but also “plant-human” and “plant-animal-human” may be other potential food chain pathways of arsenic accumulation in human body, though arsenic contaminated drinking water is the major and direct source. Adequate emphasis should be given on this matter. To figure out the fact regarding arsenic poisoning in human body through these food chain pathways, intensive investigation on a complete food chain is needed, which is our future interest.

### **Acknowledgement**

Authors are grateful to the Bangladesh Rice Research Institute (BRRI) authority for facilitating their Arsenic Laboratory, Soil Science Division, for conduction experiments. Authors are also thankful to Mr. Mosharraf Hossain for his sincere help in preparing this manuscript. The first author is thankful to the Ministry of Science, Information and Communication Technology, Government of the People’s Republic of Bangladesh, for awarding the NSICT fellowship for this research work.

### **References**

- (1) Abedin, M.J., Cresser, M.S., Meharg, A.A., Feldmann, J., Cotter-Howells, J., 2002a. Arsenic accumulation and metabolism in rice (*Oryza sativa* L.). *Environ. Sci. Technol.* 36(5), 962-968.
- (2) Abedin, M.J., Feldmann, J., Meharg, A.A., 2002b. Uptake kinetics of arsenic species in rice plants. *Plant Physiol.* 128, 1120-1128.
- (3) Ackerman, A.H., Creed, P.A., Parks, A.N., Schwegel, C.A., Creed, J.T., Heitkemper, D.T., Vela, N.P., 2005. Comparison of a chemical and enzymatic extraction of arsenic from rice and an assessment of the arsenic absorption from contaminated water by cooked rice. *Environ. Sci. Technol.* 39, 5241-5246.



- (4) Anastasia, F.B., Kender, W.J., 1973. The influence of soil arsenic on the growth of lowbush. *Environ. Qual.* 2, 335-337.
- (5) Bae, M., Watanabe, C., Inaoka, T., Sekiyama, M., Sudo, N., Bokul, M.H., Ohtsuka R., 2002. Arsenic in cooked rice in Bangladesh. *The Lancet.* 360, 1839-1840.
- (6) Bruce, S.L., Noller, B.N., Grigg, A.H., Mullen, B.F., Mulligan, D.R., Ritchie, P.J., Currey, N., Ng, J.C., 2003. A field study conducted at Kidston gold mine to evaluate the impact of arsenic and zinc from mine tailing to grazing cattle. *Toxicol. Letters.* 137, 23-34
- (7) Campbell, J.A., Stark, J.H., Carlton-Smith, C.H., 1985. International symposium on heavy metals in the environment, Athens, Vol. 1, p 478.
- (8) Carbonell, A.A., Aarabi, M.A., Delaune, R.D., Grabrell, R.P., Patrick, W.H.Jr., 1998. The influence of arsenic chemical form and concentration on *Spertina partens* and *Spertina alterniflora* growth and tissue arsenic concentration. *Plant and Soil.* 198, 33-43.
- (9) Chen, J.C., Lin, L.J., 1994. Human carcinogenicity and antherogenicity induced by chronic exposure to inorganic arsenic. In: Nriagu, J.O. (Ed.), *Arsenic in the environment: Part II; Human health and ecosystem effects.* Wiley & Sons, New York, pp. 09-157.
- (10) Das, H.K., Mitra, A.K., Sengupta, P.K., Hossain, A., Islam, F., Rabbani, G.H., 2004. Arsenic concentrations in rice, vegetables and fish in Bangladesh: a preliminary study. *Environ. Inter.* 30, 383-387.
- (11) D'Illo, S., Alessandrelli, M., Cresti, R., Forte, G., Caroli, S., 2002. Arsenic content of various types of rice as determined by plasma-based techniques. *Microchemical Journal.* 73, 195-201.

- (12) Duxbury, J.M., Mayer, A.B., Lauren, J.G., Hassan, N., 2003. Food chain aspects of arsenic contamination in Bangladesh: effects on quality and productivity of rice. *Environ. Sci. Health.* 38(1), 61-69.
- (13) Farmer, J.G., Johnson, L.R., 1990. Assessment of occupational exposure to inorganic arsenic based on urinary concentrations and speciation of arsenic. *Br. J. Ind. Med.* 47, 342-8.
- (14) Jiang, Q.Q., Singh, B.R., 1994. Effect of different form and sources of arsenite on crop yield and arsenic concentration. *Water, Air and Soil Pollut.* 74 (3-4), 321-343.
- (15) Johnson, L.R., Hiltbold, A.E., 1969. Arsenic content of soil and crops following use of methanearsonate herbicides. *Soil Sci. Soc. Am. Proc.* 33, 279-282.
- (16) Karim, M.M., 2000. Arsenic in groundwater and health problems in Bangladesh. *Water Res.* 34, 304-310.
- (17) Khattak, R.A., Page, A.L., Parker, D.R., Bakhtar, D., 1991. Accumulation and interactions of arsenic, selenium, molybdenum and phosphorus in alfalfa. *Environ. Qual.* 20, 165-168.
- (18) Klose, R., Braun, C., 1997. Studies on arsenic transfer and recommendations in agricultural use of an area in Saxony with high arsenic pollution. *Kongressband 1997 leipzig. Stoff und Energiebilanzen in der Landwirtschaft und weitere Beitrage aus den offentlichen Sitzungen*, Leipzig, Germany, 15-19 September, pp. 501-504.
- (19) Liao, C.M., Ling, M.P., 2003. Assessment of human health risks for arsenic bioaccumulation in tilapia (*Oreochromis mossambicus*) and large-scale mullet (*Liza macrolepis*) from Blackfoot disease area of Taiwan. *Arch. Environ. Contam. Toxicol.* 45, 264-272.
- (20) Liebig, G.F., 1966. Diagnostic criteria for plants and soils. In: Chapman, H.D. (Ed.) *Arsenic*. University of California press, Riverside C.A.

- (21) Marin, A.R., Masscheleyn, P.H., Patrick, W.H.Jr., 2003. Soil redox-pH stability of arsenic species and its influence on arsenic uptake by rice. *Plant and soil*. 152, 245-253.
- (22) Marin, A.R., Masscheleyn, P.H., Patrick, W.H.Jr., 1992. The influence of chemical form and concentration of arsenic on rice growth and tissue arsenic accumulation. *Plant and Soil*. 139, 175-183.
- (23) Mason, R.P., Laporte, J., Andres, S., 2000. Factors controlling bioaccumulation of mercury, methylmercury, arsenic, selenium and cadmium by freshwater invertebrates and fish. *Arch. Environ. Contam. Toxicol.* 38(3), 283-297.
- (24) Nicholson, F.A., Chambers, B.J., Williams, J.R., Unwin, R.J., 1999. Heavy metal contents of livestock feeds and animal manures in England and Wales. *Bioresour. Technol.* 70, 23-31.
- (25) NRC, 2001. *Arsenic in drinking water*. National Academy Press, Washington, DC.
- (26) Onken, B.M., Hossner, L.R., 1995. Plant uptake and determination of arsenic species in soil solution under flooded conditions. *Environ. Qual.* 24, 373-381.
- (27) O'Neil, P., 1995. Heavy metals in soils. In: Alloway B. J. (Ed.), *Arsenic*. Blackie Academic & Professional, London. Pp. 105-121.
- (28) Paul, P.C., Chattopadhyay, A., Dutta, S.K., Mazumder, D.N., Santra, A., 2000. Histopathology of skin lesions in chronic arsenic toxicity--grading of changes and study of proliferative markers. *Indian J. Pathol. Microbiol.* 43(3), 257-64.
- (29) Peryea, F.J., 1991. Phosphate induced release of arsenic from soils contaminated with lead arsenate. *Soil Sci. Soc. Am. J.* 55, 1301-1306.
- (30) Rahman, M.A., Hasegawa, H., Rahman, M.A., Rahman, M.M., Miah, M.A.M., 2006. Influence of cooking method on arsenic retention in cooked rice related to dietary exposure. *Sci. Total Environ.* 370, 51-60.

- (31) Rahman, M.A., Rahman, M.M., Miah, M.A.M, Khaled, H.M., 2004. Influence of soil arsenic concentrations on rice (*Oryza sativa* L.). Sub-trop. Agric. Res. & Dev. 2(3), 24-31.
- (32) Roychowdhury, T., Uchino, T., Tokunaga, H., Ando M., 2002. Survey of arsenic in food composites from an arsenic-affected area of West Bengal, India. Food and Chem. Toxicol. 40, 1611-1621.
- (33) Sadiq, M., 1986. Solubility relationships of arsenic in calcareous soils and its uptake by corns. Plant and Soil. 91, 241-248.
- (34) Schoof, R.A., Yost, L.J., Eickhoff, J., Crecelius, E.A., Cragin, D.W., Meacher, D.M., Menzel, D.M., 1999. A market basket survey of inorganic arsenic in food. Food and Chem. Toxicol. 37, 839-846.
- (35) Shariatpanahi, M., Anderson, A.C., 1984. Distribution and toxicity of monosodium methanearsonate following oral administration of the herbicide to dairy sheep and goats. Environ. Sci. Health. 19(4-5), 427-39.
- (36) Thornton, I., Webb, J.S., 1979. Aspects of geochemistry and health in the United Kingdom. Phy. and Chem. of the Earth. 11, 791-805.
- (37) Tsutsumi, S, Amagai, T, Kawaguchi, M, Ishizuka, S, Matsumoto, Y., 1980. Effects of arsenic trioxide ( $As_2O_3$ ) on the uptake of 3H-thymidine in lymphocytes of mice. Bull. Tokyo Dent. Coll. 21(2), 63-70.
- (38) Ullah, S.M., 1998. International conference on arsenic pollution of groundwater in Bangladesh: causes, effects and remedies, Dhaka community hospital, Dhaka, Bangladesh. p 133.
- (39) Vaughan, G.T., 1993. The environmental chemistry and fate of energy pesticides in cattle tick dip sites and banana plantations. Investigation report CET/LHIR 148, CSIRO, Division of Coal and Energy Technology, Centre for Advanced Analytical Chemistry, Sydney, Australia.

- (40) Welsch, E.P., Crock, J.G., Sanzolone, R., 1990. Trace-level determination of arsenic and selenium using continuous-flow hydride generation atomic absorption spectrophotometry (HG-AAS). In: Arbogast, B.F. (Ed.), Quality assurance manual for the Branch of Geochemistry. Open-File Rep. 90-0668. U.S. Geological Survey, Reston, VA. p. 38-45.
- (41) Walsh, L.M., Keeney, D.R., 1975. Behavior and phytotoxicity of inorganic arsenicals in soils. In: Woolson, E.A. (Ed.), Arsenical pesticides. Am. Chem. Soc. Symp. Washington D.C. pp. 35-52.
- (42) WHO, 2001. Arsenic and arsenic compounds. 2<sup>nd</sup> (Ed.), IPCS Environmental Health Criteria Document 244. International Program on Chemical Safety, WHO, Geneva.
- (43) Williams, P.N., Price, A.H., Raab, A., Hossain, A.A., Feldmann, J., Meharg, A.A., 2005. Variation in arsenic speciation and concentration in paddy rice related to dietary exposure. *Environ. Sci. Technol.* 39, 5531-5540.
- (44) Woolson, E.A., Axley, J.H., Kearney, P.C., 1973. The chemistry and phytotoxicity of arsenic in soils, effects of time and phosphorous. *Soil Sci. Soc. Am. Proc.* 37, 254-259.

Table 1: Physical properties of initial soil

Physical properties	Soil of glasshouse experiment	Field soil
% Sand (2 – 0.05 mm)	12.30	8.25
% Silt (0.05 – 0.002 mm)	53.00	27.5
% Clay (< 0.002 mm)	34.70	64.5
Textural Class	Silty-clay-loam	Clay-loam
Moisture (%)	16.04	-

Table 2: Chemical properties of initial soil

Chemical properties	Soil of glasshouse experiment	Field soil
pH (Soil : Water = 1 : 2.50)	5.27	7.07
Organic Carbon (%)	0.77	-
Organic Matter (%)	1.32	-
Total Nitrogen (%)	0.25	-
Total Phosphorus (%)	0.02	-
Total Potassium (%)	0.12	-
Total Iron (%)	2.01	0.21
Total Arsenic (mg kg <sup>-1</sup> )	3.25 (+10)*	14.51
Available Phosphorus (mg kg <sup>-1</sup> )	6.15 (+3.12)**	6.03
Total Manganese (mg kg <sup>-1</sup> )	-	247.14

\* 10 mg of As kg<sup>-1</sup> soil was spiked to the initial soil of glasshouse experiment.


\*\* After the rice harvest, available phosphate in the soil was 9.27 mg kg<sup>-1</sup>.

Table 3: Arsenic accumulation in rice plant tissues affected by soil arsenic concentrations <sup>a</sup>

Spiked arsenic to the initial soil (mg kg <sup>-1</sup> )	Arsenic content (mg kg <sup>-1</sup> dry weight)			
	straw (PI stage)	Straw (maturity stage)	Husk	Grain
Control	0.6±0.01f	0.9±0.01f	0.2±0.01e	0.2±0.01e
10	2.5±0.02e	2.1±0.01e	0.3±0.01e	0.3±0.01bce
20	4.8±0.10d	7.4±0.02d	0.6±0.03bc	0.4±0.04bc
30	6.2±0.04c	9.1±0.04d	0.4±0.02de	0.4±0.04bc
40	6.1±0.03c	12.3±0.03c	0.5±0.04cd	0.5±0.02a
50	7.6±0.22c	12.5±0.02c	0.8±0.02b	0.3±0.11bc
60	20.6±0.52a	23.7±0.44a	1.6±0.15a	0.4±0.01bc
70	12.0±0.03b	13.2±0.05c	0.6±0.01bc	0.3±0.03bce
80	10.7±0.01b	17.1±0.32b	0.2±0.01e	0.3±0.02bce
90	-	17.3±0.21b	-	-


<sup>a</sup> The values are mean ± S.D. of three replicates. The data were statistically analyzed by Duncan Multiple Range Test (DMRT) at 5% level. In a column, values having different letters (a - f) indicate significant differences ( $p < 0.05$ ) among them.

**Figure 1:**







**Figure 2:**



**Figure 3:**



**Figure 4:**



## **Figure Captions**

Figure 1: Populations of Bangladesh, one of the severely arsenic affected areas of the world, have been exposing arsenic poisoning from drinking water directly. There are some other possible food chain pathways of terrestrial ecosystem through which human may be exposed to arsenic poisoning from many sources as they are one of the topmost consumer of the ecosystem. In the above diagram, those possible food chain pathways have been shown and the “Plant-Man” food chain has been explained in this paper to understand the arsenic toxicity to human through this pathway. Some other pathways like “Plant-Animal-Man” could be potential for human exposure to arsenic and needs to be investigated.

Figure 2: Effect of soil arsenic concentrations on arsenic uptake in rice straw. Error bars represent mean  $\pm$  SEM of three replicates. Arsenic in straw was measured at two growth stages of rice plant. At the panicle initiation (PI) stage, about 30 days after transplantation and at maturity stage (after harvest).

Figure 3: Effect of soil arsenic concentrations on arsenic uptake in rice grain and husk. Error bars represent mean  $\pm$  SEM of three replicates. Raw rice was sun dried and the husk was removed from rice grain to determine arsenic.

Figure 4: Arsenic accumulation in rice tissues. Error bars represent mean  $\pm$  SEM of three replicates. Arsenic concentrations in field soil and irrigation water were  $14.51 \pm 0.21$  and  $0.07 \pm 0.02$  mg kg<sup>-1</sup>.