

A comparison of First Nations and Non-First Nations Australians' survey responses to COVID-19 risks and impacts: Implications for health communications

Julaine Allan (✉ julaine@uow.edu.au)

University of Wollongong

Jodie Kleinschafer

Charles Sturt University

Teesta Saksena

Western NSW Local Health District

Azizur Rahman

Charles Sturt University

Jayne Lawrence

Charles Sturt University

Mark Lock

Research Article

Keywords: First Nations Australians, rural health, COVID-19, Health risk communication

Posted Date: March 28th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1476169/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

This study investigated differences between rural Australian First Nations and non-First Nations survey respondents' perceptions of COVID-19 related risks; and analysed other variables that could predict an exacerbation of anxiety related to COVID-19 harms. Methods A cross-sectional online and paper survey of rural and regional residents from the western regions of NSW, Australia was conducted. Descriptive and multivariate statistical analyses were used to assess links between First Nations status and other variables. Results There were significant differences between First Nations (n=60) and non-First Nations (n= 639) respondents across all socio-demographic categories. The results reflect a significantly higher level of anxiety among the First Nations Australians in the sample: they felt afraid more often, felt it was highly likely they would catch the virus and if they did catch the virus perceived that it would be very harmful. Living with children under eighteen years of age and in small rural towns were key factors linked to feeling afraid of COVID-19 and First Nations status. Conclusion Health risk communication in pandemic response should include an equitable focus on rural areas, recognising that First Nations Australians are a significant proportion of the rural population with different risk factors and concerns than those of non-First Nations Australians. This principle of First Nations led design is critical to all health policy and planning. Governments globally should include rural areas in planning pandemic responses, recognising that First Nations populations are a significant proportion of the rural population creating syndemic conditions.

Introduction

Rural populations globally have experienced significant impacts from COVID-19. While infections spread more rapidly in highly populated areas; once the virus arrived in rural areas, mortality rates were higher and economic and social impacts more serious [1, 2]. Impacts on rural communities were attributed to generally poorer health including chronic conditions, an older population, lower education; and employment that had to be undertaken in person [3].

Australia has reported low numbers of COVID-19 infections compared to many places in the World. Geographic isolation and good infection control resulted in a small proportion of the population being affected by the disease [4]. At the beginning of the pandemic with limited experience of similar health crises the government scrambled to identify and protect the country's most vulnerable groups, including people with chronic illnesses, those in aged care and First Nations Australians. The phrase 'First Nations Peoples' refers to Indigenous peoples worldwide but especially in colonised nations of Australia, Aotearoa / New Zealand, Canada, and the United States. In this paper, the phrase 'First Nations Australians and First Nations' refers to Aboriginal and Torres Strait Islander peoples in Australia [5]. First Nations Australians were found to be at greater risk of morbidity and mortality during past influenza pandemics [6] and gaps had been identified in existing disaster response plans including a lack of information targeted to First Nations Australians [7].

People in rural and remote Australia were not specifically included as a vulnerable group in COVID response plans even though they were likely more at risk from COVID-19 for the same reasons as rural populations in other countries. First Nations Australians are a higher proportion of the population in rural areas compared to urban areas and due to widespread racism, structural disadvantage and dispossession of land are disproportionately affected by poor physical and mental health, lower incomes and crowded housing prior to COVID-19 compared to non-First Nations Australians [8, 9. NSW Ministry of Health 2020]. COVID-19 converging with existing social and health conditions including an epidemic of poor mental health has the potential to cause a syndemic for First Nations Australians [9, 10]. Already limited access to rural health care was reduced during the pandemic due to service closures and fear of disease contagion. A move to virtual healthcare delivery was constrained in many rural areas because of unreliable connections and poor coverage [11]. This has been an additional impact of COVID-19 on First Nations Australians who already had poorer access to digital devices and virtual healthcare [12].

While a NSW pandemic preparedness guideline released in July 2019 was based on extensive consultation with First Nations stakeholders, there was a notable absence of empirical research informing the strategy [13]. In addition, there was an absence of research examining rural First Nations Australian's perceptions of COVID-19 risks and information and communication needs to better inform culturally safe community management and COVID recovery plans [14]. The COVID-19 Delta variant's spread through Western NSW communities has highlighted limited access to vaccination and health workforce shortages adding to vulnerability of

First Nations communities [15]. Redistribution of vaccines from rural NSW to non-First Nations school children in Sydney reinforced long-held views of First Nations Australians being treated as second class citizens.

Health literacy is a key factor in accessing and interpreting pandemic information. Studies of the current COVID-19 and previous Swine flu pandemic have found that health messaging has been confusing and difficult to understand, with written materials requiring above average reading ability, and with limited attention paid to providing targeted messages to marginalised communities [16, 17]. Different consumer populations require nuanced communications that addresses their cultural milieu including for First Nations Australians, a distrust of government and poor health care experiences [18]. The Aboriginal Community Controlled health sector worked hard to inform communities around Australia about COVID-19 risks including closing remote communities and developing localised social media campaigns for these sites [19]. As the pandemic continues development of specific health communications for rural and remote people in NSW is required with tailored options for First Nations Australian communities in the region.

Risk perception and resultant behaviour is strongly influenced by personal, social and cultural contexts [20]. Two Australian studies have identified differences between First Nations and non-First Nations respondents in relation to COVID-19. One found that non-Caucasian people were more likely to engage in protective health behaviours and included First Nations Australians in that group [21]. And the other found the First Nations Australians perceived a greater risk from people who were not vaccinated [22]. However, there has been no investigation of the COVID-19 risk perceptions of First Nations Australians living in rural areas compared to those of non-First Nations rural and remote populations. In particular, there is an absence of research examining rural First Nations people's concerns about COVID-19 and its likely impacts; to describe and compare factors that could better inform culturally safe community management and COVID recovery plans [14, 6]. To address this gap this study investigated differences between rural dwelling First Nations and non-First Nations survey respondents' perceptions of COVID-19 related risks during the first COVID lock down in Australia; and analysed other variables that could predict an exacerbation of anxiety related to COVID-19 harms.

Methods

A cross-sectional online and paper survey of rural and regional residents from the western regions of NSW was conducted. The research team included two First Nations researchers (a Wiradjuri Woman and Ngiyampaa Man). A First Nations Reference Group was convened with members of the western NSW community and met monthly for the life of the project. This group oversaw the cultural safety and sensitivity of the project [23]. For example, they reviewed and amended the survey (e.g. changing problematic language and including additional socio-demographic questions about household types, living arrangements, the role of community leaders and the types of health services available. They also recommended data collection methods and advised on the implications of the results.

Setting

Western NSW accounts for 29% of the NSW population and has the largest population of First Nations Australians in the country [24]. It is also home to the largest language group in Australia – Wiradjuri. The First Nations population in the study region ranges from 19% in Dubbo to 79% in Brewarrina [25]. The study began three months after COVID-19 was declared a global pandemic and the Australian government began emphasising the need for personal protective measures including social distancing, hand sanitising and staying home as much as possible. At the time mask wearing had not been mandated and unlike other measures, the effectiveness of masks was being debated. No vaccine was available at this time.

Participants

Quota sampling was used to approximate a representative sample of participants [26]. Participants were aged 18 years and over, able to read and understand English and resided in Western NSW. Participants were recruited from two market research companies, Dynata and Qualtrics. Dynata and Qualtrics are large market research companies with over 600,000 panel members each in Australia. Panel members recruited via Dynata and Qualtrics were sent an email invitation to participate in the study and received points for completing the online survey, which they could redeem for gift vouchers, donations to charities or money. There was additional recruitment through personal and community networks to improve the representation of First Nations participants. To facilitate access for First Nations Australians, information about the survey was distributed through Elders groups, the project's

First Nations Reference Group and people were given the opportunity to complete a paper based version of the survey with support of First Nations members of the research team. Recruitment of all participants occurred between July and August 2020.

Survey

The online survey used in this study was an adaptation of a questionnaire developed for assessment of risk perceptions, anxiety, and behavioural responses of the general public during the early phase of the Influenza A (H1N1) pandemic [24]. The questionnaire fitted the objectives of this study given the contextual similarities, theoretical models explaining health behaviour, design and reliability (Cronbach's alpha) of its constructs ranging from 0.6 to 0.9, with trends analysed over time [27].

A number of variables were chosen because a holistic approach that includes individual demographics, cognitive, affective, social and cultural factors is recommended to assess risk perceptions [28]. Demographic measures included: First Nations status, postcode, age, gender, education, rural or town/village location, proximity to medical services and living situation. Demographic measures were selected based on advice from the Project Advisory Group. This study analysed five items related to perceptions about COVID-19: perceived likelihood of getting COVID - 19 in the next 12 months, perceived harmfulness of the virus and how often people felt afraid. Two further items were drawn from the survey's maladaptive response scale: perception about respondents' ability to do something about the virus and perceived economic impacts of the pandemic.

Data analysis

The data reported here comes from a larger study. This analysis is focused on those perceptions where there were significant differences between First Nations and non-First Nations respondents. Respondents from Murrumbidgee, Western and Far Western NSW Local Health Districts were identified by postcode. Respondents from other areas were excluded from the analysis.

Both descriptive and multivariate statistical analyses were used to assess links between First Nations status and other variables. Bivariate analyses using cross-tabulations were performed for the respondent's First Nations status variable by anxiety about COVID-19 and socio-demographic characteristics which are considered for multivariate analysis. Significant determinants were explored by Pearson's Chi-squared test and Fisher's exact test. The multinomial logistic regression modelling for multivariable analyses was carried out to determine the influence of some covariates on the likelihood of experiencing anxiety for COVID-19 such as 'feeling afraid' and 'perceiving harmful' variables included in the survey.

The significant relationships between a variable and its effects were quantified by calculating the odds ratios with 95% confidence interval measures. The odds ratio (OR) in favour of 'moderate' and 'severe' form of anxieties were computed for the selected group of covariates to suggest how many times the group of interest is more likely to belong to the target group compared to the reference group, i.e. 'no' anxiety. Two regression models were used separately for the two different anxiety related response variables related to COVID-19 perceptions. Moreover, the -2 Log Likelihood-based Chi-squared test was employed to check the statistical significance of the fitted model. Further details about these methods and analysis techniques are available in the existing literature [29]. IBM SPSS version 26 was utilised in all statistical analyses.

Ethics Approval

Ethics approval was provided by the Aboriginal Health and Medical Research Council of NSW (AH&MRC) [1668/20] to enable the analysis of First Nations responses and Charles Sturt University Human Research Ethics Committee (H20254).

Results

There were 701 completed surveys collected. Paper based surveys were completed by seven First Nations Australian respondents. The remaining 694 surveys were completed online. There were more female (n = 447) than male respondents (n = 254). Table 1 reports the survey respondent's demographic characteristics. There were significant differences between First Nations (n = 60) and non-First Nations (n = 639) respondents across all categories. First Nations respondents were significantly more likely to be female, to have a post graduate degree, to live in a regional town more than 20 kilometres away from a health service and be looking after children in their home compared to non-First Nations respondents.

Table 1
Background characteristic of the respondents and their percentage distribution by the First Nations and non-First Nations status from selected study areas in Australia, 2021 (n = 701)

Characteristic and categories	No. of participants ^a	%	First nations %	Non-First Nations or does not identify %	p-value
Sex					p = 0.004
Male	254	36.2	34.6	53.3	
Female	447	63.8	55.4	46.7	
Age (in years)					p = 0.000
18–29	151	21.6	25.0	136	
30–49	235	33.6	56.7	201	
50–69	218	31.2	13.3	210	
70 plus	95	13.6	5.0	92	
Education					p = 0.000
School up to Year 12	228	32.5	21.7	33.5	
Trade/Diploma	257	36.7	28.3	37.4	
Bachelor	158	22.5	16.7	23.1	
Postgraduate	58	8.3	33.3	5.9	
Children under 18 at home					p = 0.003
Yes	224	75.2	92.9	72.3	
No	74	24.8	7.1	27.7	
Residence					p = 0.067
Regional Town over 500 population	603	86.4	93.3	85.7	
Rural Area	95	13.6	6.7	14.4	
Nearest health service					p = 0.01
Up to 20kms	632	90.2	78.3	91.3	
More than 20kms	69	9.8	21.7	8.7	
Living situation					p = 0.000
Single person /couple without children	137	19.5	13.3	20.1	
Couples/single parent with children	265	37.8	15	39.9	
Other household types	299	42.7	71.7	39.9	
^a Total number of participants may differ due to missing data.					

Having compared the demographic characteristics of the First Nations respondents with non-First Nations members of the sample, in the next section the groups are compared in terms of their perceptions about COVID-19. Table 2 includes five measures related

to COVID-19 perceptions where there was a significant difference between the two groups: perceived likelihood of getting COVID -19 in the next 12 months, perceived harmfulness of the virus, how often people felt afraid because of COVID-19, perceptions about respondents ability to do something about the virus and perceived economic impacts of the pandemic.

Table 2
COVID-19 related perceptions and anxiety characteristic of the participants and their percentage distribution by the First Nations and non-First Nations status, 2021.

Factors	Number of observations (%)	First Nations	Non-First Nations	p-value
Likelihood of getting in next 12 months				
Not very likely	358 (52.7%)	53.90%	40.40%	p = 0.000
Moderately likely	259 (38.1%)	40.50%	12.30%	
Very likely	62 (9.1%)	5.60%	47.40%	
Perceived Harmfulness				
Not very harmful	59 (8.4%)	8.80%	3.30%	p = 0.028*
Moderately harmful	204 (29.1%)	30.20%	18.30%	
Very harmful	438 (62.5%)	61.10%	78.30%	
Feel Afraid				
Not often/Rarely	318 (45.4%)	47.00%	28.80%	p = 0.022
Sometimes	209 (29.9%)	29.30%	35.60%	
Often/All of the time	173 (24.7%)	23.70%	35.60%	
There is nothing we can do about COVID-19				
Disagree	425 (60.7%)	63.4%	18.3%	p = 0.000
Neither agree nor disagree	169 (24.1%)	25.0%	21.7%	
Agree	106 (15.1%)	11.6%	60.0%	
Economic impacts of COVID-19 not as bad as predicted				
Disagree	543 (77.7%)	80.8%	45.0%	p = 0.000
Neither agree nor disagree	89 (12.7%)	12.4%	16.7%	
Agree	67 (9.6%)	6.9%	38.3%	
*p-value estimated using the Fishers Exact Test.				

The results reflect a significantly higher level of anxiety among the First Nations group in the sample: they felt afraid more often, felt it was highly likely they would catch the virus and if they did catch the virus perceived that it would be very harmful.

- First Nations respondents (47.4%) were eight times more likely than non-First Nations respondents (5.6%) to indicate they were “very likely to get COVID-19 in the next 12 months.
- Though the majority of the sample considered the disease harmful, more First Nations respondents perceived COVID-19 was very harmful (78.3% versus 61.1%), whereas non-First Nations respondents were more likely to consider the disease moderately harmful (30.2% versus 18.3%).

- Nearly half of non-First Nations respondents (47%) reported rarely feeling afraid about COVID-19 compared to only 28.8% of those who identified as First Nations.
- Two thirds (63.6%) of First Nations respondents agreed that there was nothing they could do about COVID-19 whereas only 11.6% of the rest of the sample agreed with this statement.
- In relation to the economic impacts of COVID-19 Non-First Nations members of the sample (80.8%) were much more likely to expect a negative impact compared to First Nations respondents (45%).

Overall these findings reflect a much higher level of anxiety and fatalism amongst the First Nations respondents.

The significant differences in other demographic characteristics included in Table 1 suggested that there may be more than First Nations status influencing the perceptions of each group, for example, education, age, gender, and population where people live and whether children were present in the household. The following section reports the results of two multinomial regression analyses examining the predictive relationship between demographic characteristics of respondents and 1) how often they felt afraid of COVID-19; and 2) how harmful they perceived the virus was.

In the first regression (Table 3) how often people reported feeling afraid about the virus was considered. Non-First Nations respondents were 60% less likely to report feeling afraid sometimes than First Nations respondents, illustrating a very different level of affect between these groups. Compared to those living in rural and remote areas, those living in regional towns were less likely to feel afraid sometimes (51.1%) and all of the time (79.6%). Males were six times more likely than females to report feeling afraid all of the time, and those with children under 18 in their household were 1.8 times more likely to report feeling afraid sometimes (when compared to those who did not feel afraid). Finally, those with a trade certificate or a diploma were 71.6% less likely to say they feel afraid sometimes than those with a postgraduate degree.

Table 3

Regression coefficients and odds ratios for the likelihood of moderate and severe form of anxiety (i.e. feeling afraid) due to COVID-19 by some selected significant characteristics, including the First Nations status of the respondents^a

Feel Afraid (Reference category is not afraid)						
	Afraid sometimes			Afraid often /all of the time		
Characteristic ^b	B	Odds Ratio	95% CI	B	Odds Ratio	95% CI
Non-First Nations	-0.92	0.40***	(0.159, 1.006)	0.15	0.87	(0.124, 6.019)
Male	-0.05	0.95	(0.493, 1.825)	1.79	6.01**	(1.743, 20.721)
Children under 18	0.57	1.77*	(0.963, 3.238)	0.90	2.45	(0.605, 9.957)
Regional Town over 500 ppl	-0.73	0.49*	(0.215, 1.070)	-1.60	0.20**	(0.050, 0.832)
Education						
Year 12	-0.86	0.42	(0.141, 1.274)	0.44	1.55	(0.117, 20.461)
Trade/Diploma	-1.26	0.28**	(0.096, 0.840)	0.17	1.18	(0.091, 15.345)
Bachelor	-0.59	0.55	(0.174, 1.752)	1.69	5.39	(0.456, 63.762)
Model fitting information:						
	-2 Log Likelihood			135.208		
	Chi-squared (df)			41.994 (14)****		
^a Reference category of dependent variable is not afraid.						
^b Omitted categories (i.e. reference class for each independent variable) not shown.						
*p < 0.10; **p < 0.05; ***p < 0.01; ****p < 0.001						

In the second regression the perceived harmfulness of the virus was examined (Table 4). First Nations status was the only variable that was significantly related to perceptions of harmfulness. When compared to those who thought COVID was not harmful, Non-First Nations respondents were 66% less likely than First Nations respondents to perceive COVID-19 was moderately harmful and 93% less likely than First Nations respondents to report that the virus was very harmful. Education level does not make a difference in this model.

Table 4

Regression coefficients and odds ratios for the likelihood of moderate and severe form of anxiety (i.e. perceiving harmful) due to COVID-19 by some selected significant characteristics, including the First Nations status of the respondents^a

Harmful (reference category is not harmful)						
	Moderately harmful			Very Harmful		
Characteristic ^b	B	Odds Ratio	95% CI	B	Odds Ratio	95% CI
Non-First Nations	-1.16	0.31	(0.034, 2.899)	-2.73	0.07***	(0.008, 0.533)
Male	-0.56	0.57	(0.237, 1.365)	-0.61	0.54	(0.237, 1.240)
Children under 18	0.38	1.47	(0.619, 3.483)	0.02	1.02	(0.454, 2.279)
Regional Town over 500ppl	0.42	1.52	(0.458, 5.016)	-0.42	0.66	(0.230, 1.881)
Education Trade/Diploma	0.23	1.26	(0.546, 2.903)	-0.032	0.97	(0.439, 2.130)
Model Fitting information						
	-2 Log Likelihood		113.622			
	Chi-squared (df)		28.588 (10)***			
^a Reference category of dependent variable is not harmful.						
^b Omitted categories (i.e. reference class for each independent variable) not shown.						
*p < 0.10; **p < 0.05; ***p < 0.01; ****p < 0.001						

Discussion

There were some significant differences in risk perception and impacts of COVID-19 between First Nations and non-First Nations survey respondents; and several demographic variables that predicted responses to COVID-19. First Nations respondents perceived COVID-19 to be more harmful than non-First Nations respondents, to perceive a higher danger and vulnerability from the virus, with just under half (47.4%) predicting they would contract the virus in the next 12 months compared to 5.6% of non-First Nations respondents. At the time the survey was conducted there was no COVID-19 in Western NSW. Therefore the finding of high perceptions of vulnerability from COVID-19 may reflect the media discourse about the high level of risk that First Nations people face in relation to COVID-19 and the potential impact on First Nations communities in other parts of Australia as well as confused messaging about risks to different places and sectors of the population [16, 18, 22].

There were significantly higher levels of anxiety among First Nations respondents (based on perceptions of fear and the harmfulness of the virus). Significantly higher levels of fatalism and a huge disparity in the perceived likelihood of catching the virus in the next 12 months point to the impact of COVID-19 anxiety on top of existing high levels of mental distress and social disadvantage [9]. All factors consistent with a syndemic [10] indicating a much more serious impact of COVID-19 on First Nations Australians compared to non-First Nations Australians. First Nation's survey respondent's fears were justified because the Delta variant of COVID-19 quickly took hold in small communities with limited healthcare services reflecting the pattern seen in other countries [15, 2]. Limitations in accessing healthcare services is also reflected in the results with 21.7% of First Nations respondents reporting the nearest health service to be more than 20 km as compared to the non-First Nations (8.7%) respondents, potentially leading to heightened concern. Excluding rural and remote populations from risk groups at the start of the pandemic and limiting the supply of vaccines to rural areas because of increasing concern about urban areas was a dangerous oversight by the Australian government. The high proportion of First Nations peoples in Western NSW was not taken into account in vaccine rollout plans nor the limited availability of healthcare services.

Other factors influencing perceptions of COVID-19's harmfulness were also characteristic of many First Nations communities. Residing in a small rural town and living with children under the age of eighteen years were significantly predictive of concerns about COVID 19 harms. First Nations peoples often live in larger extended family groups including children; and in Western NSW, are more likely to live in small rural communities than on farming properties [30]. Frequently changing advice about the risks both to and from children directly impacts First Nations communities and is likely to heighten fear and concern about COVID-19 risks.

A high proportion of First Nations survey respondents had a post- school qualification (78% compared to 42% nationally) [31]. Whereas 71% of the non-first Nations respondents only had school or vocational qualifications. High levels of general literacy are likely to be consistent with good health literacy [16]. Lower levels of fear and perceptions of COVID-19 harmfulness among non-First-Nations respondents could indicate poorer health literacy within this group and potentially a blasé response to COVID-19 risks. Lower levels of concern about pandemic risks typically results in less adherence to infection control measures putting vulnerable groups at risk [32, 33].

Health literacy is not the only issue at play in effectiveness of health communications [28]. The levels of fear and perceptions of harmfulness found in the survey results should be expected when distrust of government and previous poor health care experiences are widespread for First Nations people and reinforced by vaccine unavailability [18]. There were no First Nations representatives in the daily government press conferences that delivered health advice even though there were frequent mention of risks to 'the regions' [15, 18]. Although Australian health authorities might argue that their messaging was equitable, in that it was the same message to all Australians, the contribution of the research presented here is how those messages can better target their intended audience, by directly addressing the specific concerns of First Nations communities in rural areas via co-design of communications and health care strategies with First Nations Australians [9].

Strengths And Limitations

There has not been a First Nation's informed investigation of First Nations Australian's perceptions of COVID-19 in Western NSW. The results suggest key areas for more nuanced health communications to address. However, the small sample size (n = 60) was not representative of the First Nations population in Western NSW (8.4% in the sample vs 13% in the region) and the results do not represent the experiences or perceptions of First Nations Australians in other areas. However, the proportion of First Nations respondents in this study is greater than the population proportion nationally and more than reported in other Australian studies [21, 22]. Further the study included 78% of First Nations respondents with a post- school qualification compared to 42% nationally [31]. This result suggests however that health literacy is less like to be a factor in COVID-19 related anxiety than other factors such as mental distress and poor experiences of healthcare and government interventions [18, 10].

Conclusion

Health communications for First Nations Australians in rural areas should be designed and delivered by First Nation's Australians from those areas because they understand the rural context people live in. This principle of First Nations led design is critical to all health policy and planning. Governments globally should include rural areas in planning pandemic responses, recognising that First Nations populations are a significant proportion of the rural population with different risk factors and concerns than those of non-First Nations peoples, creating syndemic conditions.

Declarations

Ethics approval

This study was performed in line with the principles of the Declaration of Helsinki. Approval was granted by the Ethics Committee of The Aboriginal Health and Medical Research Council of NSW (1668/2020) and Charles Sturt University (H20254).

Consent to participate

Informed consent to participate was obtained from all survey respondents.

Consent to publish

Prior to seeking consent to participate all survey respondents were informed results would be published and that no identifying information would be collected.

Availability of data and material

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Funding

This work was supported by a Charles Sturt University COVID-19 Research Development grant.

Author Contributions

Authors JA, JK, TS, ML and JL contributed to the study conception and design equally. AR designed and led the statistical analysis. Material preparation, data collection and analysis were performed by JA, JK, TS and AR. The first draft of the manuscript was written by JA and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors have no relevant financial or non-financial interests to disclose.

Acknowledgements

Jess Kingsford, Gail Fuller, Oliver Burmeister and the Members of the Aboriginal Reference Group, Western NSW LHD.

References

1. OECD (2020), "Policy implications of coronavirus crisis for rural development", Tackling Coronavirus (COVID-19): Contributing to a Global Effort, OECD, Paris, https://read.oecd-ilibrary.org/view/?ref=134_134479-8kq0i6epcq&title=Policy-Implications-of-Coronavirus-Crisis-for-Rural-Development.
2. Leatherby, L. (2020), "The worst virus outbreaks in the U.S. are now in rural areas", <https://www.nytimes.com/interactive/2020/10/22/us/covid-rural-us.html>.
3. Peters, D. (2020), "Community susceptibility and resiliency to COVID-19 across the rural-urban continuum in the United States", *Journal of Rural Health*, Vol. 36/3, pp. 446–456, <https://doi.org/10.1111/jrh.12477>.
4. Australian Government, Department of Health and Ageing. Corona Virus (COVID-19) At a Glance 30 June 2020. Australian Government: Canberra. Retrieved 13 December 2021 <https://www.health.gov.au/sites/default/files/documents/2020/06/coronavirus-covid-19-at-a-glance-30-june-2020.pdf>
5. AIATSIS ?
6. Rudge S, Massey P. Responding to pandemic (H1N1) 2009 influenza in Aboriginal communities in NSW through collaboration between NSW Health and the Aboriginal community-controlled health sector. *N S W Public Health Bull.* 2010; 21(2): 26–9.
7. Australian Government, Department of Health and Ageing. National pandemic influenza exercise. Exercise Cumpston 06 report. 2007. Available at: <http://www.health.gov.au/internet/wcms/publishing.nsf/Content/ohp-cumpston-report.htm>. Accessed August 16, 2007
8. Australian Institute of Health and Welfare 2020. Indigenous health and wellbeing. Canberra: AIHW. Viewed 22 March 2021, <https://www.aihw.gov.au/reports/australias-health/indigenous-healthand-wellbeing>
9. NSW Ministry of Health. (2020). NSW Aboriginal Mental Health and Wellbeing Strategy 2020–2025. Sydney: NSW Ministry of Health. Accessed 20/12/21. Available from <https://www.health.nsw.gov.au/mentalhealth/resources/Publications/aborig-mh-wellbeing-2020-2025.pdf>
10. Tsai, Alexander C et al (2017). Co-occurring epidemics, syndemics, and population health. *The Lancet*, Volume 389, Issue 10072, 978–982

11. Better Internet for Rural, Regional & Remote Australia (2018) submission to the Regional Telecommunications Independent Review Committee (RTIRC).
https://www.infrastructure.gov.au/sites/default/files/submissions/better_internet_for_rural_regional_remote_australia_birrr.pdf
12. Follent, D., Paulson, C., Orcher, P., O'Neill, B., Lee, D., Briscoe, K. and Dimopoulos-Bick, T.L. (2021), The indirect impacts of COVID-19 on Aboriginal communities across New South Wales. *Med. J. Aust.*, 214: 199–200.e1.
<https://doi.org/10.5694/mja2.50948>
13. NSW Ministry of Health, 2019. Pandemic Preparedness and Response – Aboriginal Communities. Sydney. Available:
https://www1.health.nsw.gov.au/pds/ActivePDSDocuments/GL2019_009.pdf
14. Moodie, N, Ward, J, Dudgeon, P, et al. Roadmap to recovery: Reporting on a research taskforce supporting Indigenous responses to COVID-19 in Australia. *Aust J Soc Issues.* 2021; 56: 4–16. doi: 10.1002/ajs4.133.
15. Thorpe, N. (2021). Aboriginal elders fear COVID deaths as NSW's outbreak reaches Walgett Posted Sat 14 Aug 2021 at 6:08am Saturday 14 Aug 2021 at 6:08am, updated Sat 14 Aug 2021 at 10:20am <https://www.abc.net.au/news/2021-08-14/aboriginal-elders-fear-covid-deaths-in-walgett-outbreak/100376186>
16. McCaffery KJ, Dodd RH, Cvejic E, Ayre J, Batcup C, Isautier JMJ, Copp T, Bonner C, Pickles K, Nickel B, Dakin T, Cornell S, Wolf MS. Health literacy and disparities in COVID-19–related knowledge, attitudes, beliefs and behaviours in Australia. *Public Health Res Pract.* 2020;30(4):e30342012. First published: 5 November 2020. <https://doi.org/10.17061/phrp30342012>
17. Lin L, Savioa E, Agboola F, Viswanath K. What have we learned about communication inequalities during the H1N1 pandemic: a systematic review of the literature. *BMC Public Health.* 2014;14:484.
18. Cox, L. (2007). Fear, Trust and Aborigines: The Historical Experience of State Institutions and Current Encounters in the Health System. *Health and History*, 9(2), 70–92. <https://doi.org/10.2307/40111576>
19. Eades, S., Eades, F., McCaullay, D., Nelson, L., Phelan, P., & Stanley, F. (2020). Australia's First Nations' response to the COVID-19 pandemic. *Lancet (London, England)*, 396(10246), 237–238. [https://doi.org/10.1016/S0140-6736\(20\)31545-2](https://doi.org/10.1016/S0140-6736(20)31545-2)
20. Schneider, C., Dryhurst, S., Kerr, J., Freeman, A., Recchia, G., Spiegelhalter, D. & van der Linden, S. (2021) COVID-19 risk perception: a longitudinal analysis of its predictors and associations with health protective behaviours in the United Kingdom, *Journal of Risk Research*, 24:3–4, 294–313, DOI: 10.1080/13669877.2021.1890637
21. Faasse K, Newby J. Public Perceptions of COVID-19 in Australia: Perceived Risk, Knowledge, Health-Protective Behaviors, and Vaccine Intentions. *Front Psychol.* 2020;11:551004. Published 2020 Sep 30. doi:10.3389/fpsyg.2020.551004
22. Seale, H., Heywood, A.E., Leask, J. *et al.* Examining Australian public perceptions and behaviors towards a future COVID-19 vaccine. *BMC Infect Dis* 21, 120 (2021). <https://doi.org/10.1186/s12879-021-05833-1>
23. Gwynn, J., Lock, M., Turner, N., Dennison, R., Coleman, C., Kelly, B., & Wiggers, J. (2015). Aboriginal and Torres Strait Islander community governance of health research: Turning principles into practice. *Aust J Rural Health*, 23(4), 235–242.
<https://doi.org/10.1111/ajr.12182>
24. Australian Bureau of Statistics [ABS] 2019. Census of Population and Housing: characteristics of Aboriginal and Torres Strait Islander Australians, 2016. ABS cat. no. 2076.0. Canberra: ABS.
25. Australian Bureau of Statistics [ABS] 2016. Estimates of Aboriginal and Torres Strait Islander Australians. ABS Cat no. 3238.0.55.001. Canberra: ABS
26. Hair, J. F., Black, W. C., Babin, B. J., Anderson, R. E., & Tatham, R. L. (2006). *Multivariate Data Analysis*. New Jersey, USA: Pearson Prentice Hall.
27. Bults M, Beaujean DJ, de Zwart O, Kok G, van Empelen P, van Steenberghe JE, Richardus JH, Voeten HA. Perceived risk, anxiety, and behavioural responses of the general public during the early phase of the Influenza A (H1N1) pandemic in the Netherlands: results of three consecutive online surveys. *BMC Public Health.* 2011 Jan 3;11:2. doi: 10.1186/1471-2458-11-2. PMID: 21199571; PMCID: PMC3091536.
28. Dryhurst, S., C. R. Schneider, J. Kerr, A. L. J. Freeman, G. Recchia, A. M. van der Bles, D. Spiegelhalter, and S. van der Linden. 2020. "Risk Perceptions of COVID-19 Around the World." *Journal of Risk Research* 23 (7/8): 994–4461.
doi:<https://doi.org/10.1080/13669877.2020.1758193>.
29. Rahman, A., & Harding, A. (2016). *Small area estimation and microsimulation modeling*. CRC Press.
<https://doi.org/10.1201/9781315372143>.

30. Australian Institute of Health and Welfare 2019. Rural & remote health. Cat. no. PHE 255. Canberra: AIHW. Viewed 08 January 2021, <https://www.aihw.gov.au/reports/rural-remote-australians/rural-remote-health>
31. Australian Institute of Health and Welfare <https://www.aihw.gov.au/reports/australias-welfare/indigenous-education-and-skills>
32. Barr, M., Raphael, B., Taylor, M. et al. Pandemic influenza in Australia: Using telephone surveys to measure perceptions of threat and willingness to comply. *BMC Infect Dis* 8, 117 (2008). <https://doi.org/10.1186/1471-2334-8-117>
33. Eastwood, K., Durrheim, D., Francis, J., Espaignet, E., Duncan, S., Islam, F., & Speare, R. (2009). Knowledge about pandemic influenza and compliance with containment measures among Australians. *Bulletin of World Health Organisation*, 87, 588–594 | [doi:10.2471/BLT.08.060772](https://doi.org/10.2471/BLT.08.060772)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Appendix.docx](#)