




RESEARCH ARTICLE

Association between female genital mutilation and girl-child marriage in sub-Saharan Africa

Bright Opoku Ahinkorah¹, John Elvis Hagan Jr^{2,3*} , Abdul-Aziz Seidu^{4,5,6} ,
Obasanjo Afolabi Bolarinwa⁷, Eugene Budu⁴, Collins Adu⁸ , Joshua Okyere⁴,
Anita Gracious Archer⁹ and Thomas Schack³

¹School of Public Health, Faculty of Health, University of Technology Sydney, Australia, ²Department of Health, Physical Education, and Recreation, University of Cape Coast, Cape Coast, Ghana, ³Neurocognition and Action-Biomechanics-Research Group, Faculty of Psychology and Sport Sciences, Bielefeld University, Germany, ⁴Department of Estate Management, Takoradi Technical University, Takoradi Box 256, Ghana, ⁵College of Public Health, Medical and Veterinary Sciences, James Cook University, Townsville, Queensland QLD4811, Australia, ⁶Centre for Gender and Advocacy, Takoradi Technical University, Takoradi Box 256, Ghana, ⁷Department of Public Health Medicine, School of Nursing and Public Health, University of KwaZulu-Natal, Durban, South Africa, ⁸Department of Health Promotion and Disability Studies, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana and ⁹School of Nursing and Midwifery, University of Health and Allied Sciences, Ho, Ghana

*Corresponding author. Email: elvis.hagan@ucc.edu.gh

(Received 23 May 2021; revised 16 November 2021; accepted 16 November 2021)

Abstract

Two commonly linked harmful practices that negatively impact the health of girls and women in sub-Saharan Africa, and threaten their development and quality of life, are female genital mutilation and girl-child marriage. The central focus of the study was to investigate the association between female genital mutilation and girl-child marriage in sub-Saharan Africa. Data from the most recent Demographic and Health Surveys of twelve sub-Saharan African countries were pooled. A total of 14,748 women aged 20–24 were included in the study. A multilevel logistic regression analysis was employed, with reported adjusted odds ratios (aORs) and associated 95% confidence intervals (CIs). The overall prevalence of FGM in the twelve countries was 52.19%, with the highest prevalence in Guinea (97.17%). The overall prevalence of girl-child marriage in the twelve countries was 57.96%, with the highest prevalence in Chad (78.06%). Women who had never experienced female genital mutilation were less likely to experience girl-child marriage (aOR=0.76, CI=0.71–0.82) compared with those who had ever experienced female genital mutilation. Age 24 (aOR=0.47, CI=0.43–0.52), secondary/higher level of education (aOR=0.31, CI=0.28–0.35), richest wealth quintile (aOR=0.56, CI=0.47–0.66), exposure to mass media (aOR=0.81, CI=0.74–0.88) medium community literacy level (aOR=0.63, CI=0.57–0.69) and low community socioeconomic status (aOR=0.67, CI=0.49–0.92) were found to be protective against girl-child marriage. The findings reveal that female genital mutilation is associated with girl-child marriage in sub-Saharan Africa. The continued practice will adversely affect the reproductive health outcomes of girls in the sub-region. Policies aimed at eliminating female genital mutilation and girl-child marriage should focus on compulsory basic education, poverty alleviation and increasing access to mass media. Further, campaigns should cover more communities with lower literacy levels and medium socioeconomic status.

Keywords: Child marriage; Female genital mutilation; Sub-Saharan Africa

Introduction

Traditional cultural practices are deeply entrenched in many cultures and societies across the world and mirror perceived values and beliefs held by its members. While many indigenous

practices are useful to its populace, others can be detrimental, with women and children being the most vulnerable (United Nations Office of the High Commissioner on Human Rights, 2015). Two commonly linked harmful practices that have received considerable attention and which are pervasive in sub-Saharan Africa (Ahinkorah *et al.*, 2020) are female genital mutilation (FGM) and girl-child marriage (Karumbi & Gathara, 2020). FGM has often been related to girl's/women's eligibility for marriage and has been shown to be associated with the formation of formal partnerships for girls under the age of 18, defined as 'child marriage' (Varol *et al.*, 2014; Population Reference Bureau, 2018). These traditional practices not only negatively impact on the well-being of girls and women in sub-Saharan Africa, but also threaten their progress and quality of life (Karumbi & Gathara, 2020). FGM and girl-child marriage are social practices that function alone, but can be related to each other at different levels (Mackie & LeJeune, 2009).

According to UNFPA (2012), 30 countries in sub-Saharan Africa have a girl-child marriage incidence rate of 30% or higher. It is projected that 15 million girls are likely to enter into marriages before the age of 18 in sub-Saharan Africa, and without practical measures to halt girl-child marriage, a large number will be engaged before the age of 18 – an increase from the present estimated 720 million to 1.2 billion by 2050 (UNFPA, 2014). A recent study has shown that the overall prevalence of FGM among women in sub-Saharan Africa is 51.7%, with the largest proportion in Guinea (96.1%) and the lowest in Niger (4.2%) (Ahinkorah *et al.*, 2020).

FGM is regarded as a rite of passage into womanhood in many high-prevalence countries, and has strong ancestral and socio-cultural roots (Williams-Breault, 2018). It has been linked to girl-child marriage due to it being a rite of passage to maturity (Karumbi & Jacinta, 2017). FGM and girl-child marriage, as traditional practices, are often connected by a wide range of socio-cultural drivers such as the maintenance of family values (i.e. maintaining dignity and honour), preservation of girls' virginity before marriage, promoting fidelity within marriage, societal integration of the girl and family and the offer of monetary assurance to those within the poorest wealth quintile of society (Boyden *et al.*, 2012). Furthermore, both FGM and girl-child marriage have similar undesirable outcomes, such as increased risk of maternal and newborn death, delivery difficulties, social isolation and intimate partner violence (Karumbi *et al.*, 2017).

Research using recent data from Demographic and Health Surveys (DHSs) and Multiple Indicator Cluster Surveys providing nationwide representative samples of women aged 15–49 in ten countries in sub-Saharan Africa has shown that FGM and girl-child marriage vary markedly by country, region and/or ethnic affiliation (Population Reference Bureau, 2018). Among sub-Saharan African women who report girl-child marriage, the proportion who have undergone FGM is higher than among those who do not report girl-child marriage (>50%), with significant variations in Burkina Faso, Egypt, Kenya, Senegal and Sierra Leone. Similarly, in Burkina Faso, Ethiopia, Somalia and Sudan, girls and women who report both FGM and girl-child marriage represent the largest cohort. However, girls and women who report FGM without girl-child marriage constitute the second largest subgroup. Other studies in Somalia (World Vision UK, 2013, 2014) and Ethiopia (Boyden *et al.*, 2013) have also shown a direct relationship between the two practices, with FGM being seen as a transition of a girl from 'childhood' to 'adulthood', which then facilitates girl-child marriage. These studies show the similarity of causes or underlying drivers behind FGM and girl-child marriage.

Given the overall serious negative consequences of FGM and girl-child marriage, such as poor health, unintended or early pregnancy, pregnancy difficulties, multi-morbidity and mortality, providing updated empirical evidence on subnational variations and trends in the two variables, and assessing their potential linkages to indicators such as education, residential status, age, ethnicity and wealth quintile, could help map context-specific interventions and inform policy initiatives. Using DHS data from twelve selected countries in sub-Saharan Africa, this study sought to investigate the association between FGM and girl-child marriage by examining their possible correlates and making comparisons across selected countries.

Table 1. Description of surveys and study sample of women aged 20–24 years ($N=14,748$)

Country	Survey year(s)	Weighted n	Weighted %
1.Burkina Faso	2010	2669	18.10
2.Chad	2014–15	1272	8.63
3.Ethiopia	2016	726	4.92
4.Guinea	2018	1151	7.81
5.Kenya	2014	1408	9.55
6.Mali	2018	755	5.12
7.Niger	2012	622	4.22
8.Nigeria	2018	951	6.45
9.Senegal	2010–11	1750	11.87
10.Sierra Leone	2013	1566	10.62
11.Tanzania	2015–16	1260	8.54
12.Togo	2013–14	617	4.18

Methods

Data source

The study pooled DHS data from twelve sub-Saharan Africa countries covering the period from January 2010 to December 2018. The inclusion criteria in the selection of countries was that they had a recent DHS survey (2010–2018) with data on girl-child marriage and FGM. The women's file (IR) which gathers data from women aged 15–49 years on significant indicators related to children's and women's health, including girl-child marriage and FGM was used (Corsi *et al.*, 2012). The DHS uses a stratified sampling approach, which is done in two stages from a nationally representative sample. Details of the sampling process has been provided in a previous study (Aliaga & Ruilin, 2006). This study included a total of 14,748 women aged 20–24 years for whom there was complete information on the essential variables. Table 1 shows the weighted samples for each country.

Outcome variable

The outcome variable was child marriage, defined by the United Nations as marriage involving a person under the age of 18 (United Nations, 2014). This was obtained by asking respondents their marital status and their age at first marriage. A dichotomous variable was created and grouped by whether a respondent married as a child (<18 years) or otherwise. In this study, girl-child marriage is used because of the focus on girls as the study sample.

Explanatory variable

The independent variable was FGM, derived by asking respondents if their genital area 'was nicked with nothing removed', 'had something removed' or 'was sewn shut'. The responses were 'Yes' and 'No', coded No=0, and Yes=1.

Covariates

Eight covariates were considered in the study. These were broadly grouped into individual- and community-level variables. The individual-level variables included age, educational level,

employment status, wealth quintile and exposure to mass media. Age was categorized as 20, 21, 22, 23 and 24 years. Educational level was grouped into no education, primary and secondary/higher. Employment status was categorized as not working and working. In the DHS, wealth quintile is computed based on household assets and characteristics of the household using Principal Component Analysis (PCA) and grouped into poorest, poorer, middle, richer and richest. Exposure to media was derived from questions asked on frequency of listening to the radio, reading newspapers, watching television and access to the internet. Respondents who never listened to the radio, read newspapers or watched television and who had no access to the internet were considered as those with no exposure to mass media, and otherwise as those exposed to mass media (Fatema & Lariscy, 2020). Residence, community literacy level (proportion of women who can read and write) and community socioeconomic status (proportion of women in the richest household quintile) were considered as contextual-level factors. Residence was coded as 'urban' or 'rural'. Community literacy level and community socioeconomic status were each coded as 'low', 'middle' and 'high'. These were generated as the percentage of community members in the sampling cluster that were in the richest quintile and high literacy level aggregated at the national level using PCA. Details of how these were generated have been published in a previous study (Solanke *et al.*, 2019). The selection of variables was in line with their theoretical significance and practical importance for girl-child marriage (Hotchkiss *et al.*, 2016; Bezie & Addisu, 2019; Efevbera *et al.*, 2019).

Data analyses

The study followed four steps to analyse the data using Stata version 14.0. First, the prevalences of girl-child marriage and FGM in sub-Saharan Africa were computed using frequencies and percentages. Second, the distributions of girl-child marriage across FGM and the covariates were calculated. Third, the chi-squared test of independence (χ^2) was used to examine the association between girl-child marriage, FGM and the covariates, with significance set at 95% confidence interval ($p < 0.05$) (Table 2). Statistical significant variables at the χ^2 level were further analysed using a two-level multilevel logistic regression. As per the two-level modelling used in this study, women were nested within clusters to account for the variance in Primary Sampling Units (PSUs). Clusters were regarded as random effect to take care of the unexplained variability at the contextual level (Solanke *et al.*, 2019). Prior to the multilevel logistic regression analysis, a multicollinearity test was conducted among all variables that were significant at the χ^2 test using variance inflation factor (VIF). There was no indication of multicollinearity (Mean VIF=1.57, Maximum VIF=2.32, Minimum VIF=1.01).

Three models were fitted, with Model 0 showing the variance in the outcome variable attributed to the clusters in the absence of the explanatory variable and covariates. The second model (Model 1) had only FGM and the final model (Model 2) was the complete model that had FGM, the individual-level and contextual-level factors simultaneously. For model 1 and model 2, crude odds ratios (cORs) and adjusted odds ratios (aORs) with their associated 95% confidence intervals (CIs) are presented. To compare models, Akaike's Information Criterion (AIC) test was used (see Table 3). Sample weight (v005/1,000,000) was applied to correct for over- and under-sampling, while the *svy* command was used to account for the complex survey design and generalizability of the findings.

Results

The prevalence of FGM in the twelve countries was 52.19%, with the highest prevalence in Guinea (97.17%) and the lowest in Togo (4.57%) (Figure 1). The overall prevalence of girl-child marriage in the twelve countries was 57.96%, with the highest and lowest prevalences in Chad (78.06%) and Kenya (35.45%), respectively (Figure 2).

Table 2. Distribution of girl-child marriage among young women by FGM status and socio-demographic characteristics (N=14,748)

Variable	Weighted <i>n</i>	Weighted %	Girl-child marriage (%)	χ^2 (<i>p</i> -value)
Experienced FGM				244.1 (<0.001)
No	7051	47.8	50.4	
Yes	7697	52.2	64.9	
Age (years)				342.8 (<0.001)
20	3717	25.2	68.9	
21	2154	14.6	60.4	
22	3278	22.2	57.1	
23	2734	18.5	51.4	
24	2864	19.4	49.2	
Education				6.5 (<0.001)
No education	7523	51.0	70.5	
Primary	4001	27.1	54.9	
Secondary+	3223	21.9	32.5	
Employment status				0.6 (ns)
Not working	5394	36.6	57.0	
Working	9354	63.4	58.5	
Wealth status				593.1 (<0.001)
Poorest	2834	19.2	70.6	
Poorer	3009	20.4	66.8	
Middle	2897	19.7	61.6	
Richer	3061	20.8	51.9	
Richest	2947	20.0	39.5	
Mass media exposure				322.8 (<0.001)
No	4431	30.1	70.2	
Yes	10317	69.9	52.7	
Type of place of residence				401.7 (<0.001)
Urban	4435	30.1	41.9	
Rural	10313	69.9	64.6	
Community literacy level				724.8 (<0.001)
Low	6996	47.4	69.7	
Medium	3971	26.9	45.1	
High	3781	25.6	49.8	
Community socioeconomic status				340.5 (<0.001)
Low	9958	67.5	64.0	
Medium	215	1.5	67.0	
High	4575	31.0	44.3	

Table 3. Results of multilevel logistic regression analysis of the association between FGM and child marriage among young women ($N=14,748$)

Variable	Model 0	Model 1 aOR	Model 2 aOR (95% CI)
Fixed effects			
Experienced FGM			
No		0.53*** (0.51–0.57)	0.76*** (0.71–0.82)
Yes (Ref.)			
Age (years)			
20 (Ref.)			
21			0.75*** (0.67–0.85)
22			0.63*** (0.57–0.70)
23			0.53*** (0.47–0.59)
24			0.47*** (0.43–0.53)
Education			
No education (Ref.)			
Primary			0.69*** (0.63–0.76)
Secondary+			0.31*** (0.28–0.35)
Wealth			
Poorest (Ref.)			
Poorer			0.93 (0.83–1.05)
Middle			0.84** (0.75–0.94)
Richer			0.72*** (0.64–0.82)
Richest			0.56*** (0.47–0.66)
Mass media exposure			
No (Ref.)			
Yes			0.81*** (0.74–0.88)
Type of place of residence			
Urban			0.92 (0.74–0.88)
Rural (Ref.)			
Community literacy level			
Low (Ref.)			
Medium			0.63*** (0.57–0.69)
High			0.95 (0.85–1.05)
Community socioeconomic status			
Low			0.67* (0.49–0.92)
Medium (Ref.)			
High			0.75 (0.54–1.04)
Random effect	Model 0	Model 1	Model 2

(Continued)

Table 3. (Continued)

Variable	Model 0	Model 1 aOR	Model 2 aOR (95% CI)
PSU variance (95% CI)	0.06 (0.04–0.10)	0.05 (0.03–0.09)	0.03 (0.01–0.07)
ICC	1.8%	1.6%	0.9%
Likelihood ratio test	$\chi^2=24.33, p<0.001$	$\chi^2=18.73, p<0.001$	$\chi^2=5.49, p=0.0096$
Wald's χ^2	Reference	237.21	1664.86
Log-likelihood	-9999.84	-9880.47	-9026.7533
AIC	20,003.68	19,766.95	18,091.51
N	14,748	14,748	14,748

Ref.=Reference category; aOR=Adjusted Odds Ratio; PSU=Primary Sampling Unit; ICC=Intra-Class Correlation; AIC=Akaike's Information Criterion.

* $p<0.05$; ** $p<0.01$; *** $p<0.001$.




Figure 1. Prevalence (%) of FGM in twelve countries of sub-Saharan Africa.

Distribution of girl-child marriage by FGM and women's socio-demographic characteristics

Table 2 shows the distribution of girl-child marriage by FGM and socio-demographic characteristics of the sample women. More than 50% of the women had experienced FGM. Out of the total sample, 25.2% were aged 20 and 51% had no formal education. Approximately 63% of the young women had an occupation, 20.8% were in richer wealth quintile households and 69.9% had exposure to mass media. In addition, 69.9% lived in rural areas, 47.4% in communities with low literacy levels and 67.5% in communities with low socioeconomic status. Results of the χ^2 test indicated




Figure 2. Prevalence (%) of girl-child marriage in twelve countries of sub-Saharan Africa.

that all the explanatory variables had significant associations with girl-child marriage, except employment status.

Measures of association

Table 3 shows the results of the multilevel regression analysis of the relationship between FGM and girl-child marriage among the sample women. The complete model (Model 2) presents results of the relationship between FGM and girl-child marriage, after controlling for the effect of both individual- and community-level factors. The results indicate that women who had never experienced FGM were less likely to experience girl-child marriage (aOR=0.76, CI=0.71–0.82) compared with those who had experienced FGM.

In terms of the individual-level factors, the odds of girl-child marriage were lower among women aged 24 years at the time of the survey compared with those aged 20 years (aOR=0.47, CI=0.43–0.52). Also, women with secondary education were less likely to experience girl-child marriage compared with those with no formal education (aOR=0.31, CI=0.28–0.35). The odds of girl-child marriage reduced with increasing wealth quintile, with women belonging to the richest wealth quintile less likely to experience girl-child marriage compared with those of the poorest wealth quintile (aOR=0.56, CI=0.47–0.66). Furthermore, the odds of girl-child marriage were lower among women who had exposure to mass media compared with those who were not exposed to mass media (aOR=0.81, CI=0.74–0.88). Of the community-level factors, a lower odds of girl-child marriage was found among women who lived in communities with medium literacy

level compared with those in communities with low literacy level (aOR=0.63, CI=0.57–0.69), and those who lived in communities with low socioeconomic status compared with those in communities of medium socioeconomic status (aOR=0.67, CI=0.49–0.92).

Discussion

This study investigated the association between FGM and girl-child marriage in sub-Saharan Africa. It was conducted with the intent of providing evidence of the persistence and shifts in both practices that might trigger transformative actions to enhance the elimination of the acts. Overall, the findings revealed a greater than average prevalence of both FGM (52.19%) and girl-child marriage (57.96%) in sub-Saharan Africa, with the highest prevalence in Guinea for FGM (9 in 10 persons) and Chad for girl-child marriage (8 in 10 persons). These findings are in pursuant with a study conducted in Tanzania, which concluded that the prevalences of FGM and girl-child marriage were high in the country and that the rate was similar to that of the current study (Avalos *et al.*, 2015). A study conducted in Guinea also showed a similar results (Van Rossem & Gage, 2009). Similarly, a study on girl-child marriage in 22 sub-Saharan African countries also indicated that Chad has a high prevalence of girl-child marriage (77.90%) (Yaya *et al.*, 2019). Although recent situational reports (UNICEF, 2013, 2016) have noted declines in both FGM and girl-child marriage in sub-Saharan Africa, the overall prevalence noted in this study is worrisome, especially for Guinea and Chad. This could be driven by deeply rooted socio-cultural norms and social pressures, which facilitate these harmful practices through enforcement or even manipulation (Ohchr *et al.*, 2008). Educational interventions (e.g. advocacy campaigns) and law enforcement could help minimize these harmful practices.

In this study, women who did not experience FGM were less likely to experience girl-child marriage compared with their counterparts who did experience FGM. Girl-child marriage has been found to be linked to FGM in previous studies (Sakeah *et al.*, 2018; Karumbi & Gathara, 2020). This supports the view that FGM is a major corridor to girl-child marriage among women in Africa (Karumbi & Gathara, 2020). Besides, a commonly held assumption in sub-Saharan Africa is that a woman who is circumcised is less likely to be promiscuous compared with other leading outcomes associated with the FGM practice (Varol *et al.*, 2014). It could be that the purpose of girl-child marriage and FGM is to help preserve virginity and control girls' sexuality for marriage (Karumbi & Gathara, 2020).

Previous studies on the individual-level factors influencing girl-child marriage have found age of the respondent, educational attainment, wealth quintile and media exposure to be major socio-demographic factors (Sekine & Hodgkin, 2017; Rumble *et al.*, 2018; Okiyi *et al.*, 2020). The present study found that women aged 21–24 years at the time of the survey were less likely be married before the age of 18 years than those aged 20 years. Respondents with a higher educational level, those in the higher wealth quintile and those with media exposure were less likely to experience girl-child marriage. Girls who marry early (i.e. before or at the age of 18) will not have completed their secondary education, and hence they are more likely to drop out of school. More highly educated girls are more likely to marry later than less educated girls. Previous studies in Asia (e.g. Singh & Vennam, 2016) and Ethiopia (e.g. Erulkar *et al.*, 2010; Presler-Marshall *et al.*, 2016) found education to be positively related to greater decision-making among girls (Boyden *et al.*, 2013). Education facilitates the empowerment of girls and women, making them more aware of their rights, which might delay marriage (Karumbi & Gathara, 2020).

Women living in communities with medium literacy levels were less likely to experience girl-child marriage, as were those living in low socioeconomic status communities. Community women's literacy level often predicts women's susceptibility to early marriage, and economically autonomous women have a greater tendency not to consent to girl-child marriage (Paul, 2019). Community literacy level is thus a major driver of women's autonomy and

empowerment (Duflo, 2012). Unlike the findings of previous studies indicating that women living in lower socioeconomic status communities are more likely to experience girl-child marriage (Parsons *et al.*, 2015; Hotchkiss *et al.*, 2016), the current study found that women residing in communities with medium socioeconomic status were more likely to experience girl-child marriage. Methodological and context-specific variations could account for these inconsistent results. Future research could examine trends over time to provide a better understanding based on context-variations associated with girl-child marriage taking into account other socio-cultural determinants.

The main strength of this study lies in its use of nationally representative data and its rigorous analysis. The large sample size also allowed generalization of the findings to the entire population of the countries included in this study. However, the study used cross-sectional survey data, which limits the ability to make causal inferences from the findings. Also, social desirability bias may have resulted in under- or over-reporting on sensitive issues such as FGM and girl-child marriage.

In conclusion, this study has demonstrated a significant relationship between FGM and girl-child marriage with educational attainment, mass media exposure and community literacy level in sub-Saharan Africa. Tackling these factors could help eliminate girl-child marriage and FGM. Interventions should focus on girls' socio-demographic needs, community socioeconomic level, sexual and reproductive health service provision, and information campaigns against FGM. Policies to eliminate these practices should focus on compulsory basic education, poverty alleviation and increased access to mass media. Further campaigns should cover more of communities with lower literacy level and medium socioeconomic status. These steps will not only help stop girl-child marriage but also help reduce early pregnancies, maternal and child morbidity as well as mortality.

Acknowledgments. The authors are grateful to the MEASURE DHS project for granting free access to their original data. The dataset is freely available for download on <https://dhsprogram.com/data/available-datasets.cfm>

Author contributions. BOA conceived the study, performed the analysis and wrote up the data and methods. BOA, JEH, AS, OAB, EB, CA and TS wrote the first draft of the manuscript. All authors revised and proof-read the manuscript for its scholarly content and gave consent for the final version to be published.

Funding. The authors sincerely thank Bielefeld University, Germany, for providing financial support through the Open Access Publication Fund for the article processing charge.

Conflicts of Interest. The authors declare that they have no conflicts of interest. There are no financial, copyright, trademarks or patent implications arising from this research and no organization has any vested interest in this research.

Ethical Approval. This study used data from DHS datasets, which are publicly available. ICF international and an Institutional Review Board (IRB) in each respective country approved all the DHS surveys in line with the US Department of Health and Human Services regulations for the protection of human subjects. More details regarding DHS data and ethical standards are available at: <http://goo.gl/ny8T6X>.

References

- Ahinkorah BO, Hagan JE, Ameyaw EK, Seidu AA, Budu E, Sambah F *et al.* (2020) Socio-economic and demographic determinants of female genital mutilation in sub-Saharan Africa: analysis of data from demographic and health surveys. *Reproductive Health* 17(1), 1–14.
- Aliaga A and Ruilin R (2006) *Cluster optimal sample size for Demographic and Health Surveys*. DHS Working Papers No. 30. ORC Macro, Calverton, MD.
- Avalos L, Farrell N, Stellato R and Werner M (2015) Ending female genital mutilation and child marriage in Tanzania. *Fordham International Law Journal* 38, 639.
- Bezie M and Addisu D (2019) Determinants of early marriage among married women in Injibara Town, North West Ethiopia: community-based cross-sectional study. *BMC Women's Health* 19(1), 134.
- Boyden J, Pankhurst A and Tafere Y (2012) Child protection and harmful traditional practices: female early marriage and genital modification in Ethiopia. *Development in Practice* 22(4), 510–522.

- Boyden J, Pankhurst A and Tafere Y** (2013) *Harmful Traditional Practices and Child Protection: Contested Understandings and Practices of Female Child Marriage and Circumcision in Ethiopia*. Young Lives, Oxford.
- Corsi DJ, Neuman M, Finlay JE and Subramanian SV** (2012) Demographic and Health Surveys: a profile. *International Journal of Epidemiology* **41**(6), 1602–1613.
- Duflo E** (2012) Women empowerment and economic development. *Journal of Economic Literature* **50**(4), 1051–1079.
- Efevbera Y, Bhabha J, Farmer P and Fink G** (2019) Girl child marriage, socioeconomic status, and undernutrition: evidence from 35 countries in Sub-Saharan Africa. *BMC Medicine* **17**(1), 55.
- Erulkar A, Ferede A, Ambelu W, Girma W, Amdemikael H, GebreMedhin B et al.** (2010) *Ethiopia Young Adult Survey: A Study in Seven Regions*. Population Council, Addis Ababa.
- Fatema K and Lariscy JT** (2020) Mass media exposure and maternal healthcare utilization in South Asia. *SSM-Population Health* **100**, 614.
- Hotchkiss DR, Godha D, Gage AJ and Cappa C** (2016) Risk factors associated with the practice of child marriage among Roma girls in Serbia. *BMC International Health and Human Rights* **16**(1), 6.
- Karumbi J and Gathara D** (2020) *Associations Between Female Genital Mutilation/Cutting and Early/Child Marriage: A Multi-country DHS/MICS Analysis*. Population Council, New York.
- Karumbi J, Gathara D and Muteshi J** (2017) *Exploring the Association Between FGM/C and Child Marriage: A Review of the Evidence. Evidence to End FGMC. Programme Consortium Report*. Population Council, New York.
- Karumbi J and Jacinta M** (2017) Exploring the associations between FGM/C and early/child marriage: a review of the evidence. *Female Genital Mutilation/Cutting* **75**, 35–81
- Mackie G and LeJeune J** (2009) *Social Dynamics of Abandonment of Harmful Practices: A New Look at the Theory*. UNICEF, New York.
- Ohchr U, Undp U and Unesco U** (2008) *Eliminating Female Genital Mutilation: An Interagency Statement*. WHO, Geneva.
- Okiyi G, Odionye C and Okeya A** (2020) Socio-cultural variables and media coverage of girl child marriages. *Nnamdi Azikiwe University Journal of Communication and Media Studies* **1**(2), 1–15
- Parsons J, Edmeades J, Kes A, Petroni S, Sexton M and Wodon Q** (2015) Economic impacts of child marriage: a review of the literature. *Review of Faith & International Affairs* **13**(3), 12–22.
- Paul P** (2019) Effects of education and poverty on the prevalence of girl child marriage in India: a district-level analysis. *Children and Youth Services Review* **100**, 16–21.
- Population Reference Bureau** (2018) *Exploring the Association Between Female Genital Mutilation/Cutting and Early/Child Marriage. Evidence to End FGM/C. Programme Consortium Evidence Brief*. Population Reference Bureau.
- Presler-Marshall E, Lyytikainen M and Jones N** (2016) *Child Marriage in Ethiopia: A Review of the Evidence and an Analysis of the Prevalence of Child Marriage in Hotspot Districts*. Overseas Development Institute, London.
- Rumble L, Peterman A, Irdiana N, Triyana M and Minnick E** (2018) An empirical exploration of female child marriage determinants in Indonesia. *BMC Public Health* **18**(1), 407.
- Sakeah E, Debpuur C, Oduro AR, Welaga P, Aborigo R, Sakeah JK and Moyer CA** (2018) Prevalence and factors associated with female genital mutilation among women of reproductive age in the Bawku municipality and Pusiga District of northern Ghana. *BMC Women's Health* **18**(1), 1–10.
- Sekine K and Hodgkin ME** (2017) Effect of child marriage on girls' school dropout in Nepal: analysis of data from the Multiple Indicator Cluster Survey 2014. *PLoS One* **12**(7), e0180176.
- Singh R and Vennam U** (2016) *Factors Shaping Trajectories to Child and Early Marriage: Evidence from Young Lives in India*. Young Lives, Oxford.
- Solanke BL, Oyinlola FF, Oyeleye OJ and Ilesanmi BB** (2019) Maternal and community factors associated with unmet contraceptive need among childbearing women in Northern Nigeria. *Contraception and Reproductive Medicine* **4**(1), 11.
- UNFPA** (2012) *Marrying Too Young: End Child Marriage*. URL: www.unfpa.org/sites/default/files/pub-pdf/MarryingTooYoung.pdf (accessed 15th September 2020).
- UNFPA** (2014) *Girls Not Brides Fact Sheet*. URL: www.girlsnotbrides.org/wpcontent/uploads/2014/10/GNB-factsheeton-child-marriage-numbers-Oct-2014.pdf (accessed 15th September 2020).
- UNICEF** (2013) *Female Genital Mutilation/Cutting: A Statistical Overview and Exploration of the Dynamics of Change*. URL: www.unicef.org/media/files/UNICEF_FGM_report_July_2013_Hi_res.pdf (accessed 15th September 2020).
- UNICEF** (2016) *Female Genital Mutilation/Cutting: A Global Concern*. URL: www.unicef.org/media/files/FGMC_2016_brochure_final_UNICEF_SPREAD.pdf (accessed 10th September 2020).
- United Nations** (2014) *Annual Report of the United Nations High Commissioner for Human Rights*. United Nations, New York.
- United Nations Office of the High Commissioner on Human Rights** (2015) *Harmful Practices, Especially Forced Marriage and Female Genital Mutilation (Tech. Rep.)*. URL: https://www.ohchr.org/Documents/Issues/Women/WRGS/SexualHealth/INFO_Harm_Pract_WEB.pdf (accessed 15th September 2020).
- Van Rossem R and Gage AJ** (2009) The effects of female genital mutilation on the onset of sexual activity and marriage in Guinea. *Archives of Sexual Behavior* **38**(2), 178–185.

- Varol N, Fraser IS, Ng CHM, Jaldesa G and Hall J** (2014) Female genital mutilation/cutting – towards abandonment of a harmful cultural practice. *Australian and New Zealand Journal of Obstetrics and Gynaecology* **54**(5), 400–405.
- Williams-Breault BD** (2018) Eradicating female genital mutilation/cutting: human rights-based approaches of legislation, education, and community empowerment. *Health and Human Rights* **20**(2), 223.
- World Vision UK** (2013) Untying the Knot: Exploring Early Marriage in Fragile States. URL: www.girlsnotbrides.org/reports-and-publications/untying-the-knot-exploring-early-marriage-infragile-states (accessed 15th September 2020).
- World Vision UK** (2014) Exploring the Links: Female Genital Mutilation/Cutting and Early Marriage. URL: http://worldvision.org.uk/files/4814/0068/7160/Exploring_the_links_FGM_cutting_and_early_marriage.pdf (accessed 15th September 2020).
- Yaya S, Odusina EK and Bishwajit G** (2019) Prevalence of child marriage and its impact on fertility outcomes in 34 sub-Saharan African countries. *BMC International Health and Human Rights* **19**(1), 33.

Cite this article: Ahinkorah BO, Hagan JE, Seidu A-A, Bolarinwa OA, Budu E, Adu C, Okyere J, Archer AG, and Schack T. Association between female genital mutilation and girl-child marriage in sub-Saharan Africa. *Journal of Biosocial Science*. <https://doi.org/10.1017/S0021932021000687>