Counting stabiliser codes for arbitrary dimension

Tanmay Singal, ^{1, a)} Che Chiang, ^{2, b)} Eugene Hsu, ^{3, c)} Eunsang Kim, ^{4, d)} Hsi-Sheng Goan, ^{2, 5, 6, e)} and Min-Hsiu Hsieh ^{7, f)}

In this work, we compute the number of $[[n,k]]_d$ stabiliser codes made up of d-dimensional qudits, for arbitrary positive integers d. In a seminal work by Gross (Ref. 22) the number of $[[n,k]]_d$ stabilizer codes was computed for the case when d is a prime (or the power of a prime, i.e., $d = p^m$, but when the qudits are Galois-qudits). The proof in Ref.²² is inapplicable to the non-prime case. For our proof, we introduce a group structure to $[n,k]_d$ codes, and use this in conjunction with the Chinese remainder theorem to count the number of $[[n,k]]_d$ codes. Our work overlaps with Ref.²² when d is a prime and in this case our results match exactly, but the results differ for the more generic case. Despite that, the overall order of magnitude of the number of stabilizer codes scales agnostic of whether the dimension is prime or non-prime. This is surprising since the method employed to count the number of stabilizer states (or more generally stabilizer codes) depends on whether d is prime or not. The cardinality of stabilizer states, which was so far known only for the prime-dimensional case (and the Galois qudit prime-power dimensional case) plays an important role as a quantifier in many topics in quantum computing. Salient among these are the resource theory of magic, design theory, de Finetti theorem for stabilizer states, the study and optimisation of the classical simulability of Clifford circuits, the optimal verification of stabilizer states, the study of quantum contextuality of small-dimensional systems and the study of Wigner-functions. Our work makes available this quantifier for the generic case, and thus is an important step needed to place results for quantum computing with non-prime dimensional quantum systems on the same pedestal as prime-dimensional systems.

I. INTRODUCTION

The stabilizer formalism (which will be explained in detail in Section II) has become an indispensable part of the study of quantum computing. Some of its salient applications are as follows: it forms the bedrock for the vast field of quantum error correcting codes (QECC)²⁹. Randomising over stabilizer states or Clifford unitaries (see Eq. (18) for the definition of a Clifford unitary) serves as an important application, for instance, for computing capacities of quantum channels¹, data hiding² and the study of noise-compounding in quantum circuits (randomized benchmarking)^{3,4}. It also demarcates a boundary between the classical simulability of quantum computations via the Gottesmann-Knill theorem⁵, and the onset of quantum complexity for which some non-stabilizerness is necessary. The non-universal nature of stabilizer operations⁶ and the need to compensate this deficiency with non-stabilizer operations manifests in a resource theory of "non-stabilizerness", a.k.a., magic (see

Ref. 44-46), where stabilizer operations are the free resource (see Subsection I A for an explanation of stabilizer operations and the resource theory with respect to which it is defined). While the basic mathematical preliminaries which support all the above have been developed in many works (for e.g. see Ref.⁸, ²²), there remain some gaps, particularly for quantum computing with multiqudit systems for arbitrary d. And while qubit systems are envisaged as the paradigmatic building blocks of quantum computing, qudits of larger dimensions may offer their own benefits (for e.g. see Ref. 9-14), owing to which it is imperative to study them. Among these, it is easier to study multiqudit systems when the dimension d is a prime number, or when d is the power of a prime, i.e., $d = p^m$, while, simultaneously, the configuration space of the qudit system is the Galois field \mathbb{F}_{p^m} . We call such a qudit a Galois-qudit¹⁶. Many important results which were obtained for qubits may be easily generalized to such qudit systems (this has been noted in many places, for e.g. in the introduction in Ref.⁵⁷). This contrasts with qudit systems, whose configuration space is \mathbb{Z}_d , when d is not a prime number. We refer to such qudit systems simply as qudits since this work is about such qudits, but wherever disambiguation is required, we may instead refer to them as modular qudits¹⁷. While there has been a lot of seminal work on such modular qudit systems ^{18–27}, some significant gaps remain. For instance, one of those gaps is the computation of the number of $[[n,k]]_d$ stabilizer codes of such qudits (stabilizer codes are defined

¹⁾Institute of Physics, Faculty of Physics, Astronomy and Informatics, Nicolaus Copernicus University, Grudziadzka 5/7, 87-100 Toruń, Poland

²⁾Department of Physics and Center for Theoretical Physics, National Taiwan University, Taipei 10617, Taiwan

³⁾Quantum information center, Chung Yuan Christian University, No. 200, Zhongbei Rd., Zhongli Dist., Taoyuan City 320314, Taiwan

⁴⁾Department of Applied Mathematics, Hanyang University, Ansan Kyunggi-do, Korea

⁵⁾Center for Quantum Science and Engineering, National Taiwan University, Taipei 10617, Taiwan

⁶⁾Physics Division, National Center for Theoretical Sciences, Taipei, 10617, Taiwan

⁷⁾ Hon Hai Quantum Computing Research Center, Taipei, Taiwan

a)tanmaysingal@gmail.com Corresponding author

b)cchiang314@gmail.com

c)a13579230@gmail.com

d)eskim@hanyang.ac.kr

e)goan@phys.ntu.edu.tw

f)min-hsiu.hsieh@foxconn.com

in Subsection IID). We will label this number as $\mathscr{C}[[n,k]]_d$. $\mathscr{C}[[n,k]]_2$ was computed in a seminal paper Ref.⁶ (a recent proof also appeared in Ref.⁷), and for Galois-qudit systems, $\mathscr{C}[[n,k]]_{\mathbb{F}_{p^m}}$ was computed in another seminal paper, Ref.²². In this work we compute $\mathscr{C}[[n,k]]_d$ for arbitrary $d \geq 2$. Our work overlaps with earlier works^{6,7,22} for the case when d is a prime number, and in that case our results match exactly.

Typically one is interested in the order of magnitude of $\mathscr{C}[[n,k]]_d$ since it may act as an important quantifier. For such a reader, we present here the main result.

Corollary 1 (Corollary 4). Let c = 2.17. Then number of $[[n,k]]_d$ stabiliser codes scales as

$$d^{\frac{(n-k)(n+3k+1)}{2}} \ \leq \mathscr{C}[[n,k]]_d \ < \ d^{\frac{(n-k)(n+3k+1)}{2}+c}, \eqno(1)$$

the number of $[[n,k]]_d$ stabiliser code spaces scale as

$$d^{\frac{(n-k)(n+3k+3)}{2}} \leq \mathscr{C}_{\mathrm{space}}[[n,k]]_d < d^{\frac{(n-k)(n+3k+3)}{2}+c}, \quad (2)$$

and thus the number of stabilizer states scale as

$$d^{\frac{n(n+3)}{2}} \le \mathscr{C}_{\text{state}}[[n]]_d < d^{\frac{n(n+3)}{2}+c}.$$
 (3)

Here we make a distinction between an $[[n,k]]_d$ stabilizer code, and the corresponding stabilizer code spaces, which is one of the d^{n-k} subspaces associated with such a stabilizer code (see Remark 3).

The number of stabilizer states, i.e., $\mathcal{C}_{\text{state}}[[n]]_d$, plays the role of an important quantifier in quantum computing. Before this work, $\mathcal{C}_{\text{state}}[[n]]_d$ was known only for prime d (and for \mathbb{F}_{p^m} number systems), and, thus, its application as a quantifier was limited to these cases. We list below the salient topics where it has been used as a quantifier before.

A. Resource theory of magic

To initiate the interested reader into the resource theory of stabilizer-operations we refer them to a short summary in Appendix A. For more comprehensive treatments on the topic, see Ref. 45,46,48. For some recent interesting developments in the topic, we further refer the reader to Ref. 58.

The most significant application of $\mathcal{C}_{\text{state}}[[n]]_d$ is that stabilizer states are the extremal points of the stabilizer polytope, and many measures of magic are defined as optimizations over this polytope. Since $\mathcal{C}_{\text{state}}[[n]]_d$ scales super-exponentially for prime-dimensional systems (and Galois-qudit systems), computing these measures of magic is an intractable problem for such systems. The earliest known measure in this category is the relative entropy of magic 145 , after which the robustness

of magic was defined in Ref.⁴⁶. The robustness of magic has an operational interpretation: it gives an upper bound on the classical simulation complexity of the quantum Clifford+ $\pi/8$ circuits (see also Ref.⁴⁷), and scales exponentially in the number of $\pi/8$ gates. It may also be used for the optimality of $\pi/8$ -gate counts in the gate-synthesis problem. In Ref.⁴⁸ the regularised robustness of magic was introduced, and its computation, while superpolynomially faster, still suffers from an exponential run-time. We also refer the reader to a recent work Ref.⁵⁰ which tries to optimize the classical simulability of Clifford+ $\pi/8$ circuits, in the manner introduced by Ref.⁴⁷. The intractability of these measures has also encouraged the search for new quantifiers of magic, for instance, see Ref.⁴⁹.

Before our work, it wasn't known how $\mathscr{C}_{\text{state}}[[n]]_d$ scales with n for arbitrary d. Since Corollary 1 states that $\mathscr{C}_{\text{state}}[[n]]_d$ scales super-exponentially in n for all d, we now know that the problem of computing the aforementioned measures of magic is intractable for arbitrary dimension.

B. STIM: The classical simulation of stabilizer operations

The resource theoretic perspective of Subsection I A, which deems stabilizer operations to be a free resource, is an approximation, albeit a useful one. These costs are apparent while simulating algorithms with only stabilizer operations. Such a classical simulation requires one to keep track of $\frac{1}{2}d^2$ dits (where 'dit' is the *d*-level analog of the two-level bit) during the computation. Appropriate optimisation schemes are required for such simulations, particularly, when the goal is to simulate large quantum circuits, for instance a circuit for the surface code with fifteen thousand qubits (for e.g., see Ref. 51).

C. Frame potentials for stabilizer states

In Ref.⁵², Kueng and Gross established that a uniform ensemble of multiqubit stabilizer states are complex projective 3-designs. To prove this, they computed explicitly the frame-potential associated with this ensemble of states. And this frame potential was computed using $\mathcal{C}_{\text{state}}[[n]]_2$, which was borrowed from Ref.⁶ and Ref.²². While for $d \geq 3$, it is known that multi-qudit stabilizer states aren't projective 3-designs, one may nevertheless still be interested in the projective-design which this ensemble gives rise to. While the result in Ref.²² may be employed for this purpose for only prime values of d, our results allows one to obtain the result for arbitrary values of d.

D. de Finetti theorem for stabilizer states

In Ref.⁵³ a de Finetti theorem for stabilizer states was established for quantum states on *t*-copes of *n*-qudit systems, where *d* is prime. To summarize it, let us consider the action of the *t*-th tensor power of the *n*-qudit Clifford group on *t*-copies of an *n*-qudit system, i.e., $U^{\otimes t}$, where $U \in \mathscr{C}_n^{(d)}$, and

¹ In Ref. ⁴⁵, a computable measure of magic called *mana* was also introduced. This measure, while efficiently computable, is computable for only odd-dimensional quantum systems. That being said it has a more significant operational interpretation compared to the relative entropy of magic.

let $\mathscr L$ be the commutant of this tth tensor power of the Clifford group, i.e., $L \in \mathscr L$ implies that $U^{\otimes t}L(U^\dagger)^{\otimes t}=L$. Let a quantum state ρ commute with all L in $\mathscr L$. Then ρ has the following property. Define $\rho_s=\mathrm{Tr}_{i_1,i_2,\cdots i_{t-s}}\rho$ be the reduced state, which is obtained by tracing out t-s subsystems from ρ . The subsystems which are traced out are arbitrary. Then there exists some probability distribution p_{σ_S} on the n-qudit stabilizer states σ_S such that

$$\frac{1}{2} \left\| \rho_{s} - \sum_{\sigma_{s}} p_{\sigma_{s}} \sigma_{s}^{\otimes s} \right\|_{1} \leq C d^{\mathscr{O}(n^{2})} d^{-\frac{1}{2}(t-s)}, \quad (4)$$

where C is some constant, σ_S is a stabilizer state of an n-qudit systems and p_{σ_S} is a probability distribution on these states. The exponential scaling in the number of traced out subsystems, i.e., t-s contrasts with the ordinary de Finetti result, whose scaling is of the form $\mathcal{O}(s/t)$ (see Ref. 54). The exponential scaling of the stabilizer version of the de Finetti theorem is proved using $\mathcal{C}_{\text{state}}[[n]]_d$, which was computed in Ref. 22 for prime d (and Galois qudits). We anticipate it to be possible for Corollary 1 to provide a similar exponential scaling for the case when d is non-prime.

E. Others

We mention some other topics where $\mathscr{C}_{\text{state}}[[n]]_d$ for prime d was employed. We hope that our result could find similar applications for non-prime d.

- (i) Quantum contextuality: In Ref. 55, Howard and collaborators study quantum stabilizer states, to see if these states exhibit any quantum contextual correlations. For odd prime d, the set of $\mathcal{C}_{\text{state}}[[n]]_d$ two-qudit stabilizer states is partitioned into two classes: separable states and entangled states. Each class is separately studied to see if it exhibits state dependent contextuality, and the number of elements within each class plays an important role in this.
- (ii) Disambiguation of Wigner functions for odd dimensional qudit systems: While Hudson's theorem for finite dimensional quantum systems was proved in Ref.²², the choice of the Wigner function with which it was proved wasn't disambiguated. In Ref.⁵⁶, Howard establishes that there is a unique choice of the Wigner function which supports Hudson's theorem. The proof of this result employs the explicit formula for $\mathscr{C}_{\text{state}}[[n]]_d$.
- (iii) Optimal verification of quantum stabilizer states. In Ref. 57 , optimal schemes to verify stabilizer states are given. Knowing $\mathscr{C}_{\text{state}}[[n]]_d$, or at least the order of magnitude of $\mathscr{C}_{\text{state}}[[n]]_d$ is essential for this task. While the results in Ref. 57 are given for qubit systems, Dangniam et al. believe that those results are generalizable to the case when d is an arbitrary prime number.

Our work readily generalises results for the resource theory of magic and for the classical simulability of stabilizer-only circuits. For other topics mentioned above, we anticipate that our computation of $\mathcal{C}_{\text{state}}[[n]]_d$ will facilitate comparable results for arbitrary d-dimensional systems.

The challenge in computing $\mathscr{C}[[n,k]]_d$, for non-prime d, is that \mathbb{Z}_d is not a field. In particular, only elements which are co-prime with d will have multiplicative inverses. The phase space of an n-modular qudit system is the 2n-fold Cartesian product of \mathbb{Z}_d , i.e., \mathbb{Z}_d^{2n} . We need to treat it like a vector space which is defined over a field. Towards this end, we will borrow some basics terms and definitions from the theory of vector spaces wherever this is appropriate. We stress though, that the jargon we borrow from linear algebra is actually superfluous, and all the necessary concepts can be phrased in terms of more primitive concepts in the theory of abelian groups and its subgroups. Nevertheless, a rigorous justification of viewing \mathbb{Z}_d^{2n} as a vector space and applying concepts and definitions from the theory of vector spaces will be justified in the appendix.

 $\mathscr{C}[[n,k]]_d$ is computed in two steps. (i) We introduce a group theoretic structure of $[[n,k]]_d$ stabilizer codes constructed in the following way: we take a $2n \times (n-k)$ check-matrix of a given code, and extend it to a $2n \times 2n$ symplectic matrix. This extension isn't unique, and the set of all such symplectic extensions realizable for a given $[[n,k]]_d$ code forms a unique coset of $Sp(2n,\mathbb{Z}_d)/T(2n,k,\mathbb{Z}_d)$, where $T(2n, k, \mathbb{Z}_d)$ is a subgroup of matrices in $Sp(2n, \mathbb{Z}_d)$ which forms the coset corresponding to the $[[n,k]]_d$ trivial code. Conversely, to each coset of $Sp(2n, \mathbb{Z}_d)/T(2n, k, \mathbb{Z}_d)$ one can attribute a unique $[n,k]_d$ stabilizer code, and thus a bijection between $[[n,k]]_d$ stabilizer codes and cosets of $\operatorname{Sp}(2n,\mathbb{Z}_d)/\operatorname{T}(2n,k,\mathbb{Z}_d)$ is established. This tells us that $\mathscr{C}[[n,k]]_d = |\operatorname{Sp}(2n,\mathbb{Z}_d)|/|\operatorname{T}(2n,k,\mathbb{Z}_d)|$. (ii) To compute $|T(2n,k,\mathbb{Z}_d)|$, we find a way to decompose each element of $T(2n,k,\mathbb{Z}_d)$ into an ordered product of four matrices, each of which belongs to a distinct subgroup of $T(2n, k, \mathbb{Z}_d)$. These subgroups overlaps only over the identity element. Thus $|T(2n, k, \mathbb{Z}_d)|$ is the product of the orders of these subgroups. The orders of these subgroups and of $|\operatorname{Sp}(2n,\mathbb{Z}_d)|$ are computed, first for the case when $d = p^m$, i.e., the power of a prime. This corresponds to the case when the configuration space of the qudit is \mathbb{Z}_{p^m} (and not \mathbb{F}_{p^m}). Some of the details of this computation, which employ standard tricks for such computations, are relegated to Section E1 in the Appendix. This gives us $\mathscr{C}[[n,k]]_{p^m}$. To obtain $\mathscr{C}[[n,k]]_d$, we invoke the Chinese remainder theorem, which is a ring isomorphism, as follows: $\mathbb{Z}_d \simeq \mathbb{Z}_{p_1}^{m_1} \times \mathbb{Z}_{p_2}^{m_2} \times \cdots \times \mathbb{Z}_{p_r}^{m_r}$, where $d = p_1^{m_1} p_2^{m_2} \cdots p_r^{m_r}$ is the prime factorisation of d. While the proof method depends on whether d is prime or non-prime, the number of stabilizer codes $\mathscr{C}[[n,k]]_d$, the stabilizer code subspaces $\mathscr{C}_{\text{space}}[[n,k]]_d$ and the stabilizer states $\mathscr{C}_{\text{state}}[[n]]_d$ scale agnostic of d being prime or non-prime, which is surprising.

The paper is organised as follows. In Section II we give the necessary preliminaries of the stabilizer formalism: we will introduce the *n*-qudit Weyl-Heisenberg group (also

known as the generalized Pauli group), we define stabilizer codes along with their check matrices and related concepts. In Section III, we explain the group theoretical structure of $[[n,k]]_d$ stabilizer codes, and in Section IV we show how an arbitrary element of $T(2n,k,\mathbb{Z}_d)$ may be decomposed into a product of elements from four different subgroups. In Section V we count $\mathscr{C}[[n,k]]_d$. We conclude in Section VI.

II. PRELIMINARIES

Various preliminaries are introduced in this section, along with the notations and explanations of their usage.

A. The ring Z_d

Let \mathbb{Z} denote the set of all integers. Let d be an arbitrary positive integer such that $d \geq 2$. \mathbb{Z}_d then denotes the set of integers modulo d: $\mathbb{Z}_d = \{0, 1, \cdots, d-1\}$. All arithmetic operations in \mathbb{Z}_d are carried out modulo d. This means that for any $a, b \in \mathbb{Z}_d$,

$$a \oplus_d b = (a+b) \mod d$$

 $a \times_d b = (a \cdot b) \mod d,$ (5)

where, on the left hand side of Eq. (5), \oplus_d and \times_d represent addition and multiplication operations in \mathbb{Z}_d , and on the right hand side, + and \cdot represent addition and multiplication in \mathbb{Z} . From here onward, we will use the standard notations of addition and multiplication, even in \mathbb{Z}_d , and let the context determine where these operations are meant to be performed. \mathbb{Z}_d is a cyclic group under addition, with 0 as the additive identity. When d is a composite number, for $a \in \mathbb{Z}_d$ to have a multiplicative inverse in \mathbb{Z}_d , a and d in \mathbb{Z} need to be co-prime. The set of all non-zero coprime elements form the multiplicative group \mathbb{Z}_d^{\times} . Hence, \mathbb{Z}_d is a field if and only if d is a prime number. When d is a prime, we will denote it by p and denote the corresponding field by \mathbb{F}_p .

B. Using \mathbb{Z}_d^n as a vector space

Let n be a positive integer. The n-fold Cartesian product of \mathbb{Z}_d with itself, $\mathbb{Z}_d \times \mathbb{Z}_d \times \cdots \mathbb{Z}_d$ contains all ordered n-tuples (a_1,a_2,\cdots,a_n) where $a_j \in \mathbb{Z}_d$. Addition in \mathbb{Z}_d^n is defined point-wise: for arbitrary $a,b \in \mathbb{Z}_d^n$, $(a+b)_i = a_i + b_i$, where $(a+b)_i$ denotes the i-th component of a+b. Similarly, scalar multiplication of a in \mathbb{Z}_d^n with some $\lambda \in \mathbb{Z}_d$ means $\lambda a = (\lambda a_1, \lambda a_2, \cdots, \lambda a_n)$. \mathbb{Z}_d^n satisfies all the axioms which a usual vector space will satisfy. Thus, any $a = (a_1, \cdots, a_n) \in \mathbb{Z}_d^n$ will be called a vector. That being said, since \mathbb{Z}_d is not necessarily a field, \mathbb{Z}_d^n doesn't form a vector space. Technically, \mathbb{Z}_d^n is called a module over a commutative ring²⁸. The algebra in \mathbb{Z}_d^n is richer than that for vector spaces defined over fields, and one may not blindly generalise results from vector spaces over fields to \mathbb{Z}_d^n . But we will be able to borrow the following concepts from linear algebra.

Definition 1 (Linear independence). Let $a_1, a_2, \dots, a_m \in \mathbb{Z}_d^n$ be m vectors. Then they are linearly independent in \mathbb{Z}_d^n if and only if the only solution for the unknowns $x_1, x_2, \dots, x_m \in \mathbb{Z}_d$ in the following equation

$$x_1 a_1 + x_2 a_2 + \dots + x_m a_m = 0$$
(6)

is that $x_1, x_2, \dots, x_m = 0$.

We now list some corollaries which follow from Definition

Corollary 2. (i) Any set of n LI vectors forms a basis for \mathbb{Z}_d^n .

- (ii) If $\{a_1, a_2, \dots, a_m\}$ are LI, then their linear span generates a subspace of dimension m.
- (iii) For any m < n, any set of m LI vectors may be extended to form a basis for \mathbb{Z}_d^n .
- (iv) If the columns of an $n \times m$ matrix are LI, then viewed as a linear map from $\mathbb{Z}_d^n \to \mathbb{Z}_d^m$, its range is of dimension m and kernel is of dimension n-m.

Despite the fact that \mathbb{Z}_d^n is not necessarily a vector space, the following terms in Corollary 2: basis, linear span, subspace, dimension, matrix, linear map, range and kernel can be applied and used in the same manner as done when working with a vector space over fields. The reader is further directed to the Appendix B for a rigorous justification of these terms.

C. Generalized Pauli group on n qudits

We introduce the single qudit Pauli group $P^{(d)}$ using its defining representation, which acts on \mathbb{C}^d , where d is an arbitrary positive integer greater than or equal to 2. The construction of the defining representation of $P^{(d)}$ for arbitrary d has been done earlier in many works $^{18-27}$. Among these, we choose 21 . First, let us introduce an orthonormal basis for \mathbb{C}^d : $\{|j\rangle\}_{j=0}^{d-1}$, where the label j is taken from \mathbb{Z}_d . With respect to this basis, define the linear operators X and Z as follows.

$$X|j\rangle = |j+1\rangle, Z|j\rangle = \boldsymbol{\omega}^{j}|j\rangle, \ \forall \ j \in \mathbb{Z}_{d},$$
 (7)

where $\omega := \exp 2\pi i/d$, (d-th root of unity). Since $j \in \mathbb{Z}_d$, it is understood that the operation of addition in j+1 is performed in \mathbb{Z}_d . The commutation relations between X and Z are given by

$$ZX = \omega XZ.$$
 (8)

When d=2, $X=\begin{pmatrix} 0&1\\1&0 \end{pmatrix}$, and $Z=\begin{pmatrix} 1&0\\0&-1 \end{pmatrix}$, that is, we get the well-known Pauli matrices for the qubit case. When d is odd, $P^{(d)}:=\langle X,Z\rangle$. It is easily seen that an arbitrary element takes the form $\omega^j X^a Z^b$, where $j,a,b=0,1,\cdots,d-1$. Group composition is given by

$$\left(\omega^{j}X^{a}Z^{b}\right).\left(\omega^{j'}X^{a'}Z^{b'}\right) = \omega^{j+j'+a'b}X^{a+a'}Z^{b+b'}, \quad (9)$$

which is the just the composition rule of the Heisenberg-Weyl group²². The order of each element is at most d. For the case when d is even, often it is convenient² to introduce an additional phase factor $\zeta = \omega^{\frac{1}{2}}$, and $P^{(d)} := \langle \zeta \mathbb{1}, X, Z \rangle$. Then an arbitrary element in $P^{(d)}$ is $\zeta^j X^a Z^b$, where $j \in \mathbb{Z}_{2d}$, $a,b \in \mathbb{Z}_d$. The group composition law of two arbitrary elements is given by

$$\left(\zeta^{j}X^{a}Z^{b}\right).\left(\zeta^{j'}X^{a'}Z^{b'}\right) = \zeta^{j+j'+2a'b}X^{a+a'}Z^{b+b'}. \quad (10)$$

While the order of all group elements is at most 2d in the even d case, one may suitably multiply with ζ to generate another group element with order at most d. The n-qudit Pauli group $\mathscr{P}_n^{(d)}$ is simply the n-fold tensor product of the single qudit Pauli group. The Hilbert space it acts on will be denoted by $\mathscr{H} \simeq (\mathbb{C}^d)^{\otimes n}$. For the odd d case, $\mathscr{P}_n^{(d)} := \langle X_j, Z_j \rangle_{j=1}^n$, whereas for the even d case, we have that $\mathscr{P}_n^{(d)} = \langle \zeta \mathbb{1}, X_j, Z_j \rangle_{j=1}^n$, where X_j represents the operator with X on the j-th qudit and $\mathbb{1}$ on all the remaining qudits. Neglecting the phase factors of ω^j and ζ^j , we may represent an arbitrary element of the n-qudit Pauli group as

$$g(a) = X^{u_1} Z^{v_1} \cdots X^{u_n} Z^{v_n}, \tag{11}$$

where $a := (u, v)^T \in \mathbb{Z}_d^{2n}$ is a 2*n*-ordered tuple with entries in \mathbb{Z}_d . It is easily seen that group composition law is given by²¹

$$g(a)g(b) = \omega^{a^T U b} g(a+b), \tag{12}$$

where

$$U = \begin{bmatrix} 0 & 0 \\ I_n & 0 \end{bmatrix}, \tag{13}$$

and I_n is the $n \times n$ identity matrix. From Eq. (12), the commutation relation for the Pauli operators is

$$g(a)g(b) = \boldsymbol{\omega}^{-a^T \Lambda b} g(b)g(a), \tag{14}$$

where

$$\Lambda := \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}. \tag{15}$$

Eq. (14) implies that

$$a^T \Lambda b = 0 \tag{16}$$

if and only if the two Pauli operators g(a) and g(b) commute with each other. The centre of the Pauli group $\mathscr{P}_n^{(d)}$, which is the subgroup of $\mathscr{P}_n^{(d)}$ which commutes with all elements of $\mathscr{P}_n^{(d)}$, is

$$\mathscr{Z}\left(\mathscr{P}_{n}^{(d)}\right) = \langle \chi \mathbb{1} \rangle, \tag{17}$$

where we will assume that $\chi = \omega$ when d is odd and $\chi = \zeta$ when d is even. Thus the factor group $\mathscr{P}_n^{(d)}/\mathscr{Z}\left(\mathscr{P}_n^{(d)}\right) \simeq \mathbb{Z}_d^{2n}$. This is also seen from Eq. (12), since $g(a)g(b) \propto g(a+b)$. Thus $\mathscr{P}_n^{(d)}$ is homomorphic to \mathbb{Z}_d^{2n} , and one possible homomorphism takes Pauli $g(a) \in \mathscr{P}_n^{(d)}$ to $a \in \mathbb{Z}_d^{2n}$. This homomorphism plays a very important role in the formalism of the stabiliser codes.

The Clifford group $\mathcal{C}_n^{(d)}$ for an *n*-qudit system is defined as the normaliser of the Weyl-Heisenberg group in the unitary group over the *n*-qudit system.

$$U \in \mathscr{C}_n^{(d)} \Leftrightarrow UgU^{\dagger} \in \mathscr{P}_n^{(d)}, \, \forall \, g \in \mathscr{P}_n^{(d)}.$$
 (18)

The Clifford group is homomorphic to the group of $2n \times 2n$ symplectic matrices $Sp(2n, \mathbb{Z}_d)$ over \mathbb{Z}_d .

$$M \in \operatorname{Sp}(2n, \mathbb{Z}_d) \Leftrightarrow M^T \Lambda M = \Lambda,$$
 (19)

where Λ is given in Eq. (15).

D. Stabilizer codes on n gudits

The general framework of stabiliser codes was introduced in²⁹. Among the many references available in the literature, we refer the reader to³⁶, for a beginner friendly introduction to quantum error correction.

To define an $[[n,k]]_d$ stabiliser code space, we will need to first define a stabiliser group S. Let g_1,g_2,\cdots,g_{n-k} be n-k elements in $\mathscr{P}_n^{(d)}$, with the following properties. (i) they commute with each other, (ii) the order of $S \coloneqq \langle g_1,g_2,\cdots,g_{n-k}\rangle$ is $|S|=d^{n-k}$. The final condition ensures that any non-trivial product of the g_j 's, i.e., $g_1^{x_1}g_2^{x_2}\cdots g_{n-k}^{x_{n-k}}=\mathbb{1}$ if and only if $x_1,x_2,\cdots,x_{n-k}=0$. Often, one includes another condition, i.e., spectrum of each g_j always contains +1. If any g_j doesn't satisfy this condition, one may replace g_j with $\chi^a g_j$, so that $\chi^a g_j$ has eigenvalues +1. One may associate to S, a subspace of the Hilbert space $\mathscr{C}(S)$, which is defined as

$$\mathscr{C}(S) := \left\{ |\psi\rangle \in \left(\mathbb{C}^d\right)^{\otimes n} \mid g|\psi\rangle = |\psi\rangle \text{ for all } g \in S \right\}. \tag{20}$$

 $\dim \mathcal{C}(S) = d^k$ (see Theorem 1, in²⁶). It is easily seen that $\mathcal{C}(S)$ is the unique subspace *stabilised* by S, and is hence called the stabiliser code space corresponding to S. Since $\dim \mathcal{C}(S) = d^k$, $\mathcal{C}(S)$ encodes k qudits within itself, and thus $\mathcal{C}(S)$ is said to be an $[[n,k]]_d$ stabiliser code space, where the subscript denotes the dimension d of a single qudit.

² For even d, the group $\langle X,Z\rangle$ has some undesirable properties: while $X^d=Z^d=\mathbb{1}$, $(\omega^aXZ)^d=-\mathbb{1}$ for all $a=0,1,\cdots,d-1$, the order of ω^aXZ is 2d. It is cumbersome to keep track of which operator has order d and which has order 2d. By adding ζ to the group, we can obtain another group element ζXZ whose order is d. This is preferable to us because, while $\omega^aXZ \in \langle \zeta \mathbb{1}, XZ \rangle$ continues to have order d, we can always multiply this with ζ to get an order d element. The important point is that elements like X and ζXZ are put on the same footing. See⁵⁹ for a treatment of the even d case without the redundancy ζ phase factor.

Remark 1. When k = 0, $\dim \mathcal{C}(S) = 1$. Since $\mathcal{C}(S)$ is spanned by a single vector, we refer to that vector as a stabiliser state.

Remark 2. The case when k = n covers the scenario when we're encoding the whole Hilbert space into itself. We are not interested in this scenario.

Since $g_j = g(a_j)$ for some $a_j \in \mathbb{Z}_d^{2n}$, one may identify the subgroup S in terms of the a_j 's instead of the g_j 's 36 . This is done by arranging the a_j 's in a $2n \times (n-k)$ matrix, which is called the check matrix of S.

$$H = \left[a_1 \mid a_2 \mid \dots \mid a_{n-k} \right]. \tag{21}$$

The aforementioned two conditions on g_j 's may be phrased in terms of equivalent conditions on a_j 's.

$$g_i g_j = g_j g_i \Leftrightarrow a_j^T \Lambda a_i = 0. \tag{22}$$

$$|S| = d^{n-k} \Leftrightarrow \left| \text{span} \left\{ a_j \right\}_{j=1}^{n-k} \right| = d^{n-k}.$$
 (23)

Condition Eq. (22) follows from Eq. (14), and Condition Eq. (23) is equivalent to the fact that a_j 's are linearly independent and span an (n-k)-dimensional subspace.

The $2n \times (n-k)$ matrix H can be extended to a full $2n \times 2n$ symplectic matrix M. A partial construction of this full $2n \times 2n$ matrix can be found in Ref.³⁶. We refer the reader to Appendix \mathbb{C} for such a construction.

$$M = \left[E \left| L_X \right| H \left| L_Z \right| \right], \tag{24}$$

where E is a $2n \times (n-k)$ matrix, and L_X and L_Z are $2n \times k$ matrices. The columns of L_X and L_Z correspond to the logical X and the logical Z operators respectively, whereas the columns of E may be interpreted as the *correctable* Pauli errors. An explanation of this may be found in Section IV in conjunction with Section E of the Appendix. When k = 0, H is a $2n \times n$ matrix, and M takes the following form.

$$M = \left[E \mid H , \right] \tag{25}$$

where *E* is also a $2n \times n$ matrix.

When we alter the generators g_j of S as $g_j o \omega^{a_j} g_j$, for $a_j \in \mathbb{Z}_d$, then if $a_j \neq 0$ for some j, $\omega^{a_j} g_j$ are the generators of another stabiliser group S'. By replacing S with S' in Eq. (20), one can associate to S' a unique corresponding $[[n,k]]_d$ stabiliser code space $\mathscr{C}(S)$. $\mathscr{C}(S)$ and $\mathscr{C}(S')$ are orthogonal because the +1 eigenspaces of g_j and $\omega^{a_j} g_j$ are orthogonal when $a_j \neq 0$. There are d^{n-k} possible choices for a_j (including the choice $a_j = 0$ for all j), thus we get d^{n-k} orthogonal code spaces. Since each of these code spaces is of dimension d^k , the direct sum of all these code spaces is the full Hilbert space. Since a_j do not encode the overall phase factor of the Paulis, the entries in the check-matrix don't change

when the phase factors ω^{aj} are changed in the generators. So the check matrix H and its symplectic extension M for all the above codes spaces are the same. We will refer to a *code* as the aforementioned collection of code spaces without meaning to distinguish among them. Thus a code is associated to a check matrix without any ambiguity. We will refer to a *code space* $\mathscr{C}(S)$ as the subspace of the Hilbert space $\mathbb{C}^{d^{\otimes n}}$ associated to a unique stabiliser group S. This is as per Eq. (20).

Remark 3. Two codes spaces in the same code don't differ in any of their error correcting properties (such as distance, fault-tolerance threshold corresponding to some error model, the weight distribution of the generators (whether the code is LDPC or not, etc). Since one is interested these properties (among others), the emphasis is on finding good codes, and not on the many code-spaces with which these codes are realisable.

III. GROUP STRUCTURE OF $[[n,k]]_d$ STABILISER CODES

In this section, we will obtain a group theoretic structure underpinning all $[[n,k]]_d$ stabiliser codes. Let H_1 be the check matrix of some $[[n,k]]_d$ stabiliser code, and A be an $(n-k) \times (n-k)$ invertible matrix. Then H_1A is the check-matrix of the same stabiliser code. This is because the columns of H_1A are merely (invertible) linear combinations of the columns of H_1 , i.e., the generators corresponding to the column vectors of H_1 maybe recovered from the generators corresponding to the column vectors of H_2 . Define $H_2 = H_1A$. Let M_1 and M_2 be a $2n \times 2n$ symplectic matrix obtained by extending H_1 and H_2 respectively (see Section $\mathbb C$ of the Appendix). Since $M_1, M_2 \in \operatorname{Sp}(2n, \mathbb Z_d)$, there exists a matrix M such that

$$M_2 = M_1 M. (26)$$

We first assume that $k \ge 1$. Then it is clear that M needs to have the following form so that $H_2 = H_1 A$.

$$M = \begin{bmatrix} M_{11} & M_{12} & 0 & M_{14} \\ M_{21} & M_{22} & 0 & M_{24} \\ M_{31} & M_{32} & A & M_{34} \\ M_{41} & M_{42} & 0 & M_{44} \end{bmatrix}$$
 (27)

Here M_{11} , M_{31} , A are $(n-k)\times(n-k)$ matrices, M_{22} , M_{24} , M_{42} and M_{44} are $k\times k$ sized matrices, M_{12} , M_{14} , M_{32} , M_{34} are $(n-k)\times k$ and M_{21} , M_{41} are $k\times(n-k)$ matrices. Additionally M is necessarily a symplectic matrix, since $M=M_1^{-1}M_2$, and M_1 , M_2 are symplectic. Thus M satisfies the symplectic condition Eq. (19). Substituting M in Eq. (26) into Eq. (19), we get the conditions for M to be symplectic:

$$M_{11} = (A^T)^{-1}, (28)$$

$$M_{12} = M_{14} = 0_{(n-k)\times k},\tag{29}$$

$$\begin{bmatrix} M_{22} & M_{24} \\ M_{42} & M_{44} \end{bmatrix} \in \operatorname{Sp}(2k, \mathbb{Z}_d), \tag{30}$$

$$M_{31}A^T - AM_{31}^T = A\left(M_{41}^T M_{21} - M_{21}^T M_{41}\right)A^T, \tag{31}$$

$$M_{32} = A \left(M_{41}^T M_{22} - M_{21}^T M_{42} \right), \tag{32}$$

$$M_{34} = A \left(M_{41}^T M_{24} - M_{21}^T M_{44} \right), \tag{33}$$

where $\operatorname{Sp}(2k, \mathbb{Z}_d)$ denotes the group of $2k \times 2k$ symplectic matrices over \mathbb{Z}_d . Using Eq. (28) and Eq. (29), M takes the following form

$$M = \begin{bmatrix} (A^T)^{-1} & 0 & 0 & 0 \\ M_{21} & M_{22} & 0 & M_{24} \\ M_{31} & M_{32} & A & M_{34} \\ M_{41} & M_{42} & 0 & M_{44} \end{bmatrix}.$$
 (34)

The derivation of Eq. (30)-(33) is not essential for this work, and is hence shifted to the Appendix D.

Note that when k = 0, M in Eq. (34) takes the form

$$M = \begin{bmatrix} \left(A^T \right)^{-1} & 0 \\ M_{31} & A \end{bmatrix}. \tag{35}$$

Theorem 1. The set of all $2n \times 2n$ symplectic matrices of the form given in Eq. (34) form a subgroup of $Sp(2n, \mathbb{Z}_d)$.

Proof. We use the following subgroup test: if G is a group, H a subset of G, then H is a subgroup if for all $h, g \in H$, $h^{-1}g \in H$ too.

First note that the inverse of a symplectic matrix M is given by $\Lambda^T M^T \Lambda$, since $\Lambda^T M^T \Lambda M = \Lambda^T \Lambda = I$. This gives us

$$M^{-1} = \begin{bmatrix} A^T & 0 & 0 & 0 \\ M_{34}^T & M_{44}^T & 0 & -M_{24}^T \\ -M_{31}^T & -M_{41}^T & A^{-1} & M_{21}^T \\ -M_{32}^T & -M_{42}^T & 0 & M_{22}^T \end{bmatrix}.$$
(36)

Thus we see that M^{-1} has the form given by Eq. (34). Next, consider the composition of two symplectic matrices of the form Eq. (34).

$$\begin{bmatrix} (A^{T})^{-1} & 0 & 0 & 0 \\ M_{21} & M_{22} & 0 & M_{24} \\ M_{31} & M_{32} & A & M_{34} \\ M_{41} & M_{42} & 0 & M_{44} \end{bmatrix} \begin{bmatrix} (B^{T})^{-1} & 0 & 0 & 0 \\ N_{21} & N_{22} & 0 & N_{24} \\ N_{31} & N_{32} & B & N_{34} \\ N_{41} & N_{42} & 0 & M_{44} \end{bmatrix}$$

$$= \begin{bmatrix} ((AB)^{T})^{-1} & 0 & 0 & 0 \\ L_{21} & L_{22} & 0 & L_{24} \\ L_{31} & L_{32} & AB & L_{34} \\ L_{41} & L_{42} & 0 & L_{44} \end{bmatrix}.$$
(37)

Note that the matrix on the RHS of Eq. (37) is symplectic, since it is a product of two symplectic matrices. Also it is of the form given in Eq. (34).

The subgroup given in Theorem 1 will be denoted by $T(2n,k,\mathbb{Z}_d)$. It encapsulates the degree of freedom with which one may extend the check matrix of the $[[n,k]]_d$ trivial code to a symplectic matrix. The trivial code is the code corresponding to the stabiliser group $S = \langle Z_1, Z_2, \cdots, Z_{n-k} \rangle$. At the same time, right multiplying the symplectic matrix of any $[[n,k]]_d$ stabiliser code by an element of $T(2n,k,\mathbb{Z}_d)$ yields another another symplectic matrix of the same stabiliser code. We saw this in Eq. (26). In the next section we show that $T(2n,k,\mathbb{Z}_d)$ decomposes as a product of three subgroups, and highlight the roles played by these three subgroups in quantum error correction for stabiliser codes.

IV. DECOMPOSITION OF $T(2n,k,\mathbb{Z}_d)$

The matrix M given in Eq. (34) can be decomposed as a product of three matrices.

$$M = M_T M_E M_S M_L, (38)$$

where

$$M_T := \begin{bmatrix} I_k & 0 & 0 & 0 \\ 0 & I_{n-k} & 0 & 0 \\ K_S & 0 & I_k & 0 \\ 0 & 0 & 0 & I_{n-k} \end{bmatrix}, \tag{39}$$

$$M_E := \begin{bmatrix} I_k & 0 & 0 & 0 \\ N & I_{n-k} & 0 & 0 \\ K_A & L^T & I_k & -N^T \\ L & 0 & 0 & I_{n-k} \end{bmatrix}, \tag{40}$$

$$M_{L} := \begin{bmatrix} I_{k} & 0 & 0 & 0 \\ 0 & M_{22} & 0 & M_{24} \\ 0 & 0 & I_{k} & 0 \\ 0 & M_{42} & 0 & M_{44} \end{bmatrix}$$
(41)

and $M_S := \begin{bmatrix} (A^T)^{-1} & 0 & 0 & 0 \\ 0 & I_{n-k} & 0 & 0 \\ 0 & 0 & A & 0 \\ 0 & 0 & 0 & I_{n-k} \end{bmatrix},$ (42)

where

$$N = M_{21}A^{T},$$

$$K_{S} = M_{31}A^{T} + AM_{31}^{T},$$

$$K_{A} = M_{31}A^{T} - AM_{31}^{T},$$
and $L = M_{41}A^{T}.$ (43)

Furthermore, we have the following.

Corollary 3. For any $M \in T(2n, k, \mathbb{Z}_d)$, the decomposition in Eq. (38) is unique, i.e., there are unique matrices M_S , M_L , M_E and M_T such that Eq. (38) holds.

Proof. Since M_S and M_L are constructed from some of the blocks of M, there is a unique way of obtaining them. Having obtained M_S and M_L , $M_TM_E = M(M_LM_S)^{-1}$. Since N_S and N_S are matrix blocks within N_S and N_S and N_S are itirely determined by N_S and N_S and N_S a unique N_S is obtained from N_S and N_S which also gives us a unique N_S . Hence proved.

We next note that four subsets of matrices of M_S , M_L , M_E and M_T form four distinct subgroups, which don't share any overlaps with each other.

a. The $GL(n-k,\mathbb{Z}_d)$ subgroup: The set of matrices of the form M_S , with A invertible, is readily seen to be a subgroup of $T(2n,k,\mathbb{Z}_d)$ (by setting all matrix elements in M_{21} , M_{24} , M_{31} , M_{32} , M_{34} , M_{41} and M_{42} as 0, and setting $M_{22} = M_{44} = I_{n-k}$). Moreover, this subgroup is isomorphic

to $\operatorname{GL}(n-k,\mathbb{Z}_d)$, the group of $(n-k)\times (n-k)$ general linear matrices over \mathbb{Z}_d , and hence we refer to it as the $\operatorname{GL}(n-k,\mathbb{Z}_d)$ subgroup.

b. The $\operatorname{Sp}(2k,\mathbb{Z}_d)$ subgroup: All matrices of the form M_L which satisfy Eq. (30) form a subgroup of $\operatorname{T}(2n,k,\mathbb{Z}_d)$, and this subgroup is isomorphic to $\operatorname{Sp}(2k,\mathbb{Z}_d)$. Thus we call it the $\operatorname{Sp}(n-k,\mathbb{Z}_d)$ subgroup.

Note that the $GL(n - k, \mathbb{Z}_d)$ subgroup and the $Sp(k, \mathbb{Z}_d)$ subgroup commute with each other.

c. The symmetric and antisymmetric abelian subgroups $B_S(n,k,d)$ and $B_A(n,k,d)$: Matrices of the form M_T and M_E , which satisfy Eq. (31) (see Eq. (43)), are also subgroups of $T(2n,k,\mathbb{Z}_d)$. For M_T to satisfy Eq. (31), K_S has to be symmetric but is otherwise unconstrained, while for M_E to satisfy Eq. (31), K_A has to be anti-symmetric and moreover satisfies the equation $K_A = N^T L - L^T N$ (see Eq. (43)). It is easily verified that both are abelian subgroups, and also commute with each other. We refer to them as the symmetric and antisymmetric abelian subgroups $B_S(n,k,d)$, $B_A(n,k,d)$,

$$B_S(n,k,d) = \{ \text{ all } M_T \in T(2n,k,\mathbb{Z}_d) \}.$$
 (44)

$$B_A(n,k,d) = \{ \text{ all } M_E \in T(2n,k,\mathbb{Z}_d) \}.$$
 (45)

Remark 4. When k = 0, the $B_A(n,k,d)$ and $\operatorname{Sp}(2k,\mathbb{Z}_d)$ subgroups shrink to the trivial group which contains only the identity. The decomposition of $\operatorname{T}(2n,k,\mathbb{Z}_d)$ then takes the following form.

$$M = M_T M_S. (46)$$

The uniqueness of the decomposition (i.e., Corollary 3) holds for this case well.

In the Appendix Section E, we elaborate on the significance of these subgroups in quantum error correction. While the roles played by $GL(n-k,\mathbb{Z}_d)$ and $Sp(2k,\mathbb{Z}_{\mathbb{Z}_d})$ is easily surmised to those familiar with QEC with stabilizer codes, there has been less spotlight on the roles played by $B_A(n,k,d)$ and $B_S(n,k,d)$, i.e., the fact that the choice of selecting correctable errors for the code is encapsulated by the *group actions* of $B_S(n,k,d)$ and $B_A(n,k,d)$.

V. COUNTING $[[n,k]]_d$ STABILISER CODES

One may employ Lagrange's theorem in theory of finite groups to compute the total number of $[[n,k]]_d$ stabiliser codes, which we denote as $\mathscr{C}[[n,k]]_d$.

Lemma 1. The number of $[[n,k]]_d$ stabilizer codes is

$$\mathscr{C}[[n,k]]_d = |\operatorname{Sp}(2n,\mathbb{Z}_d)|/|\operatorname{T}(2n,k,\mathbb{Z}_d)|, \qquad (47)$$

where $|\operatorname{Sp}(2n,\mathbb{Z}_d)|$ is the order of $\operatorname{Sp}(2n,\mathbb{Z}_d)$ and $|\operatorname{T}(2n,k,\mathbb{Z}_d)|$ is the order of $\operatorname{T}(2n,k,\mathbb{Z}_d)$.

Remark 5. For the case when d is a prime, i.e., d = p, the number of $[[n,k]]_d$ stabiliser codes was explicitly computed in $Ref.^{22}$ (see Theorem 20 and Corollary 21 therein). The results we obtain below agree with the results in $Ref.^{22}$ in this case. There is a discrepancy in the language and notation employed: an $[[n,k]]_d$ stabiliser code in our work corresponds to an m-dimensional isotropic subspace in Theorem 20 in $Ref.^{22}$, each $[[n,k]]_d$ code space in our work is a counted as a distinct code in Corollary 21 in $Ref.^{22}$, and finally, we use the notation $\mathscr{C}[[n,k]]_d$ to count the number of $[[n,k]]_d$ stabiliser codes, whereas it is used to count the total number of code spaces in $Ref.^{22}$.

Proof. For a given $[[n,k]]_d$ stabiliser code, let's choose a representative code space $\mathscr{C}(S)$ with the stabiliser group S= $\langle g_j \rangle_{j=1}^{n-k}$. From S one can construct a check matrix H, which may be extended to a symplectic matrix M. In Sec. III we noted that this construction of M from S has a redundancy, and that the degree of freedom within the redundancy is captured by the right action on M by the subgroup $T(2n,k,\mathbb{Z}_d)$. In other words, each symplectic matrix in any coset of $\operatorname{Sp}(2n,\mathbb{Z}_d)/\operatorname{T}(2n,k,\mathbb{Z}_d)$ is a possible symplectic construction for the same $[[n,k]]_d$ stabiliser code. This tells us that the number of $[[n,k]]_d$ codes is lesser than or equal to the number of left cosets, i.e., $|\operatorname{Sp}(2n,\mathbb{Z}_d)/\operatorname{T}(2n,k,\mathbb{Z}_d)|$. Conversely, let us start from some left coset in $Sp(2n, \mathbb{Z}_d)/T(2n, k, \mathbb{Z}_d)$, i.e., $M \times T(2n, k, \mathbb{Z}_d)$, where we designate M as the coset representative. One can associate to this coset any one of d^{n-k} mutually orthogonal $[[n,k]]_d$ stabiliser codes in the following way: extract from M a $2n \times (n-k)$ submatrix by extracting the columns with numbers n+1 to 2n-k from the left (see Eq. (24)). Call this submatrix H. Since $M \in \text{Sp}(2n, \mathbb{Z}_d)$ and satisfies $M^T \Lambda M = \Lambda$, H satisfies $H^T \Lambda H = 0$. Let the j-th column in H be the homomorphic image of some Pauli $\omega^{a_j}g_j$ (see Eq. (11)), where g_j 's are such that each has a +1 eigenspace, and the a_j 's are in \mathbb{Z}_d and arbitrary. Generate the group $S = \langle \omega^{a_j} g_j \rangle$. Note that the construction branches out into d^{n-k} different choices of S, depending on the values of the a_i 's. That $H^T \Lambda H = 0$ is equivalent to the fact that the g_i 's commute between themselves. Also, the columns of H are linearly independent since M is an invertible matrix. This ensures that any subset of the $\omega^{a_j}g_j$'s will generate a strictly smaller subgroup of S. Construct the coding space $\mathscr{C}(S)$ from S, and note that $\dim \mathcal{C}(S) = 2^k$ (see Eq. (20) and the description below it). Thus $\mathscr{C}(S)$ is a code space corresponding to an $[[n,k]]_d$ stabiliser code. The d^{n-k} distinct choices of S and $\mathscr{C}(S)$ correspond to the d^{n-k} different choices within the same stabiliser code. This tells us that the number of cosets is lesser than or equal to the number of $[[n,k]]_d$ stabiliser codes.

Note that when the code to coset construction is reversed, one retrieves the original code which one started with. This is true for all codes. This proves the theorem.

Using the decomposition of $T(2n, k, \mathbb{Z}_d)$ from Sec. IV we obtain the following theorem.

Lemma 2. Let d be a prime power, i.e., $d = p^m$, where p is a prime number and m a positive integer. The total number of

(49)

 $[[n,k]]_{p^m}$ stabiliser codes, $\mathscr{C}[[n,k]]_{p^m}$ is

$$[n,k]_{p^m}$$
 stabiliser codes, $\mathscr{C}[[n,k]]_{p^m}$ is $[n,k]_{p^m}$ where the Gaussian coefficient $[n,k]_{p^m}$ is defined as
$$[n,k]_{p^m} \coloneqq \prod_{i=0}^{n-k-1} \frac{p^{n-j}-1}{p^{n-k-j}-1}.$$

$$\mathscr{C}[[n,k]]_{p^m} = \left(p^{m-1}\right)^{\frac{(n-k)(n+3k+1)}{2}} \begin{bmatrix} n \\ n-k \end{bmatrix}_p \prod_{j=0}^{n-k-1} \left((p)^{n-j}+1\right)$$
(48)

Proof. Using Lemma 1, the decomposition in Sec. IV and Lemma 3,

$$\mathscr{C}[[n,k]]_{p^{m}} = \begin{cases} \frac{\left|\operatorname{Sp}(2n,\mathbb{Z}_{p^{m}})\right|}{\left|B_{S}\left(n,k,p^{m}\right)\right| \left|B_{A}\left(n,k,p^{m}\right)\right| \left|\operatorname{GL}\left(n-k,\mathbb{Z}_{p^{m}}\right)\right| \left|\operatorname{Sp}(2k,\mathbb{Z}_{p^{m}})\right|}, \text{ when } k \geq 1 \text{ and} \\ \frac{\left|\operatorname{Sp}(2n,\mathbb{Z}_{d})\right|}{\left|B_{S}\left(n,0,p^{m}\right)\right| \left|\operatorname{GL}\left(n,\mathbb{Z}_{p^{m}}\right)\right|}, \text{ when } k = 0. \end{cases}$$

$$(50)$$

It is straightforward to compute that $|B_A(n,k,p^m)| =$ $(p^m)^{2k(n-k)}$, since in M_E from Eq. (40), N and L are left completely unconstrained by Eq. (43), whereas simultaneously K_A is completely determined by N and L. Also note, $|B_S(n,k,p^m)| = p^{m\frac{1}{2}(n-k)(n-k+1)}$, since in M_T from Eq. (39), the only constraint which Eq. (43) imposes on K_S is that it be symmetric. $|GL(n-k,\mathbb{Z}_{p^m})|$ is computed in Corollary 2.8 in Ref. 31 (see also Ref. 39 for a more accessible arguments):

$$\left| \operatorname{GL}(n-k, \mathbb{Z}_{p^m}) \right| = p^{(m-1)(n-k)^2} \prod_{j=0}^{n-k-1} \left(p^{n-k} - p^j \right).$$
 (51)

 $|\operatorname{Sp}(2n,\mathbb{Z}_{p^m})|$ and $|\operatorname{Sp}(2k,\mathbb{Z}_{p_i^{m_i}})|$ (needed for the case $k \geq 1$) are explicitly computed in the Appendix Sec. E1 (which is based on Ref.³⁸), which gives us

$$\left| \operatorname{Sp}(2n, \mathbb{Z}_{p^m}) \right| = p^{(2m-1)n^2 + (m-1)n} \prod_{i=1}^n \left(p^{2i} - 1 \right).$$
 (52)

Then putting everything together in Eq. (50) gives us Eq. (48)

Remark 6. In a seminal paper on Wigner distributions on finite dimensional phase-space Ref.²², Gross computed the number of $[[n,k]]_d$ codes when $d=p^m$, but the phase-space is identified with $\mathbb{F}_{p^m}^2$, not $\mathbb{Z}_{p^m}^2$. That the Pauli groups for both phase spaces are different is observed from the following fact. Set n = 1. Then $\mathscr{P}^{(p^m)}$ is homomorphic to the additive abelian group of $\mathbb{Z}_{p^m}^2$ (i.e., when only looking at $\mathbb{Z}_{p^m}^2$ as an abelian group), whereas $\mathscr{P}^{(\mathbb{F}_{p^m})}$ is homomorphic to the additive abelian group of $\mathbb{F}_{p^m}^2$. Note that the abelian group of $\mathbb{Z}_{p^m}^2$ decomposes as the product of two cyclic groups: $(\mathbb{Z}_{p^m}^2,+)\simeq$

 $(\mathbb{Z}_{p^m},+) imes (\mathbb{Z}_{p^m},+)$, whereas that of $\mathbb{F}^2_{p^m}$ decomposes as a product of 2m cyclic groups, i.e., $\left(\mathbb{F}_{p^m}^2,+\right) \simeq \left(\mathbb{Z}_p,+\right)^{\times 2m}$. Furthermore these decompositions are unique by the structure theorem of finite abelian groups. When m = 1, i.e., d = p, the formula in Theorem 20 in Ref.²² matches that with Eq. (48), but for larger m, the formula differs on account of the fact that while \mathbb{F}_{p^m} is a field, \mathbb{Z}_{p^m} isn't. For completeness, in Appendix G we explain why this renders the proof for Theorem 20 in Ref.²² inapplicable to our case.

Theorem 2. When d is an arbitrary positive integer with prime factorisation $d = \prod_{i=1}^{r} p_i^{m_i}$, where p_i are distinct primes, m_i are positive integers and r, which is the number of distinct prime factors, is also a positive integer. Then total number of $[[n,k]]_d$ stabiliser codes is

$$\mathscr{C}[[n,k]]_d = \prod_{i=1}^r \mathscr{C}[[n,k]]_{p_i^{m_i}}.$$
 (53)

Thus we get

$$\mathscr{C}[[n,k]]_d = d^{\frac{(n-k)(n+3k+1)}{2}} \prod_{i=1}^r \zeta_i,$$
 (54)

where

$$\zeta_i := \prod_{j=0}^{n-k-1} \left(\frac{1 - p_i^{-2(n-j)}}{1 - p_i^{-(n-k-j)}} \right).$$
(55)

Proof. We invoke the Chinese remainder theorem³², which tells us that

$$\mathbb{Z}_d \simeq \mathbb{Z}_{p_1}^{m_1} \times \mathbb{Z}_{p_2}^{m_2} \times \cdots \mathbb{Z}_{p_r}^{m_r}, \tag{56}$$

is a ring isomorphism. A simple corollary of this ring isomorphism is the following two group isomorphisms.

$$\operatorname{Sp}(2n, \mathbb{Z}_d) \simeq \operatorname{Sp}(2n, \mathbb{Z}_{p_1^{m_1}}) \times \operatorname{Sp}(2n, \mathbb{Z}_{p_2^{m_2}}) \times \cdots \times \operatorname{Sp}(2n, \mathbb{Z}_{p_r^{m_r}})$$

$$\operatorname{T}(2n, k, \mathbb{Z}_d) \simeq \operatorname{T}(2n, k, \mathbb{Z}_{p_1^{m_1}}) \times \operatorname{T}(2n, k, \mathbb{Z}_{p_2^{m_2}}) \times \cdots \times \operatorname{T}(2n, k, \mathbb{Z}_{p_r^{m_r}}), \tag{57}$$

where the \times symbol on the RHS denotes the direct product of groups. Invoking Eq. (57) into Lemma 1 thus proves Eq. (53). For Eq. (54), we first simplify the RHS of Eq. (48) using the following.

$$\begin{bmatrix} n \\ n-k \end{bmatrix}_{p_i} \prod_{j=0}^{n-k-1} \left((p_i)^{n-j} + 1 \right)
= p_i^{\frac{(n-k)(n+3k+1)}{2}} \prod_{i=0}^{n-k-1} \left(\frac{1-p_i^{-2(n-j)}}{1-p_i^{-(n-k-j)}} \right).$$
(58)

Using Eq. (58) in Eq. (48) gives us

$$\mathscr{C}[[n,k]]_{p_i^{m_i}} = p_i^{m_i}^{\frac{(n-k)(n+3k+1)}{2}} \zeta_i, \tag{59}$$

which is then invoked into Eq. (53).

It is often important to simply get an order of magnitude of $\mathscr{C}[[n,k]]_d$. Towards that result, we obtain the following corollary as a result of Theorem 2.

Corollary 4. Let c = 2.17. Then number of $[[n,k]]_d$ stabiliser codes scales as

$$d^{\frac{(n-k)(n+3k+1)}{2}} \le \mathscr{C}[[n,k]]_d < d^{\frac{(n-k)(n+3k+1)}{2}+c}, \tag{60}$$

the number of $[[n,k]]_d$ stabiliser code spaces scale as

$$d^{\frac{(n-k)(n+3k+3)}{2}} < \mathcal{C}_{\text{space}}[[n,k]]_d < d^{\frac{(n-k)(n+3k+3)}{2}+c}, \quad (61)$$

and thus the number of stabilizer states scale as

$$d^{\frac{n(n+3)}{2}} \le \mathcal{C}_{\text{state}}[[n]]_d < d^{\frac{n(n+3)}{2}+c}. \tag{62}$$

Proof. It is not difficult to upper bound ζ_i by a constant which is independent of p_i , n and k. For instance, in Appendix Sec. F we show that

$$\zeta_i < e^{1.57}, \forall n, k, \text{ and primes } p.$$
 (63)

Thus we now use Eq. (58) and Eq. (63) in Eq. (48)

$$\mathscr{C}[[n,k]]_{p_i^{m_i}} < e^{c_1} \left(p_i^{m_i} \right)^{\frac{(n-k)(n+3k+1)}{2}}, \tag{64}$$

where $c_1 = 1.57$. Let the number of distinct prime factors of d be denoted by r(d). Then Eq. (53) and (64) tell us

$$\mathscr{C}[[n,k]]_d < e^{c_1 r(d)} d^{\frac{(n-k)(n+3k+1)}{2}}.$$
 (65)

It is known that r(d) may be upper bounded as follows (see Theorem 11, p. 369 in Ref. 42):

$$r(d) \le c_2 \frac{\log d}{\log \log d}$$
, for $d \ge 3$. (66)

where $c_2 = 1.38$. Thus we get that $e^{c_1 r(d)} \le d^c$, where $c = c_1 c_2 \approx 2.17$. Thus we get

$$\mathscr{C}[[n,k]]_d < d^{\frac{(n-k)(n+3k+1)}{2}+c}.$$
(67)

Note that for d=2, Eq. (64) is already satisfied, and hence, d=2 also satisfies Eq. (67). The number of $[[n,k]]_d$ stabilizer code spaces is $d^{n-k} \mathcal{C}[[n,k]]_d$, and the number of stabilizer states is obtained by setting k=0 in the number of stabilizer code spaces.

VI. CONCLUSIONS

In this work we count the number of $[[n,k]]_d$ stabilizer codes, $\mathscr{C}[[n,k]]_d$ for arbitrary d-level systems, where the configuration space of such systems is \mathbb{Z}_d . Since \mathbb{Z}_d is not a field when d is non-prime, the method we used for this has to differ from earlier works in Ref.⁶ for qubits and Ref.²² for prime-dimensional qudits and Galois-qudits, which relied on d being prime. Our method is broadly broken up into two parts: (i) proving a bijection between distinct $[[n,k]]_d$ QECC and cosets of $Sp(2n,\mathbb{Z}_d)/T(2n,k,\mathbb{Z}_d)$, where $\mathrm{T}(2n,k,\mathbb{Z}_d)$ is a subgroup, which corresponds to the $[[n,k]]_d$ trivial code, with stabilizer group $S = \langle Z_j \rangle_{j=1}^{n-k}$, and (ii) computing $|\operatorname{Sp}(2n,\mathbb{Z}_d)|/|\operatorname{T}(2n,k,\mathbb{Z}_d)|$, for which we make use of the Chinese remainder theorem. We find that the number of $[[n,k]]_d$ stabilizer codes $\mathscr{C}[[n,k]]_d$, the number of $[[n,k]]_d$ stabilizer code subspaces $\mathscr{C}_{\text{space}}[[n,k]]_d$ and the number of stabilizer states $\mathscr{C}_{\text{state}}[[n]]_d$ scale agnostic of whether d is prime or non-prime. This is surprising since the prime or non-prime nature of d played an important role in our computation. In Section I we listed salient topics where $\mathscr{C}_{\text{state}}[[n]]_d$ plays the role as an important quantifier: the resource theory of magic, the classical simulation of stabilizer-only circuits, projective-designs of stabilizer states, a de Finetti theorem customized for stabilizer operations, the study of quantum contextuality for small systems, the study of Wigner functions and the optimal verification of stabilizer states. Since $\mathscr{C}_{\text{state}}[[n]]_d$ was so far known only for prime d (and Galois-qudits), the results were limited to such cases. For resource theory of magic and the optimal verification of stabilizer states, our work allows one to make statements for arbitrary d. For the remaining topics, we believe that our computation of $\mathscr{C}_{\text{state}}[[n]]_d$ will prove useful for generalising the corresponding results to arbitrary d. Thus our work also contributes towards the important goal of placing results for arbitrary d gudit systems on the same pedestal as for prime qudit systems or Galois qudit systems.

- ¹Mixed-state entanglement and quantum error correction, Bennett C. H., Di-Vincenzo D. P., Smolin J. A., and Wootters W. K.; Phys. Rev. A 54, 3824 (1996)
- ²Quantum data hiding, D. P. DiVincenzo, D. W. Leung and B. M. Terhal, IEEE Transactions on Information Theory, vol. 48, no. 3, pp. 580-598, March 2002
- ³Randomized benchmarking of quantum gates, Knill E., Leibfried D., R. Reichle, J. Britton, R. B. Blakestad, J. D. Jost, C. Langer, R. Ozeri, S. Seidelin, and D. J. Wineland Phys. Rev. A 77, 012307 (2008)
- ⁴Scalable and Robust Randomized Benchmarking of Quantum Processes, Magesan E., Gambetta J. M., Emerson J., Phys. Rev. Lett. 106, 180504 (2011)
- ⁵The Heisenberg Representation of Quantum Computers, Gottesmann D., preprint at: arXiv:quant-ph/9807006.
- Journal reference: Group22: Proceedings of the XXII International Colloquium on Group Theoretical Methods in Physics, eds. S. P. Corney, R. Delbourgo, and P. D. Jarvis, pp. 32-43 (Cambridge, MA, International Press, 1999)
- ⁶Improved simulation of stabilizer circuits, Scott Aaronson and Daniel Gottesman, Phys. Rev. A 70, 052328 (2004)
- ⁷New techniques for bounding stabilizer rank, Benjamin Lovitz and Vincent Steffan", Quantum 6, 692 (2022).
- ⁸Quantum systems with finite Hilbert space, A Vourdas 2004 Rep. Prog. Phys. 67 267
- ⁹Kitaev's \mathbb{Z}_d -code threshold estimates, Duclos-Cianci G., and Poulin D., Phys. Rev. A 87, 062338 (2013)
- ¹⁰Fast decoders for qudit topological codes, Anwar H., Brown B. J., Campbell E. T., and Browne D. E., New J. Phys. 16 063038 (2014)
- ¹¹Magic-State Distillation in All Prime Dimensions Using Quantum Reed-Muller Codes, Campbell E.T, Anwar H., and Browne D. E., Phys. Rev. X 2, 041021 (2012)
- ¹²Enhanced Fault-Tolerant Quantum Computing in d-Level Systems, Campbell E. T., Phys. Rev. Lett. 113, 230501 (2014)
- ¹³Towards Low Overhead Magic State Distillation, Krishna A., and Tillich J. P., Phys. Rev. Lett. 123, 070507 (2019)
- ¹⁴ A quantum compiler for qudits of prime dimension greater than 3, Heyfron L. E. and Campbell E. T., arxiv:1902.05634 (2019)
- ¹⁵Quantum computation with realistic magic-state factories, O'Gorman J., and Campbell E. T., Phys. Rev. A 95, 032338 (2017)
- 16"Galois-qudit code", The Error Correction Zoo (V. V. Albert & P. Faist, eds.), 2022.
- 17"Modular-qudit stabilizer code", The Error Correction Zoo (V. V. Albert & P. Faist, eds.), 2022.
- ¹⁸H. Weyl, "Theory of Groups and Quantum Mechanics", Dutton, New York, (1932)
- ¹⁹Unitary Operator Basis, J. Schwinger, PNAS 46, 570 (1960)
- ²⁰Symmetric informationally complete–positive operator valued measures and the extended Clifford group, D. M. Appleby, J. Math. Phys. 46, 052107 (2005)
- ²¹ Stabilizer states and Clifford operations for systems of arbitrary dimensions and modular arithmetic, E. Hostens, J. Dehaene, and B. De Moor, Phys. Rev. A 71, 042315 (2005)
- ²²Hudson's theorem for finite-dimensional quantum systems, D. Gross , J. Math. Phys. 47, 122107 (2006)
- ²³Qudit surface codes and gauge theory with finite cyclic groups, S. S. Bullock and G. K. Brennen, 2007 J. Phys. A: Math. Theor. 40 3481
- ²⁴Homological error correction: Classical and quantum codes, Bombin H., and Martin-Delgado M. A., J. Math. Phys. 48, 052105 (2007)
- ²⁵The monomial representations of the Clifford group, Appleby D. M., Bengtsson I., Brierley S., Grassl M., Gross D., and Larsson J., Quantum Information and Computation, Vol.12 No.5 & 6, 2012
- ²⁶Standard form of qudit stabilizer groups, Gheorghiu V., Phys. Lett. A, Vol 378, No. 5–6, P 505-509, (2014)
- ²⁷Geometry of Quantum States. An Introduction to Quantum Entanglement, Bengtsson I., and Życzkowski K., 2nd Edition Cambridge: Cambridge University Press.
- ²⁸ "Algebra", Artin, M., Algebra, second ed., Prentice Hall, Boston, MA, 2011. ISBN 0321998030, 9780321998033.
- See Chapter 14, "Linearly Algebra on a Ring".

- ²⁹D. Gottesman, Stabilizer Codes and Quantum Error Correction, Ph.D. thesis (CalTech).
- ³⁰M. Newman, "Integral Matrices", Academic Press 1972
- ³¹The General Linear Group over a Ring Han, J.-C., (2006). Bull. Korean Math. Soc. 43(3), 619–626
 See Corollary 2.8 and 2.9.
- ³²Algebra, Lang S., 3rd edition, Graduate Texts in Mathematics, Springer New York, NY See Theorem 2.1 and Corollary 2.2.
- ³³ A Classical Introduction to Modern Number Theory, Ireland K., and Rosen, M., Graduate Texts in Mathematics, 84 (2nd ed.), p.35. Springer (1990)
- 34. Introductory Lectures on Rings and Modules, Beachy J, London Mathematical Society Student Texts. Cambridge: Cambridge University Press (1999).
 - See Theorem 1.2.7 (The fundamental homomorphism theorem) on p.27.
- ³⁵Classical Groups and Geometric Algebra, Grove L. C., Graduate Studies in Mathematics) (1st ed.), American Mathematical Society, 2001. Theorem 3.12, p. 27.
- ³⁶Quantum Computation and Quantum Information: 10th Anniversary Edition, Nielsen, M., & Chuang, I., Cambridge: Cambridge University Press See chapter 10 "Quantum error correction".
- ³⁷Poulain D., Optimal and efficient decoding of concatenated quantum block codes, Poulain D., Phys. Rev. A 74, 052333 (2006)
- 38 Is there a formula for the size of Symplectic group defined over a ring Z/p^kZ^2 , S. Carnahan
- ³⁹Order of $GL(n, \mathbb{Z}/m\mathbb{Z})$, Vaidyanathan P.
- ⁴⁰Introduction to analytic number theory, Apostol, Tom M., Undergraduate Texts in Mathematics, New York-Heidelberg: Springer-Verlag, (1976)
- ⁴¹What is a *q*-series?, Baruah N. D., Berndt B. C., Cooper S., Huber T., and Schlosser M. J., Ramanujan Rediscovered: Proceedings of a Conference on Elliptic Functions, Partitions, and q-Series in memory of K. Venkatachaliengar: Bangalore, 1-5 June, 2009, eds., Ramanujan Mathematical Society, Mysore, 2010, pp. 31-51
- ⁴²Estimation de la fonction de Tchebychef Θ sur le k-ième nombre premier et grandes valeurs de la fonction ω(n) nombre de diviseurs premiers de n, Robin, G., Acta Arithmetica 42 367-389, (1983)
- ⁴³Modular arithmetic before C.F. Gauss: Systematizations and discussions on remainder problems in 18th-century Germany, Bullynck M., Historia Mathematica, Vol. 36, Issue 1, 2009.
- ⁴⁴Negative quasi-probability as a resource for quantum computation, Veitch V., Ferrie C., Gross D., and Emerson J., New J. Phys. 14 113011 (2012)
- ⁴⁵The resource theory of stabilizer quantum computation, Veitch V., Mousavian S. A. H., Gottesman D., and Emerson J., New J. Phys. 16 013009 (2014)
- ⁴⁶Application of a Resource Theory for Magic States to Fault-Tolerant Quantum Computing, Howard, M. and Campbell, E., Phys. Rev. Lett. 118, 090501 (2017)
- ⁴⁷Estimating Outcome Probabilities of Quantum Circuits Using Quasiprobabilities", Pashayan H., Wallman J. J., and Bartlett S. D., Phys. Rev. Lett. 115, 070501 (2015)
- ⁴⁸Robustness of Magic and Symmetries of the Stabiliser Polytope, Heinrich, M. and Gross, D. Quantum 3, 132 (2019).
- ⁴⁹ Detecting Magic States via Characteristic Functions, Dai, H., Fu, S. & Luo, S., Int J Theor Phys 61, 35 (2022).
- ⁵⁰Comparative Study of Sampling-Based Simulation Costs of Noisy Quantum Circuits, Shigeo Hakkaku and Keisuke Fujii, Phys. Rev. Applied 15, 064027 Published 10 June 2021
- ⁵¹Stim: a fast stabilizer circuit simulator, Gidney C., Quantum 5, 497 (2021).
- ⁵²Qubit stabilizer states are complex projective 3-designs Kueng R. and Gross D., Preprint available at arXiv:1510.02767
- ⁵³Schur-Weyl Duality for the Clifford Group with Applications: Property Testing, a Robust Hudson Theorem, and de Finetti Representations, Gross, D., Nezami, S. and Walter, M., Commun. Math. Phys. 385, 1325–1393 (2021).
- ⁵⁴One-and-a-Half Quantum de Finetti Theorems, Christandl M., König R., Mitchison G. & Renner R., Commun. Math. Phys. 273, 473–498 (2007).
- ⁵⁵Quantum Contextuality with Stabilizer States, Howard M., Brennan, E., Vala, J., Entropy 15, 2340-2362 (2013).
- ⁵⁶Classical codes in quantum state space, Howard M., J. Phys. A: Math. Theor. 48 495303 (2015)

Theor. 43 075302 (2010)

⁶⁰user0, Reply on math.stackechange to query about embedding an n-tuple of \mathbb{Z}_d into an invertible $n \times n$ matrix A.

Appendix A: A short summary of the resource theory of magic

An operationally significant resource theory of magic has been studied and developed over the past decade. In this theory stabilizer operations are considered a free resource. By stabilizer operations we mean the following. (i) Initiating the starting state of a quantum computation as an eigenstate of some multi-qudit Pauli operator. Often this is the state $|0,0,\cdots,0\rangle$, which spans the $[[n,0]]_d$ trivial code for the *n*-qudit system with stabilizer group $S = \langle Z_j \rangle_{j=1}^n$. (ii) Performing Clifford operations during the computation. (iii) The only observables which are subjected to quantum measurements are multiqudit Pauli operators. It is known that stabilizer operations are not universal⁶, and universality demands the inclusion of some non-stabilizer operations, such as the $\pi/8$ gate via magic state injection. From the perspective of fault-tolerant quantum computing, performing non-stabilizer operations at the *logical* level of the quantum code, is a high-cost incurring operation (see for e.g. ¹⁵), which justifies the differentiation between the 'free' resources of stabilizer operations (the aforementioned three operations) and the costly resources of non-stabilizer operations (magic state injection).

Appendix B: Proof of Corollary 2

 \mathbb{Z}_d^n is a free- \mathbb{Z}_d module. This means that one may simply view \mathbb{Z}_d^n as an abelian group with vector addition as the corresponding group operation. Furthermore, \mathbb{Z}_d^n is then simply seen to be a direct product group of \mathbb{Z}_d with itself n times.

1. Any set of n LI vectors is a basis for \mathbb{Z}_d^n

By *basis* for \mathbb{Z}_d^n , we mean a *minimal generating set* for \mathbb{Z}_d^n as a group. Any set of n LI vectors is a minimal generating set since there are precisely d^n distinct linear combinations of these n vectors, and the LI property guarantees that each distinct linear combination results in a distinct vector in \mathbb{Z}_d^n . Since there are d^n vectors in \mathbb{Z}_d^n , each vector must be one of the d^n possible linear combinations.

2. If a_i 's are LI, then their linear span generates a subspace of dimension m

By the linear span of the a_j 's, we mean the subgroup of \mathbb{Z}_d^n , which the a_j 's generate. Since the a_j 's are LI and since they are m in number, they generate a subgroup of size d^m . However, not all subspaces of size d^m are isomorphic. It remains to justify the concept of an m dimensional subspace. We use the term m-dimensional subspace to mean a subgroup of \mathbb{Z}_d^n , which is isomorphic to \mathbb{Z}_d^m . One may easily construct such an isomorphism from span $\{a_j\}_{j=1}^m$ to \mathbb{Z}_d^m as follows: define T: span $\{a_j\}_{j=1}^m \to \mathbb{Z}_d^m$ as follows:

$$T(x_1a_1 + x_2a_2 + \dots + x_ma_m) = (x_1, x_2, \dots, x_m), \ \forall \ x_i \in \mathbb{Z}_d.$$
(B1)

T is readily seen to be an isomorphism (in the group theoretic sense).

3. For any m < n, any set of m LI vectors may be extended to form a basis for \mathbb{Z}_{d}^{n} .

We prove this by finding an invertible matrix A whose first m columns are given by the LI vectors a_1, a_2, \dots, a_m . Then we may simply add the remaining n-m columns to obtain a set of n linearly independent columns. We prove this inductively.

When m = 1. We adapt the arguments given in Ref. 603 to our purpose. To say the $a_1 = a$ is linearly independent

⁵⁷Optimal verification of stabilizer states, Dangniam N., Han Y.-G., Zhu H., Phys. Rev. Research 2, 043323 (2020)

⁵⁸The axiomatic and the operational approaches to resource theories of magic do not coincide, Heimendahl A., Heinrich M., and Gross D., arXiv:2011.11651

⁵⁹ Wigner distributions for finite-state systems without redundant phase-point operators, Chaturvedi S., Mukunda N., and Simon R., J. Phys. A: Math.

³ The proof in Ref.⁶⁰ is itself taken from Ref.³⁰ (Corollary II.I). We can't rely on the proofs in Ref.^{30,60} directly, since these results are valid only for

means that there is no non-zero $x \in \mathbb{Z}_d$ such that xa = 0. In other words, a is of order d. This implies that if $a = (a_1, a_2, \dots, a_n)^T$, then as integers in \mathbb{Z} , g.c.d. $\{a_1, a_2, \dots, a_n, d\} = 1 \in \mathbb{Z}$. We can thus apply Bézout's identity (see Ref.⁴³), in the same way that it is applied in Ref.⁶⁰. For completeness, we give the whole proof here.

Proceeding inductively, suppose that n=2 first. Bézout's identity tells us that there exist b_1 , b_2 and d' in $\mathbb Z$ such that $a_1b_2-a_2b_1+d'd=1$ in $\mathbb Z$. Thus we also get that $(a_1b_2-a_2b_1)\mod d=1\in\mathbb Z_d$. With $A=\begin{pmatrix}a_1&b_1\\a_2&b_2\end{pmatrix}\mod d$, the result is proved for n=2. Next, suppose that the result is true for n-1. We need to then prove the result for n. As before $g.c.d\{a_1,a_2,\cdots,a_n,d\}=1\in\mathbb Z$. There exists some integer g which is a common factor of a_1,a_2,\cdots and a_{n-1} such that if $b_i=a_i/g$, then $g.c.d\{b_1,b_2,\cdots,b_{n-1},d\}=1$. It isn't necessary that g equals $g':=g.c.d\{a_1,a_2,\cdots,a_{n-1},d\}$, because $g.c.d\{a_1/g',a_2/g',\cdots,a_{n-1}/g',d\}$ need not be 1. Writing d=g'h', we see that a_i/g' have no common factors with h' (otherwise g' would have been larger). Thus the only factors which $a_1/g',a_2/g',\cdots,a_{n-1}/g'$ have in common with g'0 will occur in g'1. Writing $g.c.d\{a_1/g',a_2/g',\cdots,a_{n-1}/g',g'\}=\tilde{g}'$ 2, and defining $g''=\tilde{g}'g'$ 3, we examine whether $a_1/g'',a_2/g'',\cdots,a_{n-1}/g''$ 3 and g'1 have any common factors. If not, we may choose g'2 is necessarily a factor of g'3 itself. Since the hypothesis is assumed true for g'1, corresponding to g'2, where g'3 is necessarily a factor of g'4 itself. Since the hypothesis is assumed true for g'4, corresponding to g'5. Construct an g'6, where g'6 is necessarily and the same fashion as in Ref. g'6.

$$A^{T} = \begin{pmatrix} gb_{1} & gb_{2} \cdots & gb_{n-1} & a_{n1} \\ b_{21} & b_{22} \cdots & b_{2,n-1} & 0 \\ \vdots & \vdots \ddots & \vdots & \vdots \\ b_{n-1,1} & b_{n-1,2} \cdots & b_{n-1,n-1} & 0 \\ rb_{1} & rb_{2} \cdots & rb_{n-1} & s \end{pmatrix},$$
(B2)

which gives us $\det A = (sg - a_{n1}r) \det B$, and since $\det B = 1 - d'd$, we get $\det A = (sg - a_{n1}r) (1 - d'd)$. Note that g.c.d $\{g', a_{n1}, d\} = 1$, which tells us that g' and a_{n1} are co-prime. Since g/g' is also a factor of g', we get that g.c.d $\{g, a_{n1}, d\} = 1$ as well. Hence Bézout's identity informs us that there will exist some integers s, r and d'' in \mathbb{Z} so that $sg - a_{n1}r + d''d = 1$. Hence we get that $\det A = (1 - d'd)(1 - d''d)$. In \mathbb{Z}_d , $\det A = 1$, which implies that $A \mod d$ is invertible. This proves our result for m = 1.

Assumed true for m. To prove for m+1. $a_1, a_2, \dots, a_m, a_{m+1} \in \mathbb{Z}_d^n$, such that they are LI. Our hypothesis is assumed true for m vectors, so a_1, a_2, \dots, a_m can be placed as the (left-most) columns of an $n \times n$ matrix A, which is invertible. Consider the isomorphism on \mathbb{Z}_d^n : $a \to A^{-1}a$. This maps $a_j = e_j$ for j = 1 to m, where e_j are the standard basis vectors whose only non-zero component is the j-th component, and this component is equal to 1. Let $A^{-1}a_{m+1} = (c_1, c_2, \dots, c_m, c_{m+1}, c_{m+2}, \dots, c_n)^T =: c$. Since A^{-1} is an isomorphism on \mathbb{Z}_d^n , we get that e_1, e_2, \dots, e_m and c are LI. Consider the following.

$$\begin{pmatrix} \mathbb{1}_m & 0 \\ 0 & A' \end{pmatrix} \begin{pmatrix} \mathbb{1}_m & c \\ 0 & c \end{pmatrix}, \tag{B3}$$

where the matrix on the left is an $n \times n$ matrix. The upper right 0 denotes a block matrix of size $m \times n - m$ whose entries are all 0. The lower left 0 denotes another block matrix of size $(n-m) \times m$ and whose entries are all 0. A' is an $(n-m) \times (n-m)$ matrix, as yet unspecified. The matrix on the right is of size $n \times (m+1)$, whose first m columns are the vectors e_j ordered from 1 to m, and the (m+1)-th column is c. Denote $c' := (c_{m+1}, c_{m+2}, \cdots, c_n)^T$. Our hypothesis allows us to choose A' to be invertible and such that $A'c' = (1, 0, \cdots, 0)^T$, which is a column vector of size n-m. Defining $c'' := (c_1, c_2, \cdots, c_m)^T$, so that $c^T = (c''^T, c'^T)$, define an $n \times n$ matrix T

$$T = \begin{pmatrix} \mathbb{1}_k & -c'' & 0\\ 0 & 1 & 0\\ 0 & 0 & \mathbb{1}_{n-k-1} \end{pmatrix}.$$
 (B4)

The diagonal blocks $\mathbb{1}_k$, 1 and $\mathbb{1}_{n-k-1}$ are of sizes k, 1 and n-k-1 respectively. These determine the sizes of the remaining blocks of T. When T is multiplied on the left of Eq. (B3), we get an $n \times (m+1)$ sized matrix whose columns are $e_1, e_2 \cdots, e_m$ and e_{m+1} . Denoting the left matrix in Eq. (B3) by A'', and defining $Q := TA''A^{-1}$, we see that Q is an invertible matrix, and the action of Q on a_j is to produce e_j for j=1 to m+1. This tells us that a_j is the j-th column of the matrix Q^{-1} . Thus given a_j for j=1 to m+1, we may extend it to a set of n LI vectors by adding the remaining columns as vectors to this set. This proves our result.

commutative rings with no zero divisors. In our case, when d is non-prime, \mathbb{Z}_d has zero divisors.

Appendix C: Extension of a check matrix to a symplectic matrix

Consider the stabilizer group S for an $[[n,k]]_d$ stabilizer code. For a choice of independent generators g_j of S and an ordering, one may readily construct the check matrix H using these generators (see Eq. (21)). The columns of H are linearly independent and satisfy the commutation relations given by Eq. (22). Our goal in this section of the Appendix is to prove that the $2n \times (n-k)$ matrix H may be extended to a $2n \times 2n$ symplectic matrix M.

Proposition 1. Proposition 10.4,³⁶ For each $i=1,2\cdots,n-k$, there exists some $g\in \mathscr{P}_n^{(d)}$, such that $g_ig=\omega gg_i$, and $g_jg=gg_j$ when $j\neq i$.

Proof. Our proof will be based on the proof in³⁶, which works perfectly well for the case when d=p is a prime, and $\mathbb{Z}_d=\mathbb{F}_p$ is a field. In that case, one simply seeks the solution to the following equation: $H^T\Lambda x=e_i$, where $x\in\mathbb{F}_p^{2n}$ is a column vector whose solution is sought, and $e_i\in\mathbb{F}_p^{n-k}$, is the vector whose only non-zero element is the *i*-th component, and this component is equal to 1. The existence of the solution the aforementioned equation is based on the fact that dim $\mathbb{F}_p^{n-k}=\mathrm{rank}H^T\Lambda$, thus the transformation $x\to H^T\Lambda x$ is surjective in \mathbb{F}_p^{n-k} . To establish that the same result holds true when d is a composite number, we use the following arguments: since $a_1, a_2, \cdots, a_{n-k}$ are LI, one may extend this to a basis of 2n columns vectors in \mathbb{Z}_d^{2n} (see Subsection B 3). Construct a $2n\times 2n$ matrix H_{ext} , by adding to H^T the additional n+k basis vectors, by transposing them and placing the corresponding rows below H^T . Since the rows of H_{ext} are linearly independent and since they are 2n in number, by arguments in Subsection B 1 they form a basis. Hence solutions to the equations $x^T H_{\mathrm{ext}} = e_j^T$ may be found for all $j=1,2,\cdots,2n$, which implies that H_{ext} has a left-inverse. Since the left-inverse of H_{ext} is also its right inverse, that implies that the columns of H_{ext} are also linearly independent, and since these columns are 2n in number, they also form a basis using the arguments in Subsec. B 1. Thus one may then find the solution to the equation:

$$H_{\text{ext}} x' = e_i', \tag{C1}$$

where $x', e_i' \in \mathbb{Z}_d^{2n}$. Here e_i' is the vector whose only non-zero entry is the *i*-th entry, and this entry is equal to 1, whereas x' is a vector whose solution is sought. That H_{ext} is invertible tells us that $x' = H_{\text{ext}}^{-1} e_i'$. Define $x := \Lambda^{-1} x'$. Then x satisfies the equation: $H_{\text{ext}} \Lambda x = e_i'$. One may neglect the n+k bottom most rows of H_{ext} and of e_i' (not of Λ , or of x - their rows must be maintained intact), which then gives us the equation $H^T \Lambda x = e_i$. Let g be the Pauli element whose \mathbb{Z}_d^{2n} representative is x. The commutation relations between g and the g_j 's is then determined by the equation $H^T \Lambda x = e_i$: we get that g and g_j commute when $j \neq i$ and $g_i g = \omega g g_i$. Hence proved.

Let us obtain the solutions for x for all $i = 1, 2, \dots, n - k$, and call the corresponding solutions w'_i . We want w'_i $\Lambda w_j = 0$ for all $i, j = 1, 2, \dots, n - k$. To that end, we perform a Gram-Schmidth orthogonalization procedure in the following way, starting with $w_1 := w'_1$, and starting with i = 2 to i = n - k,

$$w_i := w_i' - \sum_{j=1}^{i-1} \left(w_j^T \Lambda w_i' \right) a_j. \tag{C2}$$

If k = 0, our job is done: to extend H to a $2n \times 2n$ symplectic matrix, we need to add the columns w_i to the left (or right) of H, and this will give us our symplectic matrix M. Suppose k = 1, we need to find pairs of vectors a_n, w_n so that

$$a_n^T \Lambda a_j = 0, \ \forall \ j = 1, 2, \cdots, n,$$

$$w_n^T \Lambda w_j = 0, \ \forall \ j = 1, 2, \cdots, n,$$

$$w_n^T \Lambda a_i = \delta_{i,n}, \ \forall \ j = 1, 2, \cdots, n.$$
(C3)

At the moment, we have 2(n-1) linearly independent vectors a_j and w_j . Let a' be some other vector so that the a', a_j and w_j form a set of 2n-1 LI vectors in \mathbb{Z}_d^{2n} . Define $a_n := a' - \sum_{j=1}^{n-1} \left(w_j^T \Lambda a' \right) a_j - \sum_{j=1}^{n-1} \left(a_j^T \Lambda a' \right) w_j$. Since the linear independence of a', a_j and w_j ensures that a_n is non-zero. We have also ensured that a_n satisfies the conditions $a_j^T \Lambda a_n = w_j^T \Lambda a_n = 0$ for all $j = 1, 2, \dots, n-1$. It remains to find w_n with the properties as desired by Eq. (C3). Such a w_n may be found by using Proposition 10.3 again for an extended check matrix which is obtained by adding a_n to the original check matrix H. Thus we have obtained all the column vectors as desired, and the newly obtained column vectors may be arranged suitably to produce a $2n \times 2n$ symplectic matrix M.

If k > 1, we may apply the same procedure as above iteratively, to obtain the symplectic matrix M.

Appendix D: Symplectic conditions for M

In this section of the Appendix, we derive the symplectic conditions Eq. (30)-(33) for the matrix M in Eq. (24), which has the form:

$$M = \begin{bmatrix} M_{11} & M_{12} & 0 & M_{14} \\ M_{21} & M_{22} & 0 & M_{24} \\ M_{31} & M_{32} & A & M_{34} \\ M_{41} & M_{42} & 0 & M_{44} \end{bmatrix}.$$
(D1)

Here M_{11} , M_{31} , M_{13} (which is 0) and A are of dimensions $(n-k) \times (n-k)$, M_{22} , M_{24} , M_{42} and M_{44} are of dimensions $2k \times 2k$, M_{21} , M_{41} , M_{23} and M_{43} (which are both 0) are of dimensions $k \times (n-k)$, and finally, M_{12} , M_{14} , M_{32} and M_{34} are of dimensions $(n-k) \times k$. M satisfies the following symplectic condition.

$$M^T \Lambda M = \Lambda. \tag{D2}$$

It will be convenient to label the LHS of Eq. (D2) as follows.

$$M^{T}\Lambda M = \begin{bmatrix} Q_{11} & Q_{12} & Q_{13} & Q_{14} \\ Q_{21} & Q_{22} & Q_{23} & Q_{24} \\ Q_{31} & Q_{32} & Q_{33} & Q_{34} \\ Q_{41} & Q_{42} & Q_{43} & Q_{44} \end{bmatrix}.$$
 (D3)

Here we only consider blocks Q_{13} , Q_{23} and Q_{43} . That is the conditions given in Eq. (28) and (29):

$$Q_{13} = M_{11}^T A = I_k \Longrightarrow M_{11} = (A^{-1})^T,$$

$$Q_{23} = M_{12}^T A = 0 \Longrightarrow M_{12} = 0,$$

$$Q_{43} = M_{14}^T A = 0 \Longrightarrow M_{14} = 0.$$
(D4)

Substituing Eq. (D4) into Eq. (24) gives us Eq. (34). Now, Eq. (31), Eq.(32) and Eq. (33) are merely the equations for the blocks Q_{11} , Q_{12} and Q_{14} respectively, whereas Eq. (30) is the collective condition on $\begin{pmatrix} Q_{22} & Q_{24} \\ Q_{42} & Q_{44} \end{pmatrix}$.

Appendix E: Significance of the subgroups of $T(2n,k,\mathbb{Z}_d)$: $Sp(2k,\mathbb{Z}_d)$, $GL(n-k,\mathbb{Z}_d)$, $B_A(n,k,d)$ and $B_S(n,k,d)$

We now explain the significance of each of these subgroups. For an $[[n,k]]_d$ stabiliser code, let H be the corresponding check matrix and let a symplectic extension of H be given by M as in Eq.(24). One may obtain another symplectic matrix of the same stabiliser code by right-multiplying M with a matrix in $T(2n,k,\mathbb{Z}_d)$ with a decomposition $M_EM_SM_L$. Suppose that $M_E = M_L = I_{2n}$. Right-multiplying M with M_S transforms the check matrix H to another check-matrix HA (of the same code). Thus the $GL(n-k,\mathbb{Z}_d)$ subgroup realises the freedom of making different choices of the independent generators of said stabilizer group. Right-multiplying M with M_L transforms the columns in L_X and L_Z , while leaving E and E invariant. Thus the E invariant is subgroup realises the freedom of making different choices of the generators of the E purely logical Pauli group, which we denote by $E_n^{(d)}$. Each element of this subgroup (except the identity) will act non-trivially on the stabiliser code, i.e., codewords are mapped to other codewords under the action of these elements. Thus the E purely logical Pauli group is a subgroup of the logical Pauli group E is independent of the choice of generators of the E purely logical Pauli group (which is realised by the E purely logical Pauli group is independent of the choice of generators of the E purely logical Pauli group (which is realised by the E purely logical Pauli group). This agrees with the fact that both subgroups commute with each other.

We next explain the significance of M_E . Let g_E be a Pauli which anti-commutes with some generators of S. Consider $g_E \mathcal{L}_n^{(d)}$ which is a left coset of $\mathcal{L}_n^{(d)}$. Each element $g_E g_L$ rotates $\mathcal{C}(S)$ to the same orthogonal coding space $\mathcal{C}(S)^{\perp}$. But for different choices of g_L , $g_E g_L$ performs a distinct internal rotation within $\mathcal{C}(S)^{\perp}$. The error correction procedure is unable to distinguish between different internal rotations due to different g_L . Thus it can correct only one element in the coset $g_E \mathcal{L}_n^{(d)}$. This is true for all g_E 's which anticommute with elements in S. Which element within $g_E \mathcal{L}_n^{(d)}$ is chosen to be the correctable error depends on the error correction model. We next prove the following.

Theorem 3. The columns E in M in Eq. (24) represent correctable errors. Right multiplying M with different $M_E \in B_A(n,k,d)$ realises the freedom in making different choices of such correctable errors. The action of $M_T \in B_S(n,k,d)$ doesn't affect this choice.

Proof. That the columns of E fix the choice of the correctable errors of the stabiliser code is well-known in the stabiliser formalism for quantum error correction (see Chapter 10 in³⁶ for instance or see the beginning in Section III in³⁷). What we need to prove is that for each distinct choice of correctable errors, there is a choice of $M_E \in B_A(n,k,d)$ such that right multiplying M with M_E produces the corresponding column block of correctable errors in MM_E . To this end, suppose that the columns in E are homomorphic images of $g_1, g_2 \cdots, g_k \in \mathcal{P}_n^{(d)}$ (see Eq. (11)). Suppose that we wish for $g_1h_1, g_2h_2, \cdots, g_kh_k$ to be correctable errors instead, where $h_1, h_2 \cdots, h_k \in \mathcal{L}_n^{(d)}$. These errors may be represented in \mathbb{Z}_d^{2n} by the $2n \times k$ matrix

$$E' = E + L_X N' + L_Z L', (E1)$$

where the *j*-th columns of N' and L' are determined by how h_j decomposes into the generators of $\mathcal{L}_n^{(d)}$. Thus N' and L' are determined entirely by h_j 's. Equation (E1) may be realised by right multiplying M with a matrix of the form M_E with N' = N and L' = L, and K_A determined entirely by N and L as follows: $K_A = L^T N - N^T L$. This is just Eq. (31). Additionally, right multiplying M by any $M_T \in B_S(n,k,d)$ changes the columns in E to $E' = E + HK_S$. If $g_{E'}$ and g_E represent the $g_{E'}$ and the choice of generators in E' and E' is some element of the stabiliser group. $g_{E'}$ is determined by the $g_{E'}$ -th column of $g_{E'}$ and the choice of generators in E' and E' is some element E' and E' are both simultaneously correctable since the stabiliser element $g_{E'}$ acts trivially on the code. Hence proved.

1. Order of $Sp(2n, \mathbb{Z}_{p^m})$

A computation of the order of $\operatorname{Sp}(2n,\mathbb{Z}_{p^m})$ is given in Ref.³⁸. For the interested reader, we reproduce that computation here, while elaborating on some of the definitions and computations to make them clearer. Define a ring homomorphism $\psi_1: \mathbb{Z}_{p^m} \to \mathbb{Z}_{p^{m-1}}$ as

$$\psi_1(x) = x \bmod p^{m-1},\tag{E2}$$

where $x \in \mathbb{Z}_{p^m}$. The following remark explains Eq. (E2).

Remark 7. The ideal generated by p^{m-1} within \mathbb{Z}_{p^m} is isomorphic to \mathbb{Z}_p . We label this ideal as $p^{m-1}\mathbb{Z}_{p^m}$. The factor ring of $\mathbb{Z}_{p^m}/p^{m-1}\mathbb{Z}_{p^m}$ is isomorphic to $\mathbb{Z}_{p^{m-1}}$. ψ_1 is the map from $\mathbb{Z}_{p^m} \to \mathbb{Z}_{p^m}/p^{m-1}\mathbb{Z}_{p^m}$. ψ_1 is a ring homomorphism, and $\ker \psi_1 = p^{m-1}\mathbb{Z}_{p^m}$.

We define a group homomorphism from $\operatorname{Sp}(2n,\mathbb{Z}_{p^m}) \to \operatorname{Sp}(2n,\mathbb{Z}_{p^{m-1}})$ as follows. Let M be a $2n \times 2n$ symplectic matrix. Define $\Xi_1 : \operatorname{Sp}(2n,\mathbb{Z}_{p^m}) \to \operatorname{Sp}(2m,\mathbb{Z}_{p^{m-1}})$ as follows: the ij-th matrix element of $\Xi_1(M)$ is $\psi_1(M_{ij})$. Since $M \in \operatorname{Sp}(2n,\mathbb{Z}_{p^m})$, it satisfies the equation $M^T \Lambda M = \Lambda$ (where $\Lambda \in \operatorname{Sp}(2n,\mathbb{Z}_{p^m})$), then

$$(\Xi_1(M))^T \Lambda \Xi_1(M) = \Lambda, \text{ where } \Lambda \in \operatorname{Sp}(2n, \mathbb{Z}_{p^{m-1}}),$$
 (E3)

which implies that $\Xi_1(M) \in \operatorname{Sp}(2n, \mathbb{Z}_{p^{m-1}})$. That ψ_1 is a ring homomorphism immediately implies that Ξ_1 is group homomorphism from $\operatorname{Sp}(2n, \mathbb{Z}_{p^m})$ to $\operatorname{Sp}(2n, \mathbb{Z}_{p^{m-1}})$. ker Ξ_1 is the subgroup in $\operatorname{Sp}(2n, \mathbb{Z}_{p^m})$ which Ξ_1 maps to $I_{2n} \in \operatorname{Sp}(2n, \mathbb{Z}_{p^{m-1}})$. Note that I_{2n} (in $\operatorname{Sp}(2n, \mathbb{Z}_{p^m})$) lies in ker Ξ_1 . Thus any $M \in \ker \Xi_1$ can be written in the form $I_{2n} + K$, where K is a $2n \times 2n$ matrix with matrix elements in $p^{m-1}\mathbb{Z}_{p^m}$, and satisfies the symplectic equation

$$(I_{2n} + K)^T \Lambda (I_{2n} + K) = \Lambda,$$

$$\Rightarrow K^T \Lambda + \Lambda K + K^T \Lambda K = 0.$$
 (E4)

Note that $K^T \Lambda K = 0$ since K can be written as $K = p^{m-1}K'$, with $K' \in \{0, 1, \dots, p-1\} \subset \mathbb{Z}_{p^m}$, which gives us $K^T \Lambda K = p^m \ p^{m-2}K'^T \Lambda K'$ which is 0 since $p^m = 0$ in \mathbb{Z}_{p^m} . Thus K has to satisfy the equation $\Lambda K = -K^T \Lambda$. Decomposing K into $n \times n$ blocks as follows:

$$K = \begin{bmatrix} X & Y \\ Z & W \end{bmatrix}, \tag{E5}$$

we see that the symplectic condition becomes $X = -W^T$, $Y = Y^T$ and $Z = Z^T$. Thus K has $2n^2 + n$ unconstrained matrix elements which take values in $p^{m-1}\mathbb{Z}_{p^m}$, which has p elements. Thus $|\ker \Xi_1| = p^{2n^2+n}$ and we get

$$\left| \operatorname{Sp}(2n, \mathbb{Z}_{p^m}) \right| = p^{2n^2 + n} \left| \operatorname{Sp}(2n, \mathbb{Z}_{p^{m-1}}) \right|.$$
 (E6)

Similarly one may define $\psi_j: \mathbb{Z}_{p^{m-j+1}} \to \mathbb{Z}_{p^{m-j}}$ and $\Xi_j: \operatorname{Sp}(2n, \mathbb{Z}_{p^{m-j+1}}) \to \operatorname{Sp}(2n, \mathbb{Z}_{p^{m-j}})$, with $|\ker \Xi_2| = p^{2n^2+n}$ for $j = 2, 3, \dots, m-1$, and

$$\left| \operatorname{Sp}(2n, \mathbb{Z}_{p^{m-j+1}}) \right| = p^{2n^2+n} \left| \operatorname{Sp}(2n, \mathbb{Z}_{p^{m-j}}) \right|, \tag{E7}$$

which finally gives us

$$\left| \operatorname{Sp}(2n, \mathbb{Z}_{p^m}) \right| = p^{(m-1)(2n^2+n)} \left| \operatorname{Sp}(2n, \mathbb{Z}_p) \right|.$$
(E8)

It is known (see Ref.³⁵) that $|\operatorname{Sp}(2n,\mathbb{Z}_p)| = p^{n^2} \prod_{j=1}^n (p^{2j} - 1)$, which tells us that

$$\left| \operatorname{Sp}(2n, \mathbb{Z}_{p^m}) \right| = p^{(2m-1)n^2 + (m-1)n} \prod_{j=1}^n (p^{2j} - 1).$$
 (E9)

Appendix F: Proof of Eq. (63)

We start by noting that

$$1 \le \frac{1 - p_i^{-2(n-j)}}{1 - p_i^{-(n-k-j)}},\tag{F1}$$

we get $\left(p_i^{m_i}\right)^{\frac{(n-k)(n+3k+1)}{2}} \leq \mathscr{C}[[n,k]]_{p_i^{m_i}}$ and using Eq. (53) we immediately get $d^{\frac{(n-k)(n+3k+1)}{2}} \leq \mathscr{C}[[n,k]]_d$. For the inequality on the right, we note that $\frac{1-p^{-2(n-j)}}{1-p^{-(n-k-j)}} \leq \frac{1}{1-p^{-(n-k-j)}} \leq \frac{1}{1-2^{-(n-k-j)}}$, since $p \geq 2$. Thus

$$\prod_{j=0}^{n-k-1} \frac{1-p^{-2(n-j)}}{1-p^{-(n-k-j)}} \, \leq \, \prod_{j=0}^{n-k-1} \left(1+\frac{2^{-(n-k-j)}}{1-2^{-(n-k-j)}}\right) \, \leq \prod_{j=0}^{n-k-1} \, \left(1+2^{-j}\right), \tag{F2}$$

where we used the fact that $\frac{2^{-(n-k-j)}}{1-2^{-(n-k-j)}} \le 2^{-(n-k-1-j)}$, and re-label $j \to n-k-1-j$. and using

$$\prod_{j=0}^{n-k-1} \left(1 + 2^{-j} \right) \le \prod_{j=0}^{\infty} \left(1 + 2^{-j} \right) = \phi\left(\frac{1}{2}\right), \tag{F3}$$

where $\phi(.)$ is the Euler function⁴⁴⁰, which is a special case of the q-Pochhammer function⁴¹. Using Mathematica, this evaluates to 4.768 or $e^{1.56}$.

Appendix G: Why the proof of Theorem 20. of Ref. 22 fails for counting $[[n,k]]_d$ modular stabilizer codes

We first give the context of Theorem 20 in Ref.²²: $d=p^m$, for some positive prime integer p, and some positive integer p, and the vector space in question is $\mathbb{F}_{p^m}^{2n}$, which is defined over the field \mathbb{F}_{p^m} . Here \mathbb{F}_{p^m} is the Galois field extension of the base field $\mathbb{F}_p \simeq \mathbb{Z}_p$. Thus this overlaps with our scenario only when d=p. The proof counts the number of isotropic subspaces of $\mathbb{F}_{p^m}^{2n}$. An isotropic subspace is spanned by n-k linearly independent vectors $a_1, a_2, \cdots, a_{n-k}$, with the additional property that the symplectic inner products between any pair of these vectors is zero, i.e., $a_i^T \Lambda a_j = 0$, for all $i, j = 1, 2, \cdots, n-k$. The proof in Ref.²² begins by counting the number of possible values that a_1 can take in $\mathbb{F}_{p^m}^{2n}$. Since any non-zero vector in $\mathbb{F}_{p^m}^{2n}$ is a potential candidate, the count is $d^{2n} - 1$. Subsequently it counts the possible values which a_2 can be in $\mathbb{F}_{p^m}^{2n} \setminus \text{span}\{a_1\}$. Note that a_2 has to satisfy the additional condition that $a_1^T \Lambda a_2 = 0$. Viewing the transformation $a_2 \to a_1^T \Lambda a_2$ as a linear map on $\mathbb{F}_{p^m}^{2n}$, one notes that the symplectic condition simply demands that a_2 belongs to the kernel of this linear map. The rank-nullity theorem is then used to tell us that a_2 belongs to a 2n-1 dimensional subspace of $\mathbb{F}_{p^m}^{2n}$. But since a_1 also belongs to the kernel, and since a_2 and a_1 need to be linearly independent, we need to rule out the possibility that $a_2 = xa_1$ for all $x \in \mathbb{F}_{p^m}$. This gives us

⁴ Not the Euler totient function

the count for a_2 . The same technique is applied iteratively to give us counts for all a_j for $j = 3, \dots, n-k$. The product of these counts gives the number of distinct ordered sets of n-k "symplectic" orthogonal vectors. To obtain the number of isotropic subspaces, one needs to divide the aforementioned product by the redundancy with which each subspace is counted. It is easily seen that this redundancy is equal to the number of distinct ordered "symplectic" orthogonal basis any isotropic subspace has.

Coming to our scenario, we note that the first step of the proof of Theorem 20 in Ref.²² itself is incompatible with our context, since in our case we need to rule out all the non-zero linearly *dependent* vectors for a_1 , i.e., non-zero vectors a_1 which satisfy the equation $xa_1 = 0$, for a non-zero x. When d is not prime, such vectors exist. For example consider the vector $a = (2,0,0,0,0,0)^T$ for d = 4, n = 3 and k = 2. $2a = (4,0,0,0,0,0)^T = 0$ in \mathbb{Z}_4 . If one still manages to weed out those non-zero vectors for a_1 , one still encounters similar issues for counting a_j for $j \ge 2$. According to Lemma 1 the total number of ordered linear independent vectors a_j whose symplectic inner products are zero should be

$$\begin{split} &\frac{|\operatorname{Sp}(2n,\mathbb{Z}_d)|}{|B_S(n,k,d)|\ |B_A(n,k,d)|\ |\operatorname{Sp}(2k,\mathbb{Z}_d)|}, \text{ when } k \geq 1\\ &\frac{|\operatorname{Sp}(2n,\mathbb{Z}_d)|}{|B_S(n,0,d)|}, \text{ when } k = 0. \end{split} \tag{G1}$$

This is because the redundancy, which is equal to the number of choices for ordered bases, is $|GL(n-k,\mathbb{Z}_d)|$. Thus one may relate our method with the method used by Gross in Theorem 20 in Ref.²² in this way. It is possible that one may compute the expression in Eq. (G1) by suitably modifying the counting technique in Ref.²², but our method has the advantage of being conceptually richer.