

BMJ Open Project20: maternity care mechanisms that improve access and engagement for women with social risk factors in the UK – a mixed-methods, realist evaluation

Hannah Rayment-Jones ¹, Kathryn Dalrymple ¹, James M Harris,² Angela Harden,³ Elidh Parslow,⁴ Thomas Georgi,⁵ Jane Sandall¹

To cite: Rayment-Jones H, Dalrymple K, Harris JM, *et al*. Project20: maternity care mechanisms that improve access and engagement for women with social risk factors in the UK – a mixed-methods, realist evaluation. *BMJ Open* 2023;**13**:e064291. doi:10.1136/bmjopen-2022-064291

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-064291>).

Received 27 April 2022
Accepted 28 December 2022



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY. Published by BMJ.

¹Women and Children's Health, King's College London, London, UK

²Women's Health, Chelsea And Westminster NHS Foundation Trust, London, UK

³Department of Health Services Research and Management, City University of London, London, UK

⁴Maternity and Women's Health, North Middlesex University Hospital NHS Trust, London, UK

⁵Department of Women and Children's Health, King's College London, London, UK

Correspondence to

Dr Hannah Rayment-Jones; hannah.rayment-jones@kcl.ac.uk

ABSTRACT

Objectives To evaluate how women access and engage with different models of maternity care, whether specialist models improve access and engagement for women with social risk factors, and if so, how?

Design Realist evaluation.

Setting Two UK maternity service providers.

Participants Women accessing maternity services in 2019 (n=1020).

Methods Prospective observational cohort with multinomial regression analysis to compare measures of access and engagement between models and place of antenatal care. Realist informed, longitudinal interviews with women accessing specialist models of care were analysed to identify mechanisms.

Main outcome measures Measures of access and engagement, healthcare-seeking experiences.

Results The number of social risk factors women were experiencing increased with deprivation score, with the most deprived more likely to receive a specialist model that provided continuity of care. Women attending hospital-based antenatal care were more likely to access maternity care late (risk ratio (RR) 2.51, 95% CI 1.33 to 4.70), less likely to have the recommended number of antenatal appointments (RR 0.61, 95% CI 0.38 to 0.99) and more likely to have over 15 appointments (RR 4.90, 95% CI 2.50 to 9.61) compared with community-based care. Women accessing standard care (RR 0.02, 95% CI 0.00 to 0.11) and black women (RR 0.02, 95% CI 0.00 to 0.11) were less likely to have appointments with a known healthcare professional compared with the specialist model. Qualitative data revealed mechanisms for improved access and engagement including self-referral, relational continuity with a small team of midwives, flexibility and situating services within deprived community settings.

Conclusion Inequalities in access and engagement with maternity care appears to have been mitigated by the community-based specialist model that provided continuity of care. The findings enabled the refinement of a realist programme theory to inform those developing maternity services in line with current policy.

INTRODUCTION

As a core component of maternity care across the globe, the adequacy of antenatal care is

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The specialist models of care evaluated in this study are situated in the UK's complex maternity system and can be appropriately investigated using a mixed-methods, realist evaluation design.
- ⇒ Both qualitative and quantitative methods are used to understand not only if access and engagement are improved by specialist models of care, but how and in what context.
- ⇒ Longitudinal interviews were undertaken to increase trust between participants and the researcher and lessen the perception of surveillance.
- ⇒ 'The relatively small and varied numbers in each quantitative data group, and the multiple testing required to establish the separate effects of the model of care, place of care and service attended might result in a change in statistical power, reducing the probability of detecting effects when they do exist.'
- ⇒ The generalisability of the findings is limited by the urban location of both specialist models of care evaluated and the UK health system context.

measured by the timing of antenatal care access and the number of appointments attended.¹ The routine functions of antenatal care include support, health promotion, screening and diagnosis, disease prevention and additional care for women at higher risk.² When these functions are of high quality and care is well attended, it makes a crucially important contribution to the reduction of health inequalities.^{3 4} The WHO¹ recommends a minimum of eight antenatal appointments to reduce perinatal mortality and improve women's experiences. In high-income countries such as the UK, antenatal care coverage is consistently high and correlates with relatively low maternal and infant mortality rates when compared with the low-income and middle-income countries.⁵ Despite this, there are marked inequalities in access, health outcomes and women's




Figure 1 Programme theory (PT) refinement process leading to context+mechanism=outcome (CMO) configuration.

experiences with poor antenatal care uptake for women from lower socioeconomic and minority groups.^{6–10} The most deprived women in the UK are 60% less likely to have received any antenatal care when compared with the least deprived women.⁶

Factors associated with inequalities in maternal and neonatal outcomes include black and minority ethnicity, poverty, young motherhood, homelessness, difficulty speaking or understanding English, migrant or refugee status, domestic violence, mental illness and substance abuse.^{7 11–13} It is hypothesised that a lack of access and engagement with maternity services is directly linked to health inequalities for women with these social risk factors.^{14–17} Therefore, policies to tackle inequalities often focus on improving access to maternity services and continuity of midwifery care.^{18 19} There is a strong evidence base documenting improved maternity outcomes and experiences for women who receive continuity models of midwifery care²⁰ but less is known about what works to improve access and engagement for women with social risk factors.²¹ The impact of place-based maternity care is also poorly understood. The specialist models of care being evaluated in this study provide continuity of care across the antenatal, intrapartum and postnatal periods from a one-to-one midwife or small team of midwives whom women have had the opportunity to meet during pregnancy. One model is based in the community setting and provides care to women in the local, deprived catchment area, the other is based in a large teaching hospital and women with significant social risk factors are referred to the model of care early in pregnancy. See online supplemental file 1 for full definitions of each type of maternity care women may experience, and those being compared in this study. As part of a wider evaluation these two UK-based specialist models, this study aimed to describe and compare service use for women with and without social risk factors by asking:

1. Are women with social risk factors more likely to be offered the specialist models of care over those without? If so, do they find this acceptable?
2. Compared with standard maternity care and group practice models, do specialist models improve access and engagement with maternity services? If so, for whom, in what context and how?

3. Do specialist models of care increase the quality of relational continuity and reduce the number of appointments women miss?
4. How does the location of a model of care—community or hospital, impact on women's access and engagement with maternity services?

METHODS

Study design

Realist methodology is a theoretically-informed approach to evaluating complex interventions and often uses mixed methods to understand *how* interventions, in this case models of maternity care, do or do not work in different contexts.²² This allows those implementing services such as specialist models of care to refine, scale-up or even withdraw the service.²³ The aims of this study were approached through the testing and refinement of initial programme theories constructed in an earlier synthesis of literature⁷ and focus groups with midwives²⁴ relating to how women with social risk factors access and engage with maternity care (see figure 1 for the theory refinement process). Retroductive theorising was used to uncover meaningful causal mechanisms, often focusing on how the wider context and human response to different aspects of maternity care leads to specific outcomes.²⁵ This approach offers an epistemologically and ontologically grounded way of integrating mixed methods, often analysing qualitative data to find the causal relationship behind quantitative findings.²⁶ Routinely collected pregnancy and birth outcome data for 1000 women accessing different maternity models, including standard care, group practice and specialist models of care at two large, inner-city maternity services were prospectively collected and analysed. Longitudinal interviews with 20 women with social risk factors were conducted to uncover the mechanisms for any associations found in the cohort analysis. Online supplemental file 1 provides definitions of the two service provider settings and describes the different models of care women might experience at each. Online supplemental file 2 demonstrates the detailed, methodological process of programme theory refinement and development of the context+mechanism=outcome (CMO) configuration and provides adjusted and unadjusted data.

All participants provided written consent by signing a consent form approved by the ethics committee. The quantitative data collected were anonymised before being made available to the research team. A service user panel contributed to the study through external peer review for funding, the research focus, design and analysis of data. The Standards for Quality Improvement Reporting Excellence checklist was used to report this new knowledge for use in healthcare services.²⁷

Patient and public involvement

Multiple representative, diverse groups of service users were involved in the planning and development of this

research. They were recruited through local community groups, clinicians and existing patient involvement groups. Using participatory appraisal methods and online engagement events, recent maternity service users provided feedback on the protocol, study materials, interview guides and refinement of programme theories. They also prioritised outcome measures and reviewed the qualitative data analysis. Training needs were identified by the service users for analysis of quantitative data and further research addressing maternal health inequities.²⁸

Data collection

Quantitative data collection

A power calculation was based on a previous analysis of UK antenatal care usage⁶ and validated metrics for monitoring local inequalities in access to care at a service evaluated in this research.²⁹ We calculated that with 250 women in each group (those receiving standard maternity care and those receiving group practice or a specialist model of care), we would have 90% power to detect a 15% difference in timely access to antenatal care (before 12+6 weeks gestation) between the different models of care with 500 anonymised birth records accessed at each trust. Pregnancy and birth outcome data were extracted from computerised records at each service provider. Data collection was prospective; the demographics of the first 500 women booking for maternity care with each service provider in January 2019 were collected, with pregnancy and birth outcomes collected later in the year when all women had been discharged from maternity care. Deprivation deciles calculated using the 2019 English Indices of Deprivation³⁰ were grouped into four groups of sufficient numbers to enable comparisons between groups of similar numbers.

Semi-structured, longitudinal interviews with 20 women with low socioeconomic status and social risk factors who were receiving specialist care from one of the two service providers were carried out at approximately 28 and 36 weeks' gestation, and 6 weeks post birth. Women were identified by the specialist model midwives providing their care if they met the following inclusion criteria: A deprivation score³¹ of higher than 30 and/or secondary school as the highest level of education attained. An interview guide was developed using previous literature^{7,24,32} to elicit specific mechanisms explaining how the specialist model might improve access and engagement.

Analysis

The quantitative data were analysed using Stata V.16.0. First, women's social risk factors, ethnicity, socioeconomic status and medical characteristics were described using descriptive statistics and stratified by the service provider attended to enable comparisons of differences in the samples between each service. Variables were tested for bivariate association using χ^2 tests and t-tests, for dichotomous and continuous variables, respectively. X^2 analyses were also performed to test for associations between socioeconomic position by deprivation (indices

of multiple deprivation (IMD)) decile,³³ as well as social and medical risk factors. Second, three regression models were developed: Model 1 adjusted for ethnicity, age, parity, deprivation score, social risk factors and medical risk status; Model 2 included model 1, plus adjustment for the service provider that women attended to consider differences in organisation guidelines, processes and culture; and Model 3 included model 2, however, the place of antenatal care (hospital vs community-based antenatal care) was treated as the independent variable. This structured model allowed us to explore the association between access and engagement depending on the model of care received, while accounting for interactions between independent variables to predict the dependent variable. Risk ratios and CIs are used to demonstrate statistical significance as well as the direction and strength of the effect.³⁴ Definitions for all demographics and social risk factors are given in online supplemental file 1.

The qualitative data were coded using NVivo V.12 and analysed using a thematic framework analysis.³⁵ This allowed for the organisation of a large qualitative data set into a coding framework developed using previously constructed programme theories,^{7,24} to uncover new theories and differences in women's experiences depending on their characteristics.³⁵ The method suited the longitudinal approach to data collection as changes in women's perceptions and relationships with healthcare providers could be seen over the course of their pregnancy and post-natal period. We used existing models of data adequacy³⁶ to assess acceptable data quality. The initial programme theories relating to access and engagement were refined and constructed as a CMO configurations that present specific aspects of the specialist models of care.

RESULTS

Full pregnancy and birth data were collected and analysed from 799 women accessing care across the two service providers. A total of 201 sets of birth outcome data were missing due to sample dropout, that is, those women stopped receiving care at the service and were therefore excluded from the final analysis. The total number and reasons given for women with missing outcome data at each hospital did not differ significantly. See figure 2 for the data collection flowchart.

Demographics of women included in the quantitative data analysis

Table 1 presents the demographic profiles of the 799 women who continued their pregnancies and gave birth at the two services, with significant differences in ethnicity, social and medical risk factors, and model of care received highlighted and adjusted for in the analysis of access and engagement outcomes. See online supplemental file 1 for definitions relating to risk factors and model of care.

Characteristics of the women interviewed

Twenty pregnant women with low socioeconomic status and/or educational attainment were recruited (see




Figure 2 Quantitative data collection flowchart.

table 2). All 20 women received a specialist model of care, 8 were first time mothers and the other 12 had between one and eight children. For five of the multiparous women, this was their first pregnancy in the UK. Based on 2019 deprivation scores,³⁰ 19 participants were in the first or second most deprived deciles, with 1 participant in the third and fourth decile group. Twelve participants were born outside of the UK, and nine did not speak English and required an interpreter. All participants were experiencing between one and seven social risk factors including common or severe mental health issues, domestic violence, drug/alcohol misuse, no support, single motherhood, financial and housing problems, learning disability, sexual abuse, trafficking, female genital mutilation and no recourse to public funds. Five participants were seeking asylum, had refugee status or had had an asylum claim refused and nine had social care involvement during their pregnancy.

Model of care accessed by deprivation score and risk factors

When merging data for the two service providers table 3 shows that women in the most deprived deciles were significantly more likely to receive a specialist model of care, and women in the least deprived deciles were less likely to receive community-based antenatal care than hospital-based antenatal care. This reflects the aim of the service providers to offer women in more deprived areas both specialist and/or community-based antenatal care. A statistically significant relationship was found between deprivation score and the number of social risk factors recorded.

Testing using qualitative data

Women reported different pathways into the specialist models including direct referrals from a general practitioner (GP), midwives, sexual health clinics, social workers, emergency departments and self-referral. Overall, women

were pleased to be referred to the specialist model, however they were often not made aware of *why* they had been referred and described a lack of choice.

I don't know if it's because I have these issues, only, or if it's the team or if it's the hospital or, I don't know, or the combination. I have no idea. But I'm super-happy and feeling lucky I ended up like having all this. like how many people get the support we get here? Is it just me because I have mental health?' (community-based model (CBM)10)

Timing of access to maternity services

Quantitative analysis 1—model of care

The quantitative data were analysed to test the hypothesis that the specialist model of care has an impact on the timing of access to maternity care, known as the 'booking appointment'. Table 4 shows that no relationship was found between the model of care received and the gestation at which women attended their booking appointment. The base outcome was set for less than 10 weeks' gestation at the booking appointment to reflect the National Institute for Care Excellence guidance for women with complex social factors.¹⁵ When adjusting for women's characteristics (see online supplemental file 2 for fully adjusted outcome data tables) primiparous women (risk ratio (RR) 1.79, 95% CI 1.02 to 3.12), those with social risk factors (RR 1.93, 95% CI 1.03 to 3.62) and those with high medical risk status at the booking appointment (RR 2.49, 95% CI 1.18 to 5.25) were more likely to book for maternity care later than 20 weeks.

Quantitative analysis 2—place of antenatal care

Table 5 shows that after adjusting for the model of care received and service attended, women receiving hospital-based antenatal care were significantly more likely to attend their booking appointment after 20 weeks' gestation (RR 2.51, 95% CI 1.33 to 4.70).

Qualitative data analysis

The qualitative data were analysed to explore why timing of access to maternity care was different for women attending hospital-based antenatal care, primiparous women and those with high medical risk status and social risk factors. Many women expressed wanting to be seen earlier in pregnancy and described not feeling valued by maternity services until their pregnancy is viable. Delays in accessing maternity services was often due to difficulties in accessing GP services and a convoluted referral pathway between community and hospital services. Community-based care and the ability to self-refer to maternity services appeared to reduce the time women spent waiting for their booking appointment.

[at 3 weeks gestation] I called the GP to book an appointment. to say that I'm pregnant, and they sent me... to buy a Clearblue to check it because the list is too long. But it take a very long time, to see the GP... to refer me to the midwife, I was waiting a long time...

Table 1 Women's demographics at each service provider

Demographic variable	Service A, n (%) Total data=405	Service B, n (%) Total data=394	Total, n (%) Total data=799	X ² , p value
Ethnicity				p<0.001
Asian	37 (9)	53 (13)	90 (11)	
Black African	31 (8)	46 (12)	77 (10)	
Black Caribbean	23 (6)	16 (4)	39 (5)	
Black other	8 (2)	14 (4)	22 (3)	
Mixed	12 (3)	7 (2)	19 (2)	
White British	98 (24)	58 (15)	156 (20)	
White other	80 (20)	139 (36)	219 (27)	
Unknown	116 (29)	61 (15)	177 (22)	
Age				p=0.356
≤20	6 (1)	4 (1)	10 (1)	
21–24 years	19 (5)	32 (8)	51 (6)	
25–29 years	63 (16)	56 (14)	119 (15)	
30–34 years	134 (33)	125 (32)	259 (32)	
≥35 years	183 (45)	177 (45)	360 (45)	
Parity				p=0.167
Primiparous	212 (52)	187 (47)	399 (50)	
Multiparous	193 (48)	207 (53)	400 (50)	
Medical risk				
High at booking	118 (29)	106 (27)	224 (28)	p=0.496
High at onset of labour	152 (38)	223 (57)	375 (47)	p<0.001
Social risk factor				
Domestic abuse	23 (6)	17 (4)	40 (5)	p=0.377
Common mental health	4 (1)	34 (9)	38 (5)	p<0.001
Severe mental health	2 (<1)	8 (2)	10 (1)	p=0.051
Non-English speaking	16 (4)	48 (13)	64 (8)	p<0.001
Social care involvement	27 (7)	29 (7)	56 (7)	p=0.701
Drug/alcohol abuse	1 (<1)	10 (3)	11 (1)	p<0.001
Unsupported/single	1 (<1)	11 (3)	12 (2)	p<0.001
Financial/housing	15 (4)	31 (8)	46 (6)	p<0.001
Learning disability	6 (2)	5 (1)	11 (1)	p=0.797
Sexual abuse/trafficked	4 (2)	5 (1)	9 (1)	p=0.677
Asylum seeker/refugee	8 (2)	7 (2)	15 (2)	p=0.836
Female genital mutilation (FGM)	0	11 (3)	11 (1)	p<0.001
No recourse to public funds	6 (1)	0	6 (1)	p<0.001
No of social risk factors				p<0.001
None	337 (83)	279 (70)	616 (77)	
1	43 (11)	61 (15)	104 (13)	
2	13 (3)	26 (7)	39 (5)	
3	6 (1)	15 (4)	21 (3)	
4	5 (1)	9 (2)	14 (2)	
≥5	1 (<1)	4 (1)	5 (1)	
Name of model of care				p<0.001
Standard care	256 (63)	213 (54)	469 (59)	
Group practice	77 (19)	144 (37)	221 (28)	
Specialist	59 (15)	21 (5)	80 (10)	
Private care	13 (3)	16 (4)	29 (4)	

Continued

Table 1 Continued

Demographic variable	Service A, n (%) Total data=405	Service B, n (%) Total data=394	Total, n (%) Total data=799	χ^2 , p value
Place of model of antenatal care				p<0.001
Standard model in hospital	100 (25)	212 (54)	312 (39)	
Standard model in community	156 (40)	1 (0)	157 (20)	
Group practice in community	40 (10)	94 (24)	134 (17)	
Group practice in hospital	37 (9)	50 (13)	87 (11)	
Specialist model in community	59 (15)	2 (1)	61 (8)	
Specialist model in hospital	0	19 (5)	19 (2)	
Private care	13 (3)	16 (4)	29 (4)	
By place of antenatal care only*				p<0.001
Hospital-based	137 (35)	281 (74)	418 (54)	
Community-based	255 (65)	97 (26)	352 (46)	

*Including all models of care

when I saw the midwife the first time I was around 20 weeks. (CBM7)

Engagement—number of antenatal appointments attended

Quantitative analysis 1—model of care

Women's engagement with services was tested through the number of antenatal appointments women attended. **Table 6** shows no significant relationship between the model of care received and the number of antenatal appointments attended. Women with social risk factors were significantly more likely to have more than 15 antenatal appointments than those with no social risk factors (RR 2.57, 95% CI 1.30–5.07), as were women with high medical risk (RR 2.70, 95% CI 1.21 to 6.03) (see online supplemental file 2).

Quantitative analysis 2—place of antenatal care

Table 7 shows that women receiving hospital-based care were significantly likely to have less than the recommended number of appointments,^{1 2 15} (RR 0.61, 95% CI 0.38 to 0.99), and much more likely to have over 15 appointments (RR 4.90, 95% CI 2.50 to 9.61) than those receiving community-based care regardless of their medical risk status and other confounding factors.

Qualitative analysis

Flexible care was discussed by women accessing both specialist models, in terms of when and where appointments were scheduled and how long each appointment lasted. The community or home setting was seen as more convenient and supportive, particularly for those who were unfamiliar with UK transport systems, and those with little resources or young children. Women from both specialist models expressed ease of contacting the specialist model midwives through phone call, text messaging or email, and felt that it reduced the number of face-to-face appointments they needed and that they could rely on the midwives to remind them of appointments. Women who did not speak fluent English (45%)

particularly valued the ability to text their midwife as it gave them the opportunity to use translation technology. This flexibility appeared to encourage women to seek help more readily when they were concerned or needed reassurance. However, women in the hospital-based model did not always feel that the way care was scheduled suited their needs but structured around the hospital protocol and organisational efficiency.

If it's [antenatal care] like near to me it's OK but... I have so much financial problems so when she [HBM midwife] came to my house like, so much easier for me....and at the beginning I didn't know how to use bus... she helped me. (hospital-based model (HBM)6)

The quality of relational continuity and missed appointments

Quantitative analysis 1—model of care

The quantitative data were analysed to test the hypothesis that the specialist model of care increases the quality of relational continuity through the number of antenatal appointments and care in labour from a known healthcare professional (midwife, GP or obstetrician), and reduces the number of appointments women miss, or do not attend. **Table 8** shows that women receiving group practice and specialist models were significantly more likely to see a known healthcare professional during pregnancy. When adjusting the model for women's characteristics black African women were the least likely group to see a known healthcare professional more than five times (RR 0.17, 95% CI 0.03 to 0.82). Women receiving care in the specialist model were significantly more likely to be looked after in labour by a known healthcare professional than the group practice (RR 0.44, 95% CI 0.21 to 0.92).

Quantitative analysis 2—place of antenatal care

Table 9 shows that women receiving hospital-based antenatal care were less likely to see a known healthcare professional for their antenatal appointments compared

Table 2 Demographics of women interviewed

Characteristic	Community-based model n=10	Hospital-based model n=10	Total n (%) n=20
Ethnicity and migration status			
Born outside the UK:	7	5	12 (60)
Asian	0	2	2 (10)
Black African	3	0	3 (15)
Black Caribbean	0	1	1 (5)
White	4	2	6 (30)
Asylum seeker/refugee*	2	3	5 (25)
Born inside the UK:	3	5	8 (40)
Asian British	1	1	2 (10)
Black British	2	1	3 (15)
White British	0	3	3 (15)
Not proficient in English language/interpreter required	5	4	9 (45)
Age			
18–24	0	3	3 (13)
25–29	1	1	2 (2)
30–34	5	5	10 (50)
>34	4	1	5 (25)
Parity			
Primiparous	5	3	8 (40)
IMD decile (2019)			
Most deprived 1st+2nd	9	10	19 (95)
3rd and 4th	1	0	1 (5)
Least deprived 5th to 10th	0	0	0
No. of social risk factors			
1	3	0	3 (15)
2	0	2	2 (10)
3	2	0	2 (10)
4	1	1	2 (10)
≥5	4	7	11 (55)
Mental illness			
Common	5	9	14 (70)
Severe	1	1	2 (15)
Level of education			
Secondary school only	5	6	14 (70)
Completed college	4	3	2 (15)
Completed university	1	1	4 (20)
High medical risk at booking	7	5	12 (60)

*Including women whose asylum claim had been refused.
IMD, indices of multiple deprivation.

with those receiving community-based care. No significant relationship was found between the place of care and the number of women cared for in labour by their named midwife.

Missed appointments

Quantitative analysis 1—model of care

Table 10 shows that no significant relationship was found between model of care and the number of missed appointments. When adjusting for women's characteristics,

multiparous women were four times more likely (RR 4.50, 95% CI 1.13 to 17.82) to miss four or more appointments than primiparous women, and the older a woman was less likely that she will miss four or more appointments (RR 0.03, 95% CI 0.00 to 0.55). Black African women (RR 12.85, 95% CI 2.42 to 68.07), and women with social risk factors (RR 2.26, 95% CI 1.14 to 4.47) were more likely to miss two or more appointments (see online supplemental file 2). These findings should be viewed with caution due to the wide CIs.

Analysis 2—place of antenatal care

Table 11 shows no significant relationship was found between the number of missed appointments and the place of antenatal care.

Qualitative analysis

Women accessing the community-based model perceived a high level of continuity of care from the whole team. Women accessing the hospital-based model discussed how they knew their named midwife but not necessarily the rest of the team, impacting on how they sought help. They were also not always aware of the level of continuity offered, expressed more anxiety about their labour care and discussed the impact of midwives needing to cancel appointments. For some, cancelled appointments and a lack of continuity had a significant impact on their support and engagement, leading to social care involvement. Women in both models described feeling comfortable when rebooking appointments; they were unable to attend that appeared to reduce the number of 'missed' appointments recorded, seen as a mitigating effect of the specialist model.

If I'm running late or something comes up I just text her or give her a call and she'll muddle things around. (HBM9)

Summary of findings

Table 12 summarises the quantitative findings in relation to either the model of care received, or the place of antenatal care, or both. Characteristics of women at disproportionate risk and differences between the services providers is also presented.

Refined programme theory

The qualitative interviews with women enabled the refinement of previously constructed programme theories,^{7 24} giving insight into specific, underlying mechanisms of improved access and engagement with maternity services. The process of refinement using the data analysed is detailed in online supplemental file 2. The CMO configurations are detailed in table 13, providing a framework for those developing future models of care for women with social risk factors.

DISCUSSION

In response to concerns that the most affluent and lowest risk women are often the most likely to receive the

Table 3 Maternity care received, and risk factors recorded by deprivation score

Maternity care and risk factors	Least deprived (7th, 8th, 9th + 10th)	5th and 6th deciles	3rd and 4th deciles	Most deprived (1st + 2nd deciles)	Total n (%)	X ² p value
Model of care	Total=149	Total=158	Total=286	Total=206	N=799	p<0.001
Standard care	81 (54)	98 (62)	163 (57)	127 (62)	469 (58)	
Group practice	43 (29)	41 (26)	85 (30)	52 (25)	221 (28)	
Specialist	3 (2)	16 (10)	34 (12)	27 (13)	80 (10)	
Private care	22 (15)	3 (2)	4 (1)	0 (0)	29 (4)	
Place of antenatal care*	Total=127	Total=155	Total=282	Total=206	Total=770	p<0.001
Hospital-based	82 (65)	91 (59)	134 (48)	111 (54)	418 (54)	
Community-based	45 (35)	64 (41)	148 (52)	95 (46)	352 (46)	
Social risk factors	Total=149	Total=158	Total=286	Total=206	N=799	p<0.001
None	133 (89)	132 (84)	212 (74)	139 (67)	616 (77)	
One	10 (7)	17 (11)	50 (17)	27 (13)	104 (13)	
Two	5 (3)	4 (3)	10 (4)	20 (10)	39 (5)	
Three	1 (1)	2 (1)	10 (4)	8 (4)	21 (3)	
Four or more	0	3 (2)	4 (1)	12 (6)	19 (2)	
Medical risk						
High at booking	37 (25)	46 (29)	85 (30)	56 (27)	224 (28)	p=0.743
High at onset of labour	62 (42)	79 (50)	138 (48)	96 (47)	375 (47)	p=0.475

*Excludes private care.

highest standards of maternity care,³⁷ a clear example of Tudor Hart's Inverse Care Law,³⁸ this study examines a shift change to more proportionate universalism through services targeting those who are at highest risk of poor outcomes. The aims of the specialist models of care to reach those who are most at risk of poor outcomes and experiences, are being met. This is an important finding as it addresses the aims of the National Health Service 10-year plan³⁹ to improve outcomes and care experiences

through the offer of continuity of care to black and minority ethnic women and those living in deprived areas. The findings suggest that placing continuity of care models in areas of deprivation are likely to identify women who are experiencing social risk factors during pregnancy, thus providing them with an enhanced level of care whether they have disclosed risk factors or not. In contrast, strict inclusion criteria to hospital-based models risk missing those women who are at increased risk but are

Table 4 Gestation at booking appointment in relation to the model of care received

Gestation at booking	Model of care	Number of women (%)	Unadjusted RR (95% CI)	Model 1 Adjusted RR (95% CI)*	Model 2 Adjusted RR (95% CI)†	Model 3 Adjusted RR (95% CI)‡
<10 weeks	Standard	197 (62)	Ref	Ref	Ref	Ref
	Group	87 (27)	Ref	Ref	Ref	Ref
	Specialist	35 (11)	Ref	Ref	Ref	Ref
10–12	Standard	166 (58)	1.01 (0.59 to 1.73)	0.91 (0.52 to 1.61)	0.77 (0.42 to 1.40)	0.76 (0.42 to 1.38)
	Group	90 (32)	1.24 (0.70 to 2.12)	1.18 (0.65 to 2.16)	1.09 (0.59 to 2.02)	0.86 (0.46 to 1.63)
	Specialist	29 (30)	Ref	Ref	Ref	Ref
13–20	Standard	53 (61)	0.94 (0.43 to 2.02)	1.14 (0.50 to 2.61)	0.88 (0.37 to 2.09)	0.89 (0.37 to 2.13)
	Group	24 (28)	0.96 (0.41 to 2.22)	1.14 (0.47 to 2.77)	1.04 (0.42 to 2.56)	0.75 (0.29 to 1.94)
	Specialist	10 (11)	Ref	Ref	Ref	Ref
>20 weeks	Standard	53 (68)	1.56 (0.62 to 3.92)	2.19 (0.82 to 5.80)	1.29 (0.46 to 3.60)	1.27 (0.45 to 3.56)
	Group	19 (24)	1.27 (0.46 to 3.45)	1.75 (0.61 to 4.99)	1.29 (0.43 to 3.82)	1.15 (0.38 to 3.44)
	Specialist	6 (8)	Ref	Ref	Ref	Ref

*Model 1: Adjusted for demographics ethnicity, age, parity, IMD score, any social and medical risk factors at booking and onset of labour.

†Model 2: Model 1 + adjustment for place of antenatal care (community or hospital).

‡Model 3: Model 2 + adjustment for service provider attended (A or B).

IMD, indices of multiple deprivation; RR, risk ratio.

Table 5 Gestation at booking appointment in relation to the place of antenatal care

Gestation at booking	Place of antenatal care	Number of women (%)	Unadjusted RR	Model 1 Adjusted RR (95% CI)*	Model 2 Adjusted RR (95% CI)†	Model 3 Adjusted RR (95% CI)‡
<10	Hospital	154 (48)	Ref	Ref	Ref	Ref
	Community	165 (52)	Ref	Ref	Ref	Ref
10–13 weeks	Hospital	154 (54)	1.25 (0.91–1.73)	1.19 (0.85 to 1.66)	1.30 (0.91 to 1.86)	1.00 (0.67 to 1.49)
	Community	131 (46)	Ref	Ref	Ref	Ref
13–20	Hospital	54 (62)	1.75 (1.07–2.84)	1.59 (0.95 to 2.66)	1.66 (0.97 to 2.87)	1.05 (0.54 to 2.01)
	Community	33 (38)	Ref	Ref	Ref	Ref
>20 weeks	Hospital	56 (72)	2.72 (1.58–4.67)	2.89 (1.65 to 5.10)	2.81 (1.56 to 5.06)	2.51 (1.33 to 4.70)
	Community	22 (28)	Ref	Ref	Ref	Ref

*Model 1: Adjusted for demographics ethnicity, age, parity, IMD score, social risk, and medical risk factors at booking and onset of labour.

†Model 2: Model 1 + adjusted for model of care.

‡Model 3: Model 2 + adjusted for service provider attended.

IMD, indices of multiple deprivation; RR, risk ratio.

yet to disclose social risk factors, leading to fragmented care and further risk of ‘falling through the gaps’. The wider Project20 evaluation tests the theory that women will feel more able to disclose risk factors when they have developed a trusting relationship with their care provider (paper currently under review; WOMBI-D-22–00309 R2)

Overall women found the referral to both specialist models of care acceptable, but some expressed a lack of choice, and perceived discrimination about being left in the dark for the reason for referral. There is a wealth of literature around healthcare professionals’ reluctance to address sensitive issues such as mental health, social risk factors and cultural differences with women.^{40–47} In addition to this, socioeconomic status and ethnicity are often associated with stigma and discrimination in healthcare services.^{48–51} These factors might contribute to midwives’ feelings of discomfort around informing women why

they have been referred to a specialist model. Perceived discrimination could be avoided through community-based care within areas of deprivation rather than inclusion criteria based on these social risk factors.

The findings showed no differences between the model of care received and timing of access to maternity care or the number of antenatal appointments women attended. Considering women in the specialist models were more likely to have low socioeconomic status and social risk factors and therefore more likely to struggle to access and engage.^{6 52 53} This indicates that the inequality in access may have been mitigated by the specialist model of care. This theory would need to be tested with a larger sample. Regardless of the model of care women received, those receiving hospital-based care were more likely to attend their first maternity appointment after 20 weeks’ gestation, this was thought to be partly due to convoluted

Table 6 Number of antenatal appointments attended in relation to the model of care accessed

Number of antenatal appointments	Model of care	Number of women (%)	Unadjusted RR	Model 1 Adjusted RR (95% CI)*	Model 2 Adjusted RR (95% CI)†	Model 3 Adjusted RR (95% CI)‡
1–6	Standard	173 (63)	0.58 (0.31–1.09)	0.64 (0.35 to 1.30)	0.86 (0.44 to 1.69)	0.87 (0.44 to 1.70)
	Group	63 (23)	0.41 (0.21–0.80)	0.42 (0.22 to 0.93)	0.43 (0.21 to 0.86)	0.55 (0.25 to 1.15)
	Specialist	37 (14)	Ref	Ref	Ref	Ref
7–9§	Standard	135 (61)	Ref	Ref	Ref	Ref
	Group	70 (31)	Ref	Ref	Ref	Ref
	Specialist	17 (8)	Ref	Ref	Ref	Ref
10–14	Standard	120 (64)	0.94 (0.45–1.95)	1.13 (0.53 to 2.42)	0.80 (0.36 to 1.78)	0.82 (0.37 to 1.81)
	Group	52 (28)	0.78 (0.36–1.70)	0.90 (0.40 to 2.02)	0.82 (0.36 to 1.86)	0.97 (0.42 to 2.24)
	Specialist	16 (8)	Ref	Ref	Ref	Ref
≥15	Standard	38 (46)	0.47 (0.20–1.13)	0.68 (0.26 to 1.74)	0.43 (0.16 to 1.17)	0.43 (0.16 to 1.14)
	Group	35 (42)	0.85 (0.35–2.05)	1.25 (0.47 to 3.28)	1.02 (0.38 to 2.75)	1.28 (0.47 to 3.45)
	Specialist	10 (12)	Ref	Ref	Ref	Ref

*Model 1: Adjusted for demographics ethnicity, age, parity, IMD score, any social and medical risk factors at booking and onset of labour.

†Model 2: Model 1 + adjustment for place of antenatal care (community or hospital).

‡Model 3: Model 2 + adjustment for service provider attended (A or B).

§Set as base as WHO recommends eight antenatal appointments.¹

IMD, indices of multiple deprivation; RR, risk ratio.

Table 7 Number of antenatal appointments attended in relation to the place of antenatal care

Number of antenatal appointments	Model of care	Number of women (%)	Unadjusted RR	Model 1 Adjusted RR (95% CI)*	Model 2 Adjusted RR (95% CI)†	Model 3 Adjusted RR (95% CI)‡
1–6	Hospital	106 (39)	0.53 (0.37–0.77)	0.56 (0.38 to 0.82)	0.47 (0.31 to 0.71)	0.61 (0.38 to 0.99)
	Community	167 (61)	Ref	Ref	Ref	Ref
7–9§	Hospital	120 (54)	Ref	Ref	Ref	Ref
	Community	102 (46)	Ref	Ref	Ref	Ref
10–14	Hospital	130 (69)	1.90 (1.26–2.86)	2.22 (1.44 to 3.41)	2.27 (1.43 to 3.58)	2.70 (1.62 to 4.49)
	Community	58 (31)	Ref	Ref	Ref	Ref
≥15	Hospital	60 (72)	2.21 (1.28–3.83)	2.92 (1.59 to 5.34)	3.36 (1.80 to 6.25)	4.90 (2.50 to 9.61)
	Community	23 (28)	Ref	Ref	Ref	Ref

*Model 1: Adjusted for demographics ethnicity, age, parity, IMD score, social risk, and medical risk factors at booking and onset of labour.

†Model 2: Model 1 + adjusted for model of care.

‡Model 3: Model 2 + adjusted for service provider attended.

§Set as base as WHO recommends eight antenatal appointments.¹

IMD, indices of multiple deprivation; RR, risk ratio.

referral systems between the community and hospital. Many women expressed wanting to be seen earlier in pregnancy, challenging the notion that women with social risk factors do not prioritise their maternity care.⁵⁴ Women also described difficulties in registering or booking an appointment with their GP when they found out they were

pregnant, particularly if they did not speak English, were homeless or unfamiliar with the system. The wider literature has identified similar barriers to access including difficulties navigating the health system and service delays in the processing of referrals.^{52 55 56} A UK study⁵² of women accessing antenatal care in a multicultural,

Table 8 Number of appointments and support in labour by known healthcare professional

Number of antenatal appointments with a known professional	Model of care	Number of women (%)	Unadjusted OR	Model 1 Adjusted RR (95% CI)*	Model 2 Adjusted RR (95% CI)†	Model 3 Adjusted RR (95% CI)‡
None	Standard	313 (69)	Ref	Ref	Ref	Ref
	Group	84 (19)	Ref	Ref	Ref	Ref
	Specialist	53 (12)	Ref	Ref	Ref	Ref
1–2	Standard	108 (73)	6.09 (1.86–19.9)	4.43 (1.30 to 15.0)	2.75 (0.78 to 9.67)	2.12 (0.57 to 7.87)
	Group	36 (24)	7.57 (2.21–25.8)	7.30 (2.05 to 26.0)	6.81 (1.89 to 24.5)	1.81 (0.44 to 7.45)
	Specialist	3 (2)	Ref	Ref	Ref	Ref
3	Standard	36 (47)	0.76 (0.33–1.72)	0.69 (0.27 to 1.77)	0.47 (0.17 to 1.29)	0.34 (0.11 to 1.06)
	Group	32 (42)	2.52 (1.08–5.89)	3.33 (1.27 to 8.75)	3.15 (1.18 to 8.38)	0.82 (0.24 to 2.75)
	Specialist	8 (11)	Ref	Ref	Ref	Ref
4	Standard	7 (15)	0.29 (0.83–1.04)	0.23 (0.05 to 0.89)	0.16 (0.03 to 0.67)	0.10 (0.02 to 0.53)
	Group	34 (76)	5.36 (1.80–15.9)	5.99 (1.78 to 20.1)	5.19 (1.62 to 18.7)	1.72 (0.39 to 1.47)
	Specialist	4 (9)	Ref	Ref	Ref	Ref
>5	Standard	5 (10)	0.76 (0.02–0.23)	0.05 (0.01 to 0.19)	0.03 (0.00 to 0.13)	0.02 (0.00 to 0.11)
	Group	35 (68)	2.00 (0.93–4.29)	2.50 (0.96 to 6.49)	2.38 (0.90 to 6.32)	0.82 (0.23 to 2.94)
	Specialist	11 (22)	Ref	Ref	Ref	Ref
Looked after in labour by a known midwife	Standard	235 (63)	0.41 (0.24–0.71)	0.44 (0.24 to 0.82)	0.62 (0.32 to 1.19)	0.59 (0.30 to 1.17)
	Group	81 (23)	0.24 (0.13–0.42)	0.23 (0.12 to 0.44)	0.24 (0.12 to 0.47)	0.44 (0.21 to 0.92)
	Specialist	53 (14)	Ref	Ref	Ref	Ref

*Model 1: Adjusted for demographics ethnicity, age, parity, IMD score, any social and medical risk factors at booking and onset of labour.

†Model 2: Model 1 + adjustment for place of antenatal care (community or hospital).

‡Model 3: Model 2 + adjustment for service attended (A or B).

IMD, indices of multiple deprivation; RR, risk ratio.

Table 9 Number of appointments with a known healthcare professional in relation to place of care

Number of antenatal appointments with a known professional	Model of care	Number of women (%)	Unadjusted OR	Model 1 Adjusted RR (95% CI)*	Model 2 Adjusted RR (95% CI)†	Model 3 Adjusted RR (95% CI)‡
None	Hospital	215 (48)	Ref	Ref	Ref	Ref
	Community	235 (52)	Ref	Ref	Ref	Ref
1–2	Hospital	109 (74)	3.15 (2.08–4.76)	2.44 (1.56 to 3.83)	2.71 (1.63 to 4.50)	0.24 (0.10 to 0.59)
	Community	38 (26)	Ref	Ref	Ref	Ref
3	Hospital	45 (59)	1.59 (0.97–2.61)	1.23 (0.71 to 2.13)	2.25 (1.19 to 4.25)	0.23 (0.08 to 0.62)
	Community	31 (41)	Ref	Ref	Ref	Ref
4	Hospital	20 (44)	0.87 (0.47–1.62)	0.61 (0.31 to 1.22)	2.02 (0.92 to 1.47)	0.30 (0.11 to 0.83)
	Community	25 (56)	Ref	Ref	Ref	Ref
>5	Hospital	28 (55)	1.33 (0.74–2.39)	0.70 (0.35 to 1.39)	2.66 (1.24 to 5.70)	0.35 (0.12 to 0.96)
	Community	23 (45)	Ref	Ref	Ref	Ref
Looked after in labour by a known midwife	Hospital	153 (41)	0.34 (0.25–0.46)	0.46 (0.33 to 0.64)	0.39 (0.27 to 0.57)	0.89 (0.59 to 1.36)
	Community	216 (59)	Ref	Ref	Ref	Ref

*Model 1: Adjusted for demographics ethnicity, age, parity, IMD score, social risk, and medical risk factors at booking and onset of labour.
 †Model 2: Model 1 + adjusted for model of care.
 ‡Model 3: Model 2 + adjusted for service attended.
 IMD, indices of multiple deprivation; RR, risk ratio.

deprived area found that women want to access care in early pregnancy but perceive antenatal care for viable and continuing pregnancies at a later gestation. If women do not feel that a service is open to them, or that maternity services only value those who carry a viable pregnancy, they may internalise this as a prioritisation of the

well-being of the fetus over their own emotional, physical and social needs. The wider Project20 evaluation previously tested programme theory relating to interpreter services for pregnant women with social risk factors, finding that despite accessing a specialist model of care women experienced a lack of regulation and access to

Table 10 Number of missed appointments in relation to model of care received

Number of missed appointments	Model of care	Number of women (%)	Unadjusted OR	Model 1 Adjusted RR (95% CI)*	Model 2 Adjusted RR (95% CI)†	Model 3 Adjusted RR (95% CI)‡
None	Standard	352 (63)	Ref	Ref	Ref	Ref
	Group	147 (26)	Ref	Ref	Ref	Ref
	Specialist	62 (11)	Ref	Ref	Ref	Ref
1	Standard	62 (56)	1.36 (0.62–2.99)	1.58 (0.67 to 3.67)	1.46 (0.61 to 3.48)	1.44 (0.61 to 3.49)
	Group	40 (37)	2.10 (9.33–4.76)	2.41 (1.01 to 5.77)	2.38 (0.99 to 5.72)	1.97 (0.80 to 4.84)
	Specialist	8 (7)	Ref	Ref	Ref	Ref
2	Standard	37 (67)	1.62 (0.56–4.73)	2.27 (0.72 to 7.13)	1.79 (0.54 to 5.84)	1.81 (0.53 to 6.19)
	Group	14 (26)	1.47 (0.46–4.66)	1.83 (0.54 to 6.20)	1.79 (0.52 to 6.14)	0.86 (0.22 to 3.33)
	Specialist	4 (7)	Ref	Ref	Ref	Ref
3	Standard	14 (52)	1.23 (0.27–5.55)	2.71 (0.33 to 22.0)	2.01 (0.23 to 17.2)	2.02 (0.22 to 18.4)
	Group	11 (41)	2.31 (0.49–10.7)	4.54 (0.54 to 37.9)	4.41 (0.51 to 37.6)	2.03 (0.25 to 23.6)
	Specialist	2 (7)	Ref	Ref	Ref	Ref
≥4	Standard	4 (24)	0.17 (0.42–0.72)	0.20 (0.04 to 0.98)	0.22 (0.04 to 1.14)	0.23 (0.04 to 1.22)
	Group	9 (53)	0.94 (0.28–3.19)	0.94 (0.22 to 4.02)	0.94 (0.21 to 4.05)	0.49 (0.08 to 2.71)
	Specialist	4 (23)	Ref	Ref	Ref	Ref

*Model 1: Adjusted for demographics ethnicity, age, parity, IMD score, any social and medical risk factors at booking and onset of labour.
 †Model 2: Model 1 + adjustment for place of antenatal care (community or hospital).
 ‡Model 3: Model 2 + adjustment for service provider attended (A or B).
 IMD, indices of multiple deprivation; RR, risk ratio.

Table 11 Number of missed appointments in relation to place of care

Number of missed appointments	Place of care	Number of women (%)	Unadjusted OR	Model 1 Adjusted RR (95% CI)*	Model 2 Adjusted RR (95% CI)†	Model 3 Adjusted RR (95% CI)‡
None	Hospital	292 (52)	Ref	Ref	Ref	Ref
	Community	269 (48)	Ref	Ref	Ref	Ref
1	Hospital	64 (58)	1.28 (0.84–1.93)	1.11 (0.71 to 1.72)	1.21 (0.76 to 1.94)	0.95 (0.56 to 1.62)
	Community	46 (42)	Ref	Ref	Ref	Ref
2	Hospital	38 (69)	2.06 (1.13–3.74)	2.07 (1.12 to 3.88)	1.96 (1.01 to 3.78)	0.64 (0.26 to 1.58)
	Community	17 (31)	Ref	Ref	Ref	Ref
3	Hospital	18 (67)	1.84 (0.81–4.17)	1.82 (0.74 to 4.46)	2.18 (0.83 to 5.67)	1.02 (0.32 to 3.20)
	Community	9 (33)	Ref	Ref	Ref	Ref
≥4	Hospital	6 (35)	0.50 (0.18–1.38)	0.48 (0.15 to 1.48)	0.77 (0.23 to 2.51)	0.36 (0.08 to 1.55)
	Community	11 (65)	Ref	Ref	Ref	Ref

*Model 1: Adjusted for demographics ethnicity, age, parity, IMD score, social risk, and medical risk factors at booking and onset of labour.

†Model 2: Model 1 + adjusted for model of care.

‡Model 3: Model 2 + adjusted for service provider attended.

IMD, indices of multiple deprivation; RR, risk ratio.

high-quality interpretation services.⁵⁷ As well as impacting on women's engagement with services, these factors can impact on the safety of women with complex needs such as those who are experiencing abuse, poor mental health or need to discuss a termination of pregnancy. Previous research has found a correlation between continuity models of care and increased disclosure and referral to support services,^{58 59} but it must be acknowledged that under ascertainment of sensitive issues such as mental health and domestic abuse remains likely while women perceive services as a form of surveillance and risk.⁵⁹

A recent review of the literature on how women with social risk factors experience maternity care in the UK found reasons for late access include the denial of services based on a lack of documentation, fear of disclosure to immigration services, language and financial barriers, cultural differences, unfamiliarity, a lack of trust and a perception that maternity services act as a system of surveillance rather than support.⁷ Practice recommendations

detailed in the refined programme theory suggest that all women are made aware of the possibility to self-refer directly to maternity services at the first point of contact with health services, using language appropriate information. Strict inclusion criteria for access to specialist models that restrict women who have booked late for maternity care should be reviewed as these women are often high risk and can benefit from the specialist model regardless of their gestation. An evaluation of group antenatal care⁶⁰ included women who had booked later in pregnancy with positive feedback and no negative impact on group dynamics. With the national expansion of continuity of care models,¹⁸ inclusion criteria could be relaxed somewhat as the demands on the service are more evenly distributed. Future research should assess the impact of place-based midwifery continuity of care on health inequalities to test the outcomes of this preference.

This study reflects findings in the wider literature^{6 61} highlighting service use inequities for women

Table 12 Overview of quantitative outcomes

Access and engagement outcome variable	Characteristics of women at disproportionate risk when adjusting (online supplemental file 2)	Significant effect of specialist model of care	Significant effect of hospital-based antenatal care	Significant effect of service
Access to specialist model	Most deprived, social risk factors, black African ethnicity	↑	↓	=
Late gestation at booking appointment (>20/40)	Primiparous, high medical risk, social risk factors	=	↑	=
Number of antenatal appointments outside of recommendations ^{1 2}	Social risk factors and high medical risk status (>15 appointments)	=	↓/↑	B↓
Appointments with a known healthcare professional	Black African (least likely to see a known healthcare professional)	=	↓	B↓
Missed appointments	Multiparous, black African, social risk	=	=	B↑

↑=statistically significant increase (p<0.05), ↓=statistically significant decrease (p<0.05), =no significant relationship detected, 'A' and 'B' refer to service providers.

Table 13 Refined programme theory—access and engagement with maternity services

Context +	Mechanism =	Outcome
Women who struggle to access services and are at greater risk of booking for maternity care at a late gestation. Women who are unfamiliar with the UK health system or have difficulties in registering with health services. These women are often experiencing social risk factors that might lead to chaotic lives, social isolation, lack of resource, lack of support. Primiparous women, those with any social risk factor, and high medical risk are more likely to book late for maternity care. Multiparous women, black African women and those with social risk factors are more likely to miss appointments. Black African women have less appointments with a known healthcare professional.	(M1) If maternity care provision commences when a woman accesses services regardless of her gestation and women have 24/7 access to a small team of midwives whom they have had the opportunity to meet during pregnancy and are encouraged to contact via a phone call, text message or free technology.	(O1) Then women would not feel unsupported, anxious and that the service does not value them until they have a viable pregnancy. This might also improve early access to safe abortion and family planning services. Engagement with services will improve through needs-based communication and appropriately timed antenatal appointments. This open access can work both ways through midwives reminding them of appointments, this leads to women feeling more 'cared for'.
	(M2) If women are made aware of the possibility and how to self-refer to maternity services and specialist models of care by administrative staff at the first point of contact.	(O2) Then difficulties trying to access a GP will be overcome, the time spent waiting for a GP appointment reduced and long referral processes between primary and secondary services will be avoided.
	(M3) If women can access a community-based service where GP's and midwives regularly communicate with each other.	(O3) Then the timing of access to a booking appointment with a midwife will be improved and convoluted referral pathways between community and hospital services avoided.
	(M4) If women living in areas of deprivation are prioritised to receive continuity of care through community-based models.	(O4) Then services are likely to provide an enhanced package of care to women with social risk factors who have not previously disclosed these issues with professionals, and care is less likely to be disrupted during pregnancy when a disclosure is made.
	(M5) If women are informed of the reasons why they have been allocated a continuity of care model or specialist service, or are able to self-refer to them if they feel they are eligible for their care.	(O5) Then the development of a trusting and open relationship with their healthcare provider will be enabled and feelings of suspicion and surveillance reduced. This transparency may also reduce feelings of discrimination for women in marginalised groups.
	(M6) If women who book late for pregnancy care are eligible for specialist models of care where they have a named midwife or small team of midwives.	(O6) Then the benefits of these models of care and highlighted mechanisms may protect them from the disproportionately poor outcomes associated with late booking.
	(M7) If women can reschedule appointments easily, and do not fear judgement or reproach when they miss appointments.	(O7) Then they will perceive the maternity environment as a place of safety and their engagement with flexible services will improve.
	(M8) If women have the opportunity to get to know all the midwives in a small team throughout their pregnancy and the number, time and place of appointments is co-planned with women to meet their individual needs.	(O8) Then they will not feel disappointment or let down when their named midwife is unable to attend an appointment. Care, information and responsibility will be shared across the team, women will be better able to engage, not perceive the pressure of time and feel more able to seek help and disclose information thus improving safety.

with characteristics such as multiparity, age and ethnicity and presents an opportunity for those designing models of care to focus on these demographics. Where poor engagement with services is often associated with women's priorities and behaviours,^{7 52 62} the women in this study highlighted system barriers. Interestingly, many women in this study felt that the ability to contact a known midwife anytime reduced the number of face-to-face appointments they needed. Where engagement in this study was measured through the number of antenatal appointments attended, it would be useful for future research to measure other forms of contact between women and healthcare professionals. Women from the hospital-based specialist model discussed the detrimental impact of midwives needing to cancel their appointments. Rather than a place-based issue this seems to relate to how the teams are organised as those in the hospital-based model were allocated one midwife who they saw for most of their

appointments, whereas women accessing the community-based model described being cared for by the whole team. Finally, women from both community and hospital-based models expressed a preference for care to be based in the community or home setting as they felt it was more accessible and supportive of their needs.

The aim of both the group practice and specialist models evaluated appear to be being met with women more likely to receive more antenatal appointments with a known healthcare professional from these models. Women who received hospital-based antenatal care had significantly lower levels of continuity in the antenatal period. Women receiving care in the specialist model were more likely to be looked after in labour by a known healthcare professional compared with the other models. The qualitative data revealed this continuity of care lessened anxiety, the need to repeat often complex social and medical histories to numerous professionals and increased disclosure

of social risk factors, and preparedness for labour and birth, particularly for those who had the opportunity to meet the rest of the team. Multiparous, younger and black African women were more likely to miss antenatal appointments, as well as those with social risk factors. Black African women were also less likely to see a known healthcare professional than women of other ethnicities, perhaps reflecting an underlying mechanism of the health inequalities they experience.⁶³ There is a significant paucity of literature around black women's experiences of UK maternity care, but a US review⁶⁴ concluded that midwifery care can accommodate their specific needs through attentive provider, continuity of care and empowerment. The review stated that 'researchers must meet, consult with and listen to black women and hear their stories to understand the significance of midwifery to them. Only then, in partnership with, and based on suggestions for change from black women themselves, can health care providers and researchers begin to make changes in the health care system to facilitate improved antenatal care'.⁶⁴ This recommendation is as relevant to the UK context given the known stark inequalities.

Strengths and limitations

Women interviewed during the study may have perceived the study questions to be testing them about their willingness to engage with their care. This limitation may have been lessened through the trust built between the participant and the researcher over the course of the longitudinal interviews.^{65 66} The small and varied numbers in each quantitative data group should be taken into consideration due to the significant amount of multiple testing required to establish the separate effects of the model of care, place of care and service attended. This presents a potential limitation as the use of multiple testing can result in erroneous inferences, reducing the probability of detecting effects when they do exist.⁶⁷ This could be overcome in future research using larger sample sizes, and Bayesian analysis to test the apparent mitigating effects of the specialist models of care on inequalities in access and engagement.⁶⁸ That said, the claim to the causality of the mechanisms is strengthened as both qualitative and quantitative data analysis point to a causal link between the mechanism(s) and the outcomes studied.⁶⁸

The analysis of quantitative data is also limited by the availability and depth of information routinely recorded in maternity services. It is recommended that more granular ethnicity categories, using the Office for National Statistics 18+ guidance⁶⁹ is used, as well as migration details such as country of birth, national identity, length of time in the UK, visa conditions and language proficiency. Finally, the generalisability of the findings is limited by the urban location of both specialist models of care evaluated and the UK's health system context. This is particularly significant when reflecting on the outcomes relating to place-based care what may have yielded significant outcomes in a densely populated, multicultural community, may yield very different results elsewhere. Research

is needed to test the generalisability of the findings to rural and other community settings.

CONCLUSION

This research highlights how carefully considered place-based care with a focus on continuity of carer can improve access and engagement with maternity services for women with social risk factors, but it is not a panacea. Hospital-based models of care with strict inclusion criteria may risk excluding women at increased risk who are yet to disclose social risk factors. Rather than the often-assumed 'relaxed maternal care-seeking behaviour' and 'women with social risk factors deprioritising maternity care' explanations for late booking and missed appointments, the findings highlight system barriers. Women want to be able to access care earlier in pregnancy, and there appears to be a lack of information regarding their choices in doing so, particularly for those who do not speak English or are unfamiliar with the system. The identification of specific mechanisms that improve access and engagement with services are outlined in the refined programme theory, which will enable those developing maternity services to structure models of care around women's individual needs.

Twitter Hannah Rayment-Jones @HannahTLGG

Acknowledgements The authors would like to thank student midwives Laura Peazold, Mary Newman, Natalie Goodyear and Micaela Anthony for help with data collection and anonymisation, Justin Jagosh for invaluable advice on realist methodology and Sergio Silverio for support with qualitative data analysis.

Contributors HR-J, JMH, AH and JS contributed to the conceptualisation of the research question and methodology. EP and TG organised collection, anonymisation and analysis of the quantitative data. HR-J and KD analysed the quantitative data. HR-J collected qualitative data and analysed it with JMH. All authors interpreted the data analysis, read and approved the final manuscript. HR-J is the acting guarantor.

Funding This report is an independent research supported by the National Institute for Health Research (NIHR Doctoral Research Fellowship, HR-J, award no DRF-2017-10-033). AH is supported by the NIHR Applied Research Collaboration North Thames. JS (King's College London) is supported by the NIHR Applied Research Collaboration South London (NIHR ARC South London). JS is also a NIHR Senior Investigator. JMH is supported by a Post-doctoral Fellowship from Wellbeing of Women (Award Ref PRF006). KD is funded by the Medical Research Council (MRC) (grant number: MR/V005839/1). The views expressed in this article are those of the author(s) and not necessarily those of the National Health Service, the NIHR, the MRC or the Department of Health and Social Care.

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Ethics approval This research was approved by the London Brent Research Ethics Committee (HRA) REC Reference 18-L0-0701. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Full quantitative data analysis attached as additional file or contact the lead author HR-J for qualitative and raw data files that are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and

responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Hannah Rayment-Jones <http://orcid.org/0000-0002-3027-8025>

Kathryn Dalrymple <http://orcid.org/0000-0003-0958-6725>

REFERENCES

- World Health Organization (WHO). WHO recommendations on antenatal care for a positive pregnancy experience: summary; 2018.
- National Institute for Clinical Excellence (NICE). Antenatal care. quality standard [QS22]; 2012.
- Nam JY, Cho E, Park EC. Do severe maternal morbidity and adequate prenatal care affect the delivery cost? A nationwide cohort study for 11 years with follow up. *BJOG* 2019;126:1623–31.
- Renfrew MJ, McFadden A, Bastos MH, *et al*. Midwifery and quality care: findings from a new evidence-informed framework for maternal and newborn care. *Lancet* 2014;384:1129–45.
- Shaw D, Guise J-M, Shah N, *et al*. Drivers of maternity care in high-income countries: can health systems support woman-centred care? *Lancet* 2016;388:2282–95.
- Lindquist A, Kurinczuk JJ, Redshaw M, *et al*. Experiences, utilisation and outcomes of maternity care in England among women from different socio-economic groups: findings from the 2010 national maternity survey. *BJOG* 2015;122:1610–7.
- Rayment-Jones H, Harris J, Harden A, *et al*. How do women with social risk factors experience United Kingdom maternity care? A realist synthesis. *Birth* 2019;46:461–74.
- Downe S, Finlayson K, Tunçalp Ö, *et al*. What matters to women: a systematic scoping review to identify the processes and outcomes of antenatal care provision that are important to healthy pregnant women. *BJOG* 2016;123:529–39.
- Finlayson K, Downe S. Why do women not use antenatal services in low- and middle-income countries? A meta-synthesis of qualitative studies. *PLoS Med* 2013;10:e1001373.
- Arsenault C, Jordan K, Lee D, *et al*. Equity in antenatal care quality: an analysis of 91 national household surveys. *Lancet Glob Health* 2018;6:e1186–95.
- Nair M, Knight M, Kurinczuk JJ. Risk factors and newborn outcomes associated with maternal deaths in the UK from 2009 to 2013: a national case-control study. *BJOG* 2016;123:1654–62.
- Amjad S, Chandra S, Osornio-Vargas A, *et al*. Maternal area of residence, socioeconomic status, and risk of adverse maternal and birth outcomes in adolescent mothers. *J Obstet Gynaecol Can* 2019;41:1752–9.
- Knight M, Bunch K, Kenyon S, *et al*. Saving lives, improving mothers' care. Lessons learned to inform maternity care from the UK and Ireland confidential enquiries into maternal deaths and morbidity 2015–17; 2019.
- Ki-Moon B. Global strategy for women's and children's health. In: *Maternal Mortality, Human Rights and Accountability*. 2013.
- N C C for W and C H Pregnancy and Complex Social Factors. Pregnancy and complex social factors: a model for service provision for pregnant women with complex social factors. UK; 2010.
- Organization, world health. WHO recommendation on antenatal care for positive pregnancy experience. WHO recommendation on antenatal care for positive pregnancy experience; 2016, ISBN 978 92 4 154991 2.
- Kennedy HP, Yoshida S, Costello A. Asking different questions: research priorities to improve the quality of care for every woman, every child. *Lancet Glob Health* 2016;4:e777–9.
- NHS England. The NHS long term plan: maternity and neonatal services. 2019.
- National Maternity Review. Better births: improving outcomes of maternity services in England; 2016.
- Sandall J, Soltani H, Gates S, *et al*. Midwife-led continuity models versus other models of care for childbearing women. *Cochrane Database Syst Rev* 2016;4:CD004667.
- Hollowell J, Kurinczuk JJ, Oakley L, *et al*. A systematic review of the effectiveness of antenatal care programmes to reduce infant mortality and its major causes in socially disadvantaged and vulnerable women. National Perinatal Epidemiology Unit (NPEU); 2009.
- Pawson R. *The science of evaluation: A realist manifesto*. 2014.
- Jagosh J. Realist synthesis for public health: building an ontologically deep understanding of how programs work, for whom, and in which contexts. *Annu Rev Public Health* 2019;40:361–72.
- Rayment-Jones H, Silverio SA, Harris J, *et al*. Project 20: midwives' insight into continuity of care models for women with social risk factors: what works, for whom, in what circumstances, and how. *Midwifery* 2020;84:102654.
- Jagosh J. Retroductive theorizing in pawson and tilley's applied scientific realism. *J Crit Realism* 2020;19:121–30.
- Mukumbang FC, Kabongo EM, Eastwood JG. Examining the application of retroductive theorizing in realist-informed studies. *Int J Qual Methods* 2021;20:160940692110535.
- Ogrinc G, Davies L, Goodman D, *et al*. Squire 2.0 (standards for quality improvement reporting excellence): revised publication guidelines from a detailed consensus process. *BMJ Qual Saf* 2016;25:986–92.
- Fernandez Turienzo C, Newburn M, Agyepong A, *et al*. Addressing inequities in maternal health among women living in communities of social disadvantage and ethnic diversity. *BMC Public Health* 2021;21:176.
- Murray SF, Buller AM, Bewley S, *et al*. Metrics for monitoring local inequalities in access to maternity care: developing a basket of markers from routinely available data. *Qual Saf Health Care* 2010;19:e39.
- Ministry of Housing, C. & L. G. English indices of deprivation 2019. National Statistics; 2019.
- Office for National Statistics. The english indices of deprivation 2015. Neighbourhoods Statistical Release; 2015. Available: <http://www.communities.gov.uk/publications/corporate/statistics/indices2010technicalreport>
- Manzano A. The craft of interviewing in realist evaluation. *Evaluation* 2016;22:342–60.
- Ministry of Housing Communities and Local Government. English indices of deprivation 2019. National Statistics; 2019.
- du Prel J-B, Hommel G, Röhrig B, *et al*. Confidence interval or P-value?: part 4 of a series on evaluation of scientific publications. *Dtsch Arztebl Int* 2009;106:335–9.
- Gale NK, Heath G, Cameron E, *et al*. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol* 2013;13:117.
- Vasileiou K, Barnett J, Thorpe S, *et al*. Characterising and justifying sample size sufficiency in interview-based studies: systematic analysis of qualitative health research over a 15-year period. *BMC Med Res Methodol* 2018;18:148.
- Kirkham M, Stapleton H, Curtis P, *et al*. The inverse care law in antenatal midwifery care. *British Journal of Midwifery* 2002;10:509–13.
- Tudor Hart J. The inverse care law. *The Lancet* 1971;297:405–12.
- Charles A, Ewbank L, McKenna H, *et al*. The NHS long-term plan explained. Long read. 2019.
- Fontein-Kuipers YJ, Budé L, Ausems M, *et al*. Dutch midwives' behavioural intentions of antenatal management of maternal distress and factors influencing these intentions: an exploratory survey. *Midwifery* 2014;30:234–41.
- Everitt L, Homer C, Fenwick J. Working with vulnerable pregnant women who are at risk of having their babies removed by the child protection agency in new south wales, australia. *Child Abuse Rev* 2017;26:351–63.
- Higgins A, Carroll M, Downes C, *et al*. *Perinatal mental health: an exploration of practices, policies, processes and education needs of nurses and midwives within maternity and primary care services in Ireland*. Dublin: Health Services Executive, 2017.
- Baldwin S, Griffiths P. Do specialist community public health nurses assess risk factors for depression, suicide, and self-harm among South Asian mothers living in London? *Public Health Nurs* 2009;26:277–89.
- Jomeen J, Glover L, Jones C, *et al*. Assessing women's perinatal psychological health: exploring the experiences of health visitors. *J Reprod Infant Psychol* 2013;31:479–89.
- Jane R. Midwives, society and childbirth: debates and controversies in the modern period. *Women's Hist Rev* 1999;8:169–88.
- Mulherin K, Miller YD, Barlow FK, *et al*. Weight stigma in maternity care: women's experiences and care providers' attitudes. *BMC Pregnancy Childbirth* 2013;13:19.
- Radcliffe P. Substance-misusing women: stigma in the maternity setting. *Br J Midwifery* 2011;19:497–506.



- 48 Stepanikova I, Oates GR. Perceived discrimination and privilege in health care: the role of socioeconomic status and race. *Am J Prev Med* 2017;52:S86–94.
- 49 Tajeu GS, Cherrington AL, Andreae L, et al. “We’ll get to you when we get to you”: exploring potential contributions of health care staff behaviors to patient perceptions of discrimination and satisfaction. *Am J Public Health* 2015;105:2076–82.
- 50 Phelan JC, Lucas JW, Ridgeway CL, et al. Stigma, status, and population health. *Soc Sci Med* 2014;103:15–23.
- 51 Walker R. *The shame of poverty*. 2014.
- 52 Hatherall B, Morris J, Jamal F, et al. Timing of the initiation of antenatal care: an exploratory qualitative study of women and service providers in east london. *Midwifery* 2016;36:1–7.
- 53 Rowe RE, Magee H, Quigley MA, et al. Social and ethnic differences in attendance for antenatal care in England. *Public Health* 2008;122:1363–72.
- 54 Haddrill R, Jones GL, Mitchell CA, et al. Understanding delayed access to antenatal care: a qualitative interview study. *BMC Pregnancy Childbirth* 2014;14:207.
- 55 Hollowell J, Oakley C, Vigurs E, et al. Increasing the early initiation of antenatal care by black and minority ethnic women in the united kingdom: a systematic review and mixed methods synthesis of women’s views and the literature on intervention effectiveness. Social Science Research Unit, Institute of Education, University of London; 2012.
- 56 Cresswell JA, Yu G, Hatherall B, et al. Predictors of the timing of initiation of antenatal care in an ethnically diverse urban cohort in the UK. *BMC Pregnancy Childbirth* 2013;13:103.
- 57 Rayment-Jones H, Harris J, Harden A, et al. Project20: interpreter services for pregnant women with social risk factors in england: what works, for whom, in what circumstances, and how? *Int J Equity Health* 2021;20:233.
- 58 Rayment-Jones H, Murrells T, Sandall J. An investigation of the relationship between the caseload model of midwifery for socially disadvantaged women and childbirth outcomes using routine data—a retrospective, observational study. *Midwifery* 2015;31:409–17.
- 59 Rayment-Jones H, Harris J, Harden A, et al. Project20: maternity care mechanisms that improve (or exacerbate) health inequalities. A realist evaluation. *Women Birth* 2022.
- 60 Hunter LJ, Da Motta G, McCourt C, et al. Better together: a qualitative exploration of women’s perceptions and experiences of group antenatal care. *Women Birth* 2019;32:336–45.
- 61 Higginbottom GMA, Evans C, Morgan M, et al. Experience of and access to maternity care in the UK by immigrant women: a narrative synthesis systematic review. *BMJ Open* 2019;9:e029478.
- 62 Docherty A, Bugge C, Watterson A. Engagement: an indicator of difference in the perceptions of antenatal care for pregnant women from diverse socioeconomic backgrounds. *Health Expect* 2012;15:126–38.
- 63 MBRRACE-UK. Saving lives, improving mother’s care- lay summary. action against medical accidents); 2018.
- 64 Yoder H, Hardy LR. Midwifery and antenatal care for black women: a narrative review. *SAGE Open* 2018;8:215824401775222.
- 65 Grinyer A, Thomas C. The value of interviewing on multiple occasions or longitudinally. In: *The SAGE Handbook of Interview Research: The Complexity of the Craft*. 2012.
- 66 Bhopal K, Deuchar R. *Researching marginalized groups*. 2015.
- 67 Miller R. Multiple comparisons-I. In: *Encyclopedia of Statistical Sciences*. 2006.
- 68 Ravn R. Testing mechanisms in large-N realistic evaluations. *Evaluation* 2019;25:171–88.
- 69 Office for National Statistics. Ethnic group, national identity and religion. n.d. Available: <https://www.ons.gov.uk/methodology/classificationsandstandards/measuringequality/ethnicgroupnationalidentityandreligion>