



Correction to: Projected ocean acidification and seasonal temperature alter the behaviour and growth of a range extending tropical fish

Angus Mitchell¹ · Chloe Hayes¹ · David J. Booth² · Ivan Nagelkerken¹

© The Author(s) 2023

Correction to: Coral Reefs (2023) 42:919–929
<https://doi.org/10.1007/s00338-023-02396-2>

The original article has been corrected: The ‘Materials and Methods’ section was incomplete. The information added is:

"On days 6, 17, and 25 of the experiment, all tropical and temperate fish underwent lateralization trials (reported in Mitchell et al. 2022a). Directly following our behavioural video observations on days 19 and 27 in the present study, all fish also underwent a brief startle escape response test through release of a ceramic weight from above the water surface (Mitchell et al. 2022a). We assumed the lateralization trials had no significant effect on our behavioural observations because a 24-hr rest period is adequate in alleviating potential acute stress responses in fishes (e.g., handling; Pankhurst, 2011). The API Stress Coat™ that was added to tanks daily has also been shown to reduce stress responses in fishes due to handling and transport (Vanderzwalmen et al., 2019)."

Mitchell A, Booth DJ, Nagelkerken I (2022a) Ocean warming and acidification degrade shoaling performance and lateralisation of novel tropical–temperate fish shoals. *Glob Change Biol* 28(4):1388–1401. <https://doi.org/10.1111/gcb.16022>

Pankhurst NW (2011) The endocrinology of stress in fish: an environmental perspective. *Gen Comp Endocrinol* 170(2):265–275. <https://doi.org/10.1016/j.ygcen.2010.07.017>

Vanderzwalmen M, Eaton L, Mullen C, Henriquez F, Carey P, Snellgrove D, Sloman, KA (2019) The use of feed and water additives for live fish transport. *Rev Aquac* 11(1):263–278. <https://doi.org/10.1111/raq.12239>

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

The original article can be found online at <https://doi.org/10.1007/s00338-023-02396-2>.

✉ Ivan Nagelkerken
ivan.nagelkerken@adelaide.edu.au

¹ Southern Seas Ecology Laboratories, School of Biological Sciences, The University of Adelaide, DX 650 418, Adelaide, SA 5005, Australia

² School of the Life Sciences, University of Technology Sydney, Ultimo, NSW 2007, Australia

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.