

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

Cost of illness and health-related quality of life for stuttering: Two systematic reviews

Alicia Norman <sup>a, b</sup>, Robyn Lowe <sup>c</sup>, Mark Onslow <sup>c</sup>, Sue O’Brian <sup>c</sup>, Ann Packman <sup>c</sup>, Ross  
Menzies <sup>c</sup>, Liz Schroeder <sup>a, d</sup>

<sup>a</sup> Macquarie University Centre for the Health Economy, Macquarie University Business  
School, NSW, Australia.

<sup>b</sup> Australian Institute for Health Innovation, NSW, Australia.

<sup>c</sup> University of Technology Sydney, Australian Stuttering Research Centre, NSW, Australia

<sup>d</sup> Present address: University of Oxford, Nuffield Department of Primary Care Health  
Sciences, Oxford, UK

\*Corresponding author: Alicia Norman, Macquarie University Centre for the Health  
Economy, Macquarie University Business School, and Australian Institute for Health  
Innovation. Level 5, 75 Talavera Road, North Ryde, NSW, 2113, AUSTRALIA.  
e-mail: alicia.norman@mq.edu.au

*Keywords:* stuttering, adult, quality of life, cost of illness, systematic review

26

**Abstract**

27 *Background:* For those who stutter, verbal communication is typically compromised in social  
28 situations. This may attract negative responses from listeners and stigmatization by society.  
29 These have the potential to impair health-related quality of life across a range of domains,  
30 including qualitative and quantitative impacts on speech output, mental health issues, and  
31 failure to attain educational and occupational potential. These systematic reviews were  
32 designed to explore this matter using traditional health economics perspectives of utility  
33 measures and cost of illness.

34 *Methods:* Studies were included if they involved children, adolescents, or adults with  
35 stuttering as a primary diagnosis. The quality of life search strategy identified 2,607 reports,  
36 of which three were included in the quality of life analysis. The cost of illness search strategy  
37 identified 3,778 reports, of which 39 were included in the cost of illness analysis.

38 *Results:* Two of the three studies included in the quality of life analysis had a high risk of  
39 bias. When measured using utility scores, quality of life for people who stutter was in the  
40 range of those reported for chronic health conditions such as diabetes mellitus, cardiovascular  
41 disease, and cancer. However, there is little such evidence of quality of life impairment  
42 during the preschool years. Studies included in the cost of illness analysis carried  
43 considerable risk of bias overall.

44 *Conclusions:* For people who stutter, there are substantive direct and indirect costs of illness.  
45 These include impairment, challenges, and distress across many domains throughout life,  
46 including income, education, employment, and social functioning. Evidence of quality of life  
47 impairment using utility measures is extremely limited. If this situation is not remedied, the  
48 lifetime impairment, challenges, and distress experienced by those who stutter cannot be  
49 documented in a form that can be used to influence health policy and healthcare spending.

50

**Introduction****51 Health-related quality of life and health economic evaluation**

52 Health economics relates decisions about the use of health care to the costs and outcomes  
53 of that health care. A key consideration within health economics is health-related quality of  
54 life, which is a multidimensional concept that extends beyond clinical measures of health,  
55 such as diagnosis and physiological function, to include physical and social functionality,  
56 energy level, and mental and physical wellbeing (Wilson & Cleary, 1995). Health-related  
57 quality of life includes interaction between those variables, individual characteristics, and the  
58 environment in which the individual functions.

59 All health issues impact not just the individuals and families who experience them, but  
60 also health care systems and the societies they serve. Health economic evaluation informs  
61 what is therefore a “range of very different but unavoidable decisions in health care”  
62 (Drummond et al., 2015, p. 3). Government and private health care funders, and health care  
63 policy makers, need to compare different interventions for different diseases and make  
64 decisions about where to direct health care spending to maximize benefits. This can be done  
65 with overarching measures that assess health-related quality of life across many diseases and  
66 support health economic evaluations.

67 Utility values are fundamental data for conducting health economic evaluations, being a  
68 means to measure health-related quality of life across different diseases. Utility values  
69 represent an individual’s preference for different health states, traditionally measured on a  
70 scale between 0 and 1, spanning the extremes of death and perfect health (Drummond et al.,  
71 2015). The other fundamental datum for health economics is cost of illness, which is a  
72 monetary measure of the burden of an illness to society. The costs of such a burden involve  
73 the fundamental expense to the individual in terms of receiving treatment and lost work  
74 hours, as well as wider societal costs.

75 Economic evaluations of healthcare interventions can be conducted through cost-utility  
76 analysis. The summary outcome of a cost-utility analysis is cost per quality-adjusted life year.  
77 A quality-adjusted life year is obtained by multiplying a utility value by the number of years  
78 lived in a certain health state. This provides an index that combines quality of life and length  
79 of life obtained after treatment. One quality-adjusted life year is equivalent to one year lived  
80 in perfect health. Cost-utility analyses allow health care funders and policy makers to  
81 compare different interventions for the same disease, and for different diseases, using one  
82 summary outcome. This facilitates maximum health benefit from healthcare spending.

### 83 **Health-related quality of life and stuttering**

84 Stuttering is a prevalent and potentially lifelong disorder, with a lifetime incidence in the  
85 range of 8–10% (Bloodstein et al., 2021; Yairi & Ambrose, 2013). While around 10% of  
86 preschool children start to stutter, most recover naturally within a few years of onset, leaving  
87 around 1% to stutter through childhood, adolescence, and adulthood. Verbal communication  
88 is compromised for many people who stutter, and this has the potential to impact health-  
89 related quality of life. This is reflected in how commonly treatment is sought. A study by  
90 Boyce et al. (2022) reported that 73% of adults who stuttered received treatment from a  
91 speech-language-pathologist. In that report, 92% of parents indicated that their children  
92 (mean age 11 years) had received treatment. The following sections give an overview of the  
93 many features of the disorder that are potential sources of such impact.

### 94 **Qualitative and quantitative aspects of speech output**

95 Fundamentally, the speaker experiences disruptive speech behaviors. These have been  
96 categorized as various types of repeated movements, fixed postures (“blocks”) with or  
97 without audible airflow, and verbal or nonverbal superfluous behaviors (Teesson et al., 2003).  
98 Those disruptive speech behaviors, more often than not, occur together during stuttering  
99 moments (O’Brian et al., 2022). Based on a 12-hour speaking day, a report found a mean

100 number of 33,617 syllables spoken for adults with a mean 7.8 percent syllables stuttered  
101 (Karimi et al., 2013a). Those data suggest that participants stuttered around 2–3 thousand  
102 times per day. Because stuttering moments occur so frequently, and each of them is time  
103 consuming, adults who stutter take, on average, three times as long as others to convey their  
104 message (Johnson, 1961; Spencer et al., 2009).

105 It seems that the complexity of stuttering moments increases during life, with repeated  
106 movements being predominant at onset during the preschool years (Ambrose & Yairi, 1999;  
107 Reilly et al., 2009; Yairi & Lewis, 1984) and with evidence of their advancing behavioral  
108 complexity from adolescence to adulthood (O’Brian et al., 2022). Part of stuttering causality  
109 appears to involve anomalies of brain structure and function in areas subserving spoken  
110 language (Chang et al., 2018; Packman, 2012). Perhaps for that reason, the broad notion “loss  
111 of control” has been considered as fundamental to the disorder (Perkins, 1983), and around  
112 half of adults who stutter report a loss of control while speaking, “either often or always”  
113 (Tichenor & Yaruss, 2019, p. 4339).

114 The experience of anticipating stuttering during speech is common (Jackson et al., 2015;  
115 Johnson & Solomon, 1937; Martin & Haroldson, 1967; Milisen, 1938). Word avoidance and  
116 circumlocution are common responses to anticipation (Crichton-Smith, 2002; Jackson et al.,  
117 2015; Martens & Engel, 1986; Vanryckeghem et al., 2004). According to analyses using  
118 Systemic Functional Linguistics, adults who stutter have been shown to consciously change  
119 their use of language in order to limit conversational interaction. (Lee et al., 2015; Spencer et  
120 al., 2005, 2009). Systemic Functional Linguistics reflects how people modulate language in  
121 different contexts and situations. There is also some evidence of that effect during childhood  
122 (Weiss & Zebrowski, 1994), but there is no evidence that children who stutter have reduced  
123 language ability per se (Nippold, 2019). Constant effort to conceal stuttering with word  
124 avoidance and circumlocution, and also situation avoidance, occurs commonly (Boyle &

125 Gabel, 2020; Douglass et al., 2018) and has been labelled “covert stuttering” (Murphy et al.,  
126 2007) or “interiorized stuttering” (Sønsterud et al., 2022).

### 127 **Stuttering variability**

128 Stuttering severity varies across individuals, from mild to severe, and also within  
129 individuals. Survey responses of 204 adults (Tichenor & Yaruss, 2021) indicated that 97% of  
130 them experienced variability of their stuttering across time and situations. That result is  
131 consistent with a control chart study of stuttering variability during a day for 10 adults  
132 (Karimi et al., 2013b), which showed that half of them had stuttering that varied more than  
133 three standard deviations from their mean daily score. The majority of the Tichenor and  
134 Yaruss (2021) participants attributed frustration with the disorder to stuttering variability  
135 across times and situations. Stuttering severity will vary according to the nature of an  
136 audience, particularly its size (Porter, 1939; Siegel & Haugen, 1964; Steer & Johnson, 1936).  
137 Stuttering severity can vary from situation to situation (Ulliana & Ingham, 1984) and can  
138 vary when the speaker is in the same situation (Constantino et al., 2016).

### 139 **Social anxiety**

140 Stuttering moments may invoke social penalty by drawing attention to speech and  
141 interfering with communication. Accordingly, the disorder is commonly associated with  
142 mental health issues associated with social anxiety. A large body of evidence confirms this to  
143 be the case, with a meta-analysis showing those who stutter to be a standard deviation above  
144 controls for social anxiety scores (Craig & Tran, 2014). In particular, those who stutter are at  
145 high risk of social anxiety disorder diagnosis (Blumgart et al., 2010b; Iverach & Rapee, 2014;  
146 Iverach et al., 2016; Stein et al., 1996). It appears that the origins of these issues can be  
147 detected during early childhood (Briley et al., 2019; Langevin et al, 2010; McAllister, 2016).  
148 Briley et al. (2019) and McAllister (2016) used the Strengths and Difficulties Questionnaire  
149 in two large population studies; the questionnaire measures behavioral, emotional, and social

150 well-being, with high scores indicating anxiety. Both studies reported significantly higher  
151 scores for preschool-age children who were stuttering up to age 5 years, compared with non-  
152 stuttering children. Langevin et al. (2010) reported the findings of a questionnaire sent to 77  
153 parents of preschoolers who stutter, in which 90% of respondents reported clear signs of  
154 anxiety in their child, such as withdrawal and avoidance. One report of older children  
155 (Iverach et al., 2016) found that 24% of 7–12-year-old children who stutter received a social  
156 anxiety disorder diagnosis, compared with 4.6% of control children. For social anxiety in  
157 general, a systematic review and meta-analysis (Bernard et al., 2022) concluded that children  
158 and adolescents who stutter present with increased anxiety symptoms in comparison to their  
159 non-stuttering peers. The connection between stuttering and social anxiety is probably due to  
160 marginalisation caused by negative social stereotypes, microaggressions, bullying, stigma,  
161 and social exclusion (Boyle, 2018; Coalson et al. 2022; Doody et al., 1993; Erickson &  
162 Block, 2013; Ham, 1990; White & Collins, 1984). Negative social stereotypes pertain to all  
163 ages (Horsley & Fitzgibbon, 1987; Woods & Williams, 1976; Yairi & Williams, 1970).  
164 There is evidence that those stereotypes extend to perceptions of physical attractiveness (Van  
165 Borsel et al., 2011) and that stuttering can affect personal relationships (Connery et al., 2020).

#### 166 **Educational and occupational attainment**

167 Stuttering may lead to impaired educational and occupational attainment. For adults, an  
168 inverse relationship has been reported for stuttering severity and educational attainment  
169 (Boyce et al., 2022; O’Brian et al., 2011). A large cohort report by Boyle et al. (1994) found  
170 significantly more chance of repeating a grade for children who stutter compared with control  
171 children. That effect was replicated by Berchiatti et al. (2020). There is evidence that children  
172 experience social isolation in school from fear of speaking in the classroom, and that they  
173 habitually avoid it (Daniels et al., 2012; Klompas & Ross, 2004). College professors in the  
174 US have been shown to have more negative perceptions of students who stutter than controls,

175 and students who stutter were less comfortable than controls when approaching professors  
176 (Werle & Byrd, 2022). Not surprisingly, then, adults who stutter tend to have restricted  
177 occupational opportunities and outcomes (Blumgart et al., 2010a; Gerlach et al., 2018; Klein  
178 & Hood, 2004; McAllister et al., 2012).

### 179 **The present reviews**

180 In summary, stuttering is a prevalent and potentially lifelong disorder, which is capable of  
181 impairing health-related quality of life across a range of domains, including qualitative and  
182 quantitative impacts on speech output, mental health issues, and failure to attain educational  
183 and occupational potential. Although some individuals find strength, emotional growth and  
184 relationship benefits through their stuttering (Boyle et al, 2019; Constantino, 2016), a review  
185 of qualitative studies of the matter (Connery et al., 2020) concluded that stuttering has a  
186 “profound and predominantly negative impact” (p. 2232) on the experiences of those  
187 affected.

188 The present systematic review was designed to explore this matter by seeking information  
189 from existing literature to understand how stuttering affects quality of life and how  
190 interventions have potential to change quality of life. For the review, we did not explore  
191 measures that pertain specifically to the disorder of stuttering. Instead, we sought overarching  
192 quality of life utility measures from traditional health economics that are pertinent to health  
193 policy and funding. Such measures place health care for stuttering into a broader perspective.  
194 The review was also designed to explore economic impact of stuttering: the cost of illness or  
195 the economic costs of stuttering interventions. Costs involved are to the individual in terms of  
196 treatment expenses and lost work hours, along with carer burden, health and education  
197 system costs, and wider societal costs. Those costs are an essential comparison against  
198 quality of life and how it can be changed by intervention, and how it can guide health care



199 funders and health care policy makers to compare different interventions for different  
200 diseases.

## 201 **Methods**

### 202 **Inclusion criteria**

203 Studies were included if they involved children, adolescents, or adults with stuttering as a  
204 primary diagnosis. Participants with comorbid disorders were excluded because the outcomes  
205 of treating them would not be generalisable to participants with stuttering only. The review  
206 included quality of life instruments that directly measured health-related quality of life with  
207 utility scores. For the cost analysis all studies that detailed direct, indirect, and societal costs  
208 of stuttering were included. All interventions and all study types were included.

### 209 **Search strategies**

210 An initial search of Embase and MEDLINE Medical Subject Headings was conducted to  
211 identify articles relating to the costs and outcomes of stuttering therapy. Key words from the  
212 titles, abstracts, subject headings, and other index terms were used to develop a full Boolean  
213 search strategy. The search strategy was refined at a multidisciplinary meeting that included  
214 speech-language pathologists, health economists, and a clinical psychologist. The details of  
215 the search strategy are presented in Supplemental Material S1.

216 The databases searched were: Embase; MEDLINE; PsycINFO; Cumulative Index to  
217 Nursing and Allied Health Literature (CINAHL); the Cochrane Library, including the  
218 Cochrane Database of Systematic Reviews and Cochrane Central Register of Controlled  
219 Trials; ProQuest; Web of Science; Scopus; Paediatric Economic Database Evaluation  
220 (PEDE); and the National Health Service Economic Evaluation Database (NHS EED). To  
221 identify unpublished literature The Australian New Zealand clinical trials registry  
222 (ANZCTR), United States National Institutes of Health trial register, and Open Grey were

223 searched. Searches were limited to English language. No limits on publication year were  
224 applied.

### 225 **Search dates**

226 The cost of illness and quality of life searches were run between 19<sup>th</sup> January 2021 and  
227 20<sup>th</sup> May 2021. After the initial results were retrieved, searches of Embase, MEDLINE,  
228 PsycINFO, and Scopus were automatically re-run at weekly intervals and searches of  
229 CINAHL, ProQuest and Web of Science were re-run at monthly intervals until 13<sup>th</sup> August  
230 2021. Database searches were re-run between the 14<sup>th</sup> and 15<sup>th</sup> February, 2022, to identify  
231 records indexed after 13<sup>th</sup> August, 2021. The Cochrane Library, ANZCTR, and the United  
232 States National Institutes of Health trial register were manually re-searched on 13<sup>th</sup> August  
233 2021 and 14–15<sup>th</sup> February 2022. PEDE, NHS EED, and Open Grey ceased to be updated on  
234 31<sup>st</sup> Dec 2020, 31<sup>st</sup> March 2018, and 27<sup>th</sup> July 2018, respectively so were not re-searched in  
235 2022. Additional records identified were added to the search results. Citations included in the  
236 reference lists of included studies were manually searched to identify additional papers of  
237 interest.

### 238 **Search procedures**

239 Following the searches, citations were collated in Excel, and duplicates removed. Titles  
240 and abstracts were screened against the inclusion criteria and potentially relevant studies  
241 retrieved in full. Retrieved full text articles were screened in detail against the inclusion  
242 criteria, and relevant articles were included in the review. Eligibility decisions at the  
243 title/abstract and full text stage were recorded in a spreadsheet. Two investigators  
244 independently conducted title/abstract and full text reviews for relevance and any  
245 disagreement was resolved through consensus. The search strategies are presented in  
246 PRISMA flow diagrams (Moher et al., 2009): quality of life in Figure 1 and cost of illness in  
247 Figure 2.

248

249

INSERT FIGURES 1 and 2 ABOUT HERE

250

**251 Data extraction form**

252

A data extraction form for included studies was developed, based on the Cochrane checklist of items to consider in data collection (Li et al., 2021), and piloted prior to use. The form is presented in Supplemental Material S2. Relevant data were extracted from each study and recorded in Microsoft Access. The data extracted included details about the study (design, methods, year, location), participants (age, gender, comorbidity, socioeconomic status, ethnicity), intervention (setting, duration), and outcomes (costs and health outcomes). No summary measures were pre-specified.

259

**Risk of bias assessment**

260

The risk of bias was assessed at the study level using the RoB 2 tool for randomized trials (Sterne et al., 2019), and the ROBINS-I tool (Sterne et al., 2016) for non-randomized studies. Qualitative research was assessed using the CASP qualitative studies checklist (Critical Appraisal Skills Programme, 2018). The CASP checklist is cited by The Cochrane Qualitative and Implementation Methods Group as “the most commonly used tool in qualitative evidence synthesis in Cochrane and World Health organisation guideline processes” (Noyes et al., 2018, p. 50). Risk of bias details for included studies are presented in Supplemental Material S3.

268

**Cost standardisation**

269

Costs were inflated to 2020/21 values using an inflation calculator relevant to the reported currency, then converted into US dollars where necessary using the exchange rate listed at [www.xe.com](http://www.xe.com) on 20 August, 2021.

270

271

272

**Results****273 Quality of life**

274 The quality of life search strategy identified 2,067 reports, of which three were included in  
275 the quality of life analysis. From these reports, 30 studies were excluded during the review  
276 because they did not generate a utility measure. Of these 30 studies, 19 used eleven different  
277 instruments directly pertinent to quality of life. Details are presented in Figure 1 and Table 1.

278 Three reports measured quality of life using utility measures (de Sonnevile-Koedoot et  
279 al., 2015; McAllister et al., 2017; Omori et al., 2021). These involved 242 participants: 199  
280 children, 31 adults, 12 in a mixed cohort. The proportion of participants who were male  
281 ranged by treatment group from 60% to 94%. The three studies are summarized in Table 2.

282

283

INSERT TABLE 1 and 2 ABOUT HERE

284

285 The utility measures used were the Health Utilities Index 3 (HUI3), the 3-level version of  
286 EuroQol-5 Dimension (EQ-5D-3L), and the 5-level EQ-5D questionnaire. HUI3 considers  
287 eight attributes: vision, hearing, speech, ambulation, dexterity, emotion, cognition, and pain.  
288 EQ-5D considers five dimensions: mobility, self-care, usual activities, pain/discomfort, and  
289 anxiety/depression.

290

291 In the de Sonnevile-Koedoot et al. (2015) study, using the HUI3 instrument with  
292 preschool children, the mean baseline utility score was 0.88 for both treatment groups  
293 (RESTART-DCM and Lidcombe Program). At 18-months follow-up, the mean utility score  
294 improved for both groups by 0.07 in the Lidcombe Program group and 0.06 for the  
295 RESTART-DCM group. In the Omori et al. (2021) study of adults, the mean baseline utility  
296 score was 0.80 for the intervention group (Cognitive Behavior Therapy (CBT) + speech  
treatment) and 0.71 for the control group (speech treatment), using the 5-level EQ-5D

297 instrument. After 4 months of treatment, the mean intervention group (CBT + speech  
298 treatment) utility score increased by 0.10 utility score values, and with the control group  
299 (speech treatment), it increased in a favourable direction by 0.06 utility score values. In the  
300 McAllister et al. (2017) study of adults, the mean baseline utility score was 0.82 for the  
301 intervention group and 0.83 for the control group, using the EQ-5D-3L instrument. At 4  
302 months post intervention, the mean intervention group utility score worsened slightly in both  
303 groups by 0.02 utility score values. No study reported a statistically significant difference in  
304 utility scores between interventions.

305 The McAllister et al. (2017) randomized trial was considered to have a low risk of bias,  
306 and the de Sonnevile-Koedoot et al. (2015) trial was considered to have a high risk of bias  
307 due to the potential for loss of allocation concealment. The Omori et al. (2021) non-  
308 randomized trial had a high risk of bias for several reasons (see Supplemental Material S3).

### 309 **Cost of illness**

#### 310 **Interventions and Comparators**

311 The cost of illness search strategy identified 3,778 reports. From these reports, 39 studies  
312 were included in the cost of illness analysis. Of these studies, 36 did not include interventions  
313 or comparators. Details are presented in Figure 2. One report compared two stuttering  
314 interventions (the Lidcombe Program and RESTART-DCM) (de Sonnevile-Koedoot et al.,  
315 2015), and one report compared a web-based cognitive bias modification program with  
316 placebo (McAllister et al., 2017). One report presented long-term follow-up data for two  
317 single-arm trials of smooth speech treatment for stuttering (Craig & Calver, 1991).

#### 318 **Outcomes**

319 The direct costs of treatment, indirect costs connected with treatment, and societal costs of  
320 stuttering were of interest for the cost analysis. Societal costs were grouped into three themes:  
321 education, employment, and social outcomes. In total, 155 cost outcomes were extracted. Of

322 the 39 included studies, six provided evidence on healthcare utilisation, six provided evidence  
323 on direct and indirect costs of stuttering, 20 provided evidence for the impact of stuttering on  
324 educational outcomes, 22 provided evidence on employment outcomes, and 11 provided  
325 evidence on social outcomes, and one presented information on society's willingness to pay  
326 for stuttering treatment. A summary of studies included in the cost of illness analysis is  
327 presented in Supplemental Material S4, along with overall risk of bias for the studies (see  
328 Supplemental Material S3 for further risk of bias details).

### 329 **Healthcare utilisation**

330 There is evidence of trends in health care utilisation for those who stutter, though most  
331 findings were not statistically significant. Children who stutter were (a) more likely to be  
332 hospitalized and spend more days in hospital than children with no developmental disabilities  
333 (Boyle et al., 1994); (b) more likely to attend hospital emergency departments ( $p < .01$ ) and  
334 undergo surgical or medical procedures ( $p < .05$ ) than children with no developmental  
335 disabilities (Boulet et al., 2009); (c) more likely to make frequent doctor visits than children  
336 with no developmental disabilities (Boyle et al., 1994); (d) more likely to visit specialists ( $p <$   
337  $.05$ ), allied health professionals ( $p < .01$ ), and mental health professionals (difference not  
338 significant); and (e) more likely to take medication for longer than 3 months and require  
339 special equipment than children with no developmental disabilities (Boulet et al., 2009).

340 A retrospective analysis of health insurance data by Sommer et al. (2021) reported that  
341 approximately 45% of individuals of any age diagnosed with stuttering seek speech treatment  
342 within a year of diagnosis and receive an average of 13.5 speech therapy sessions in the first  
343 year of treatment. A report by de Sonnevile-Koedoot (2015) indicated that children treated  
344 with the Lidcombe Program received 20 hours of stuttering therapy over an 18-month period,  
345 compared with 18 hours with RESTART-DCM. McAllister et. al. (2017) collected data at  
346 baseline and 4 months follow-up for adults on hospitalisations, day hospital, primary care

347 visits, specialists, other therapy, medication and special equipment. In this report, individuals  
348 in the intervention group, at baseline, had received 3.1 primary care visits and 2.4  
349 prescriptions in the previous 4 months. Individuals in the placebo group received 2.0 primary  
350 care visits and 4.5 prescriptions at baseline. Mean specialist visits, other therapy,  
351 hospitalisations, and special equipment use were all below 1.0 at baseline in the intervention  
352 and control groups.

### 353 **Direct and indirect costs of stuttering**

354 A study by Blumgart et al. (2010a) reported the mean cost of stuttering treatment over five  
355 years to be the equivalent of US\$2,528. The overall direct cost of stuttering over five years,  
356 including speech treatment, other treatment, technology, self-help, and other costs was  
357 US\$5,101 per adult client (Blumgart et al., 2010b). The mean cost of stuttering treatment in  
358 one randomized controlled trial of early intervention was US\$1,976 for the Lidcombe  
359 Program and US\$1,788 for RESTART-DCM (de Sonnevle-Koedoot et al., 2015). Total  
360 direct costs were US\$3,681 for the Lidcombe Program and US\$3,541 for RESTART-DCM:  
361 including additional allied health, home therapy, travel costs, and parent time associated with  
362 home therapy. Total costs, including absenteeism and productivity losses over the 18-month  
363 study period were US\$4,444 for the Lidcombe Program and US\$4,212 for RESTART-DCM.

364 One qualitative research report conducted by Georges (2017) considered the financial  
365 impact of stuttering. Seven of 10 participants reported no financial strain associated with  
366 stuttering. However, the same number noted that their health insurance did not cover  
367 stuttering treatment. These individuals described finding alternative methods to fund therapy  
368 or seeking alternatives to treatment. A report of the impact of stuttering on adolescents and  
369 their families by Erickson and Block (2013) indicated that stuttering placed at least some  
370 financial strain on 61% of families. Of the parents interviewed, 19% missed work and 10%  
371 had made time and financial sacrifices due to their child's stuttering.

372 In a cross-sectional study investigating society's willingness to pay for stuttering  
373 treatment, Franic et al. (2012) reported that the mean amount respondents were willing to pay  
374 ranged from US\$24,927 for an intervention that improved stuttering from severe to mild to  
375 US\$61,810 for an intervention that stopped severe stuttering; this was equivalent to between  
376 two and four times their annual income.

### 377 **Education**

378 Stuttering impacted education for a range of measures, including performance, attainment,  
379 attendance, and the requirement of special support services. In two reports, McClure and  
380 Yaruss (2003) reported 80% of respondents stating that stuttering interfered with their  
381 schoolwork, and Hayhow et al. (2002) reported 95%. Two reports indicated academic  
382 difficulties associated with stuttering that were indirectly linked to the occurrence of bullying  
383 (Erickson & Block, 2013; Hugh-Jones & Smith, 1999). Another two reports indicated that  
384 children who stutter were significantly more likely to require special education services or  
385 early intervention compared with children who have no developmental disabilities (Boyle et  
386 al., 1994; Boulet et al., 2009). Two further reports noted a link between stuttering and  
387 avoiding school, mediated by the presence of bullying (Erickson & Block, 2013; Hugh-Jones  
388 & Smith, 1999); and another reported the mean number of school days lost each year by  
389 children who stutter to be 7.7 compared with 3.0 days for children with no developmental  
390 disabilities ( $p < .01$ ) (Boyle et al., 1994).

391 Three studies reported that children who stutter perform worse academically than children  
392 who do not stutter (Berchiatti et al., 2020; Calnan & Richardson, 1977; Williams et al. 1969).  
393 Boyle et al. (1994) reported that 29% of children who stutter had repeated a grade, compared  
394 with 13% of children with no developmental disabilities ( $p < .01$ ). Conversely, a study of  
395 college instructors evaluating oral presentations reported that the performances of individuals  
396 who stutter were evaluated significantly higher than fluent controls ( $p < .01$ ), equating to a



397 “full letter” grade difference in total score (Werle & Byrd, 2022). Another study reported a  
398 significant inverse relationship between stuttering severity and highest educational  
399 achievement (O’Brian et al., 2022); and another reported that stuttering predicted lower  
400 educational attainment ( $p < .01$ ) (Rosenbaum, 2018). However, two additional follow-up  
401 studies reported that after controlling for confounding variables such as comorbidities and  
402 family background, stuttering was not a significant contributor to educational attainment  
403 (McAllister et al. 2012; Rees & Sabia, 2014).

404 Similar findings were observed in qualitative research. Participants in one study reported  
405 that stuttering had a negative effect on academic performance (Silverman & Zimmer, 1982).  
406 Another study contained reports of being marked down for not attending university seminars,  
407 and even choosing university courses with less interaction (Butler, 2013). The impact on  
408 educational attainment was mixed, with some studies reporting high levels of tertiary  
409 completion for adults who stutter (Leko Krhen et al., 2021; Silverman & Zimmer, 1982).  
410 Participants in other studies reported low educational achievement and leaving school  
411 prematurely (Butler, 2014; Crichton-Smith, 2002; Johnson, 1934).

#### 412 **Employment**

413 Stuttering impacted employment for a range of measures, including job performance,  
414 occupational choice, unemployment, underemployment, workplace discrimination,  
415 promotion, and income. From three reports, a range of 69–86% of respondents stated that  
416 stuttering interfered with their job performance (Klein & Hood, 2004; McClure & Yaruss,  
417 2003; Rice and Kroll, 2006). Another report stated that stuttering severity had a medium to  
418 high correlation with self-rated job difficulties ( $p < .01$ ) (Iimura & Miyamoto, 2022). Two  
419 reports indicated that approximately 50% of respondents stated that stuttering had affected  
420 their occupational choice (Hayhow et al., 2002; Rice & Kroll, 2006), with a similar  
421 proportion in two other reports stating that finding employment was difficult (Blumgart et al.,

422 2010a; Boyle, 2018). A workforce analysis comparing adults who stutter with controls  
423 reported significant differences between the groups in terms of choice of industry (Plexico et  
424 al., 2019); it was also reported that there were more adults who stutter than controls in the  
425 information industry ( $p = .008$ ) and fewer in sales or service roles ( $p = .02$ ).

426 An assessment of United States labor market outcomes found no significant difference  
427 between adults who stutter and controls in terms of the number of people who were employed  
428 in the labor force, underemployed, or receiving public assistance after controlling for  
429 confounding (Gerlach et al., 2018). However, a similar study conducted in the United  
430 Kingdom found that, although stuttering at age 16 was not a significant predictor of  
431 unemployment or socioeconomic class of occupation at 23, it was a significant predictor of  
432 socioeconomic class of occupation at 50 ( $p = .047$ ) (McAllister et al., 2012).

433 In a report by Boyle (2018), 70% of respondents indicated discrimination in the workplace  
434 (Boyle, 2018). The workforce analysis by Plexico et al. (2019) demonstrated that adults who  
435 stutter reported higher levels of discrimination than controls ( $p < .01$ ). In another report, 8%  
436 of respondents said their employment was terminated due to their stuttering (Blumgart et al.,  
437 2010a). A further two reports described discrimination in employment (Hayhow et al., 2002;  
438 Rice & Kroll, 1994). Reports indicate that 20–27% of respondents turned down a job or  
439 promotion, and 28–38% were denied a promotion due to their stuttering (Blumgart et al.,  
440 2010a; Klein & Hood, 2004; Rice & Kroll, 2006). The workforce analysis by Plexico et al.  
441 reported significantly fewer adults who stutter than controls in management roles ( $p = .04$ )  
442 and supervision ( $p = .01$ ) roles. Craig and Calver (1991) found that within 10 months of  
443 completing a speech treatment program, 44% of participants who stutter who were eligible  
444 for promotion. Evidence for the impact of stuttering on income is mixed, Gerlach et al.  
445 (2018) reporting that stuttering is associated with an annual income deficit of at least  
446 US\$9,054 ( $p < .05$ ) and McAllister et al. (2012) reporting that stuttering at 16 years old is not

447 a significant predictor of income at 23 or 50 years. The workforce analysis by Plexico et al.  
448 reported that adults who stutter had less income than controls ( $p < .05$ ).

449 Many participants in qualitative studies reported that stuttering had affected their working  
450 lives (Crichton-Smith, 2002; Georges, 2017; Silverman & Zimmer, 1982). In a report by  
451 Nang et al. (2018), all the women interviewed indicated that stuttering limited their ability to  
452 secure work and perform work-related tasks. There were several other reports of stuttering  
453 influencing occupational choice (Butler, 2014; Crichton-Smith, 2002; Georges, 2017;  
454 Johnson, 1934). There were also reports of discrimination (Nang et al., 2018), challenges in  
455 recruitment (Butler, 2013; Crichton-Smith, 2002), and challenges in promotion (Bricker-Katz  
456 et al., 2013). However, in one report by Leko Krhen et al. (2021), none of the women  
457 indicated that stuttering had affected their occupation choice or presented significant  
458 professional challenges.

#### 459 **Social**

460 Stuttering also has a social cost, affecting individuals through bullying, stigma, and social  
461 exclusion, with the potential to negatively impact friendships and life relationships. There are  
462 reports indicating that 53–77% of respondents were teased or bullied and 20–55% were  
463 avoided by others or excluded from social situations (Boyle, 2018; Erickson & Block, 2013).  
464 Boyle (2018) reported that 58–81% of respondents experienced other types of enacted  
465 stigma, including discrimination: treated unfairly, considered as inferior, considered less  
466 seriously, or being patronized. A reported personal effect of bullying can be a difficulty in  
467 forming relationships (Hayhow et al., 2002; Hugh-Jones & Smith, 1999). In the Hayhow et  
468 al. (2002) report, 61% of respondents reported that stuttering had impacted their life  
469 relationships, and 64% of respondents reported that stuttering had impacted their friendships.  
470 An analysis of school children by Berchiatti et al. (2020) reported that children who stutter

471 were less popular ( $p < .01$ ) and more rejected ( $p < .001$ ) in the peer group than control  
472 children.

473 Thematic results of qualitative research support these findings. Responses suggest that  
474 stuttering can lead to anxiety, avoidance, embarrassment, and frustration in those who stutter  
475 and their life partners (Beilby et al., 2013). Some respondents reported problems in life  
476 relationships (Georges, 2017). One report noted the emotional impact on life partners  
477 (Boberg & Boberg, 1990). However, a key theme was the importance of supportive partners,  
478 many of whom were not bothered by stuttering (Beilby et al., 2013; Leko Krhen et al., 2021;  
479 Nang et al., 2018).

## 480 **Discussion**

### 481 **General study limitations**

482 Interpretation of the present results requires consideration of their limitations. Results  
483 cannot be generalized to non-English literature, because only English language publications  
484 were searched. As with all systematic reviews, results were limited by the comprehensiveness  
485 of the databases searched. For the present review, 14 databases were searched, and Open  
486 Grey was searched for unpublished data. Additionally, manual searches of the reference lists  
487 of all identified studies served as a cross-check. However, it is conceivable that some data  
488 published in textbooks and presented in unpublished conference proceedings may have been  
489 overlooked. The sparse finding of only three studies identified in the quality of life search  
490 needs to be considered in light of two of them being classified as having a high risk of bias.  
491 This limits the extent to which their results can be considered admissible. The exclusion of  
492 participants with diagnosed comorbid disorders could also be seen as a limitation. This  
493 exclusion criterion was intended to clarify the findings for stuttering alone. However,  
494 stuttering and comorbidity is, of course, of great interest for health economics research and  
495 requires further exploration.

**496 Quality of life****497 Summary of results**

498 This systematic review involved a full-text review of 135 articles, and only three provided  
499 standard utility values that could be incorporated into health economics analyses. Baseline  
500 utility scores for stuttering adult groups in two studies ranged from 0.71–0.83, and for  
501 preschool children in one study, they were 0.88. Two of the three studies had a high risk of  
502 bias. Post-intervention improvements in utility values were reported in the range of 0.06–0.10  
503 for two studies, and a slight worsening was reported in one study. No study reported a  
504 statistically significant difference in utility scores between treatment groups.

**505 Instrument limitations**

506 Age restrictions for the quality of life test instruments limit the usefulness of results  
507 identified in this review. The cost utility analysis based on HUI3 utilities in de Sonnevile-  
508 Koedoot et al. (2015) is limited by the number of participants 5 years of age or older for  
509 whom the instrument was applicable, which was 25% of total participants. The cost utility  
510 analysis based on transformed EQ-VAS scores—a visual analogue scale from 0 to 100—used  
511 by de Sonnevile-Koedoot et al. is not supported by a published mapping algorithm. The  
512 Child Health Utility 9 dimensions (CHU-9D) is an alternative quality of life instrument  
513 appropriate for children 4–17 years of age (Rowen et al., 2020). Using the CHU-9D instead  
514 of HUI3 by de Sonnevile-Koedoot et al. would have expanded the cohort with applicable  
515 utility scores to children older than 4 years of age, increasing the sample size with utility data  
516 to 120 participants (60% of the study total).

517 There is currently no generic quality of life instrument appropriate for children younger  
518 than 4 years of age (Rowen et al., 2020). However, the Pediatric Quality of Life Inventory  
519 measures quality of life in children 2 years of age and older, and it could be used in stuttering  
520 research to maximize the amount of quality of life data available for economic evaluation.

521 Test instrument sensitivity is also a limitation of the quality of life assessments identified in  
522 this review. The EQ-5D and HUI3 consider psychological distress; however, both were found  
523 to be less sensitive to mental health quality of life than other instruments, such as Assessment  
524 of Quality of Life–Eight Dimension Scale (Mihalopoulos et al., 2014).

#### 525 **Conversion of instruments to utility measures**

526 It is possible that two of the quality of life instruments excluded during this review could  
527 be converted into utility measures: EQ-VAS and SF-36 (the 36-Item Short Form Health  
528 Survey). Although, a recent review of mapping algorithms that link results from quality of  
529 life instruments to EQ-5D utility values did not identify a method to convert EQ-VAS to EQ-  
530 5D (Dakin et al., 2018). However, a scoring system linking the visual analogue scale to the  
531 utility measure of the EQ-5D youth version (EQ-5D-Y) exists (Wu, 2014), which suggests  
532 that conversion of EQ-VAS scores to utility measures is possible. The results of SF-36 can be  
533 mapped to multiple generic utility measures, including HUI3 and EQ-5D (Brazier et al.,  
534 2010). Given that 21 studies evaluated the impact of stuttering interventions on quality of life  
535 using the Quality of Life scale of the Overall Assessment of the Speaker’s Experience of  
536 Stuttering (OASES), mapping OASES to EQ-5D or another generic quality of life measure  
537 would facilitate the inclusion of valuable existing research in future economic evaluations.

#### 538 **Interpretation of reported stuttering utility values**

539 The baseline utility scores for the two studies with adults (McAllister et al., 2017; Omori  
540 et al., 2021) were limited by low participant numbers. McAllister et al. (2017) reported utility  
541 scores for 22 participants, and Omori et al. (2021) reported utility scores for 12 participants.  
542 With the caveat of limited sampling, those data did substantiate an earlier suggestion (Craig  
543 et al., 2009) that the quality of life impact of stuttering is similar to other conditions that are  
544 of prime concern to health care provision. Stuttering utility scores for adults in the range of  
545 0.71–0.83 are broadly consistent with values associated with utility scores for some

546 conditions reported in another systematic review (Zhou et al., 2021). For example, Zhou et al.  
547 (2021) reported a meta-analytic utility estimate of 0.83 for diabetes mellitus, and pooled  
548 values of 0.77 for cardiovascular disease, 0.75 for cancer, 0.84 for HIV, and 0.70 for chronic  
549 kidney disease.

550 Sampling was more substantive in the de Sonnevile-Koedoot et al. (2015) report for  
551 preschoolers, although, as noted above, it was limited to participants 5 years of age or older  
552 for whom standard utility scores were available. Baseline utility scores for children (mean =  
553 0.88) were higher than for adults, reflecting less quality of life impact. It is intuitively correct  
554 that quality of life impact is lower for preschoolers than adults. This is because shortly after  
555 stuttering potentially debilitating factors of mental health and educational and occupational  
556 disadvantage would have had limited influence.

#### 557 **Interpretation of treatment results**

558 The Omori et al. (2021) study involved basic stuttering control training for both groups,  
559 with the intervention group receiving a 7-week CBT package. Results were encouraging after  
560 4 months, with the intervention group increasing its utility score by 0.10 and the standard  
561 care group by 0.06. Arguably, that was a conservative estimate, considering that the standard  
562 stuttering control training was limited to “20-minute speech therapy sessions once every three  
563 or four weeks for four months” (p. 59). With preschool participants in the de Sonnevile-  
564 Koedoot et al. (2015) report for whom standard utility scores were available, results are a  
565 little more convincing, with similar utility score improvements in both intervention groups at  
566 18 months follow-up: 0.07 for Lidcombe Program and 0.06 for RESTART-DCM. However,  
567 arguably again, those results were conservative estimates because at follow-up, 28% of  
568 children in the Lidcombe Program arm and 35% in the RESTART-DCM arm had not  
569 completed their treatments (Franken, 2016). The de Sonnevile-Koedoot et al. report did not

570 involve a no-treatment control group, which makes it unclear to what extent follow-up  
571 improvements were due to natural recovery (Bergþórsdóttir & Ingham, 2017).

## 572 **Cost of illness**

### 573 **Summary of results**

574 Overall, this body of research carries considerable risk of bias. Regardless, it seems that  
575 those who stutter—children and adults—consume more health care resources than their non-  
576 stuttering peers, including those relating to hospitalisations, outpatient visits, consultations  
577 with mental health professionals, and medications. Many of the results in the reviewed  
578 reports were not statistically significant. However, that could be attributed to study designs,  
579 in particular, limited controls for confounding variables. The direct, overall costs of  
580 treatments obtained for the adult population are in the vicinity of US\$1,000 per year, and the  
581 total costs of well-known early interventions are around US\$4,000-4,500. There is some  
582 evidence of financial strain on families with adolescents who are receiving treatment. One  
583 report considered willingness to pay for stuttering interventions. Respondent willingness to  
584 pay for treatment to improve or cure stuttering was quantified as two to four times their  
585 annual income.

586 Stuttering appears to be associated with lower academic performance in childhood and  
587 may contribute to reduced attainment of tertiary qualifications. Children who stutter are more  
588 likely than their peers to require special education services. They are twice as likely to not  
589 attend school days, apparently because of their stuttering, and this has been found to be  
590 related to bullying. It is possible that there is an inverse relationship between stuttering  
591 severity and educational attainment, and stuttering has been linked to leaving school  
592 prematurely. However, those findings are potentially confounded by uncontrolled variables.  
593 One report suggested that tertiary educators may overcorrect performance assessments to



594 account for stuttering. The authors of that report note that such overcorrection may ultimately  
595 have a negative impact and limit long-term academic achievement.

596 There is clearly a relationship between stuttering and employment, where those who  
597 stutter are limited in terms of income, promotion, and socioeconomic status of occupation.  
598 Those limitations may affect women particularly. Stuttering is reported to affect occupational  
599 choice, and it is associated with workplace discrimination. There is some evidence  
600 associating stuttering with reduced income.

601 There are social costs associated with stuttering, involving bullying, stigma, and social  
602 exclusion. Stuttering has the potential to negatively impact friendships and life relationships.  
603 Children who stutter are less popular and rejected more often than their non-stuttering peers.  
604 Stuttering can lead to anxiety, avoidance, embarrassment, and frustration in those who stutter  
605 and their life partners. Supportive life partners assist those who stutter to deal with the  
606 condition.

### 607 **Conclusions**

608 In health economics terms, stuttering has substantive direct and indirect costs of illness. It  
609 is likely to cause impairment, challenges, and distress across many domains throughout life,  
610 including income, education, employment, and social functioning. There is some evidence  
611 that, in health economics terms, stuttering causes impairment of health-related quality of life  
612 and that it can be improved by treatment. However, that evidence is extremely limited and is  
613 derived from only 85 participants across three studies. This evidence gap limits economic  
614 evaluation and cost-utility analyses of stuttering interventions.

615 There is a growing body of clinical trials evidence to support treatments for stuttering. For  
616 adult and adolescents, variations of speech restructuring have been evaluated in more than 30  
617 clinical trials (Brignell et al., 2020). There are eight randomized controlled trials of the  
618 Lidcombe Program for children younger than 6 years (Sjøstrand et al., 2021). Additionally,

619 there are five successful clinical trials of CBT specifically for the social anxiety of those who  
620 stutter (Gunn et al., 2019; Helgadóttir et al., 2014; Menzies et al., 2008, 2009, 2019). Yet, to  
621 date, that evidence has been overlooked for its capacity to improve quality of life for children  
622 and adults who stutter.

623 Eleven instruments pertinent to quality of life were excluded from the analysis because  
624 they did not generate a utility measure. Most strikingly, the Overall Assessment of the  
625 Speakers Experience of Stuttering, with its Quality of Life subscale, was presented in 21  
626 reports but was excluded from this review. Because of this, the literature to date contains  
627 much potentially useful information that cannot be used to inform cost-utility analyses of  
628 stuttering interventions. Without utility values, treatment for the impairment, challenges, and  
629 distress experienced by those who stutter cannot be presented in a cost-utility analysis,  
630 conveying cost per quality-adjusted life year. This means that the health-related quality of life  
631 impairment of those who stutter is not documented in a form that can be used to influence  
632 health policy and healthcare spending. That problem could be remedied if future clinical trials  
633 of stuttering treatment routinely incorporate a health-related quality of life measure that can  
634 be converted to a utility score. That course of action is our recommendation from this  
635 systematic review.

636

637

### **Acknowledgements**

638 This research was supported by National Health and Medical Research Council Program  
639 Grant number #1132370.

640 The authors would like to acknowledge Jeremy Cullis and Mary Simons (clinical  
641 librarians at Macquarie University Library) for their assistance with designing search  
642 strategies, navigating databases, and collating output in a consistent format. The authors

643 would also like to thank Damien Liu-Brennan for his scientific copyediting contribution to  
644 this publication.

645 **Figure Captions**

646 Figure 1. Results of the quality of life search strategy.

647 Figure 2. Results of the cost of illness search strategy.

648

**References**

- 649 Ambrose, N. G., & Yairi, E. (1999). Normative disfluency data for early childhood  
650 stuttering. *Journal of Speech, Language, and Hearing Research*, 42(4), 895–909.  
651 <https://doi.org/10.1044/jslhr.4204.895>
- 652 Beilby, J. M., Byrnes, M. L., Meagher, E. L., & Yaruss, J. S. (2013). The impact of stuttering  
653 on adults who stutter and their partners. *Journal of Fluency Disorders*, 38(1), 14–29.  
654 <https://doi.org/10.1016/j.jfludis.2012.12.001>
- 655 Berchiatti, M., Badenes-Ribera, L., Ferrer, A., Longobardi, C., & Gastaldi, F. G. M. (2020).  
656 School adjustment in children who stutter: The quality of the student-teacher relationship,  
657 peer relationships, and children’s academic and behavioral competence. *Children and*  
658 *Youth Services Review*, 116, Article 105226.  
659 <https://doi.org/10.1016/j.chilyouth.2020.105226>
- 660 Bergþórsdóttir, Í. Ö., & Ingham, R. J. (2017). Putting the cart before the horse: A cost  
661 effectiveness analysis of treatments for stuttering in young children requires evidence that  
662 the treatments analyzed were effective. *Journal of Communication Disorders*, 65, 65–67.  
663 <https://doi.org/10.1016/j.jcomdis.2016.04.006>
- 664 Bernard, R., Hofslundsengen, H., & Frazier Norbury, C. (2022). Anxiety and depression  
665 symptoms in children and adolescents who stutter: A systematic review and meta-  
666 analysis. *Journal of Speech, Language, and Hearing Research*, 65(2), 624–644.  
667 [https://doi.org/10.1044/2021\\_JSLHR-21-00236](https://doi.org/10.1044/2021_JSLHR-21-00236)
- 668 Bloodstein, O., Bernstein Ratner, N., & Brundage, S. B. (2021). *A handbook on stuttering*  
669 (7th ed.). Plural Publishing.
- 670 Blumgart, E., Tran, Y., & Craig, A. (2010a). An investigation into the personal financial  
671 costs associated with stuttering. *Journal of Fluency Disorders*, 35(3), 203–215.  
672 <https://doi.org/10.1016/j.jfludis.2010.03.002>

- 673 Blumgart, E., Tran, Y., & Craig, A. (2010b). Social anxiety disorder in adults who stutter.  
674 *Depression and Anxiety*, 27(7), 687–692. <https://doi.org/10.1002/da.20657>
- 675 Boberg, J. M., & Boberg, E. (1990). The other side of the block: The stutterer's  
676 spouse. *Journal of Fluency Disorders*, 15(1), 61-75. [https://doi.org/10.1016/0094-](https://doi.org/10.1016/0094-730X(90)90033-O)  
677 [730X\(90\)90033-O](https://doi.org/10.1016/0094-730X(90)90033-O)
- 678 Boulet, S. L., Boyle, C. A., & Schieve, L. A. (2009). Health care use and health and  
679 functional impact of developmental disabilities among US children, 1997–2005. *Archives*  
680 *of Pediatrics and Adolescent Medicine*, 163(1), 19–26.  
681 <https://doi.org/10.1001/archpediatrics.2008.506>
- 682 Boyce, J. O., Jackson, V. E., van Reyk, O., Parker, R., Vogel, A. P., Eising, E., Horton, S. E.,  
683 Gillespie, N. A., Scheffer, I. E., Amor, D. J., Hildebrand, M. S., Fisher, S. E., Martin, N.  
684 G., Reilly, S., Bahlo, M., & Morgan, A. T. (2022). Self-reported impact of developmental  
685 stuttering across the lifespan. *Developmental Medicine and Child Neurology*, 64(10),  
686 1297–1306. <https://doi.org/10.1111/dmcn.15211>
- 687 Boyle, M. P. (2018). Enacted stigma and felt stigma experienced by adults who  
688 stutter. *Journal of Communication Disorders*, 73, 50-61.  
689 <https://doi.org/10.1016/j.jcomdis.2018.03.004>
- 690 Boyle, C. A., Decoufle, P., & Yeargin-Allsopp, M. (1994). Prevalence and health impact of  
691 developmental disabilities in US children. *Pediatrics*, 93(3), 399–403.  
692 <https://doi.org/10.1542/peds.93.3.399>
- 693 Boyle, M. P., Beita-Ell, C., & Milewski, K. M. (2019). Finding the good in the challenge:  
694 benefit finding among adults who stutter. *Perspectives*, 4(6), 1316–1326.  
695 [https://doi.org/10.1044/2019\\_PERS-SIG4-2019-0024](https://doi.org/10.1044/2019_PERS-SIG4-2019-0024)  
696

- 697 Boyle, M. P., & Gabel, R. M. (2020). “Openness and progress with communication and  
698 confidence have all gone hand in hand”: Reflections on the experience of transitioning  
699 between concealment and openness among adults who stutter. *Journal of Fluency  
700 Disorders*, 65, Article 105781. <https://doi.org/10.1016/j.jfludis.2020.105781>
- 701 Brazier, J. E., Yang, Y., Tsuchiya, A., & Rowen, D. L. (2010). A review of studies mapping  
702 (or cross walking) non-preference based measures of health to generic preference-based  
703 measures. *The European Journal of Health Economics*, 11(2), 215–225.  
704 <https://doi.org/10.1007/s10198-009-0168-z>
- 705 Bricker-Katz, G., Lincoln, M., & Cumming, S. (2013). Stuttering and work life: An  
706 interpretative phenomenological analysis. *Journal of Fluency Disorders*, 38(4), 342–355.  
707 <https://doi.org/10.1016/j.jfludis.2013.08.001>
- 708 Brignell, A., Krahe, M., Downes, M., Kefalianos, E., Reilly, S., & Morgan, A. T. (2020). A  
709 systematic review of interventions for adults who stutter. *Journal of Fluency Disorders*,  
710 64, Article 105766. <https://doi.org/10.1016/j.jfludis.2020.105766>
- 711 Briley, P. M., O’Brien, K., & Ellis, C. (2019). Behavioral, emotional, and social well-being in  
712 children who stutter: Evidence from the National Health Interview Survey. *Journal of  
713 Developmental and Physical Disabilities*, 31, 39–53. [https://doi.org/10.1007/s10882-018-  
714 9625-x](https://doi.org/10.1007/s10882-018-9625-x)
- 715 Butler, C. (2013). ‘University?... hell no!’: Stammering through education. *International  
716 Journal of Educational Research*, 59, 57–65. <https://doi.org/10.1016/j.ijer.2013.03.002>
- 717 Butler, C. (2014). Wanted—straight talkers: Stammering and aesthetic labour. *Work,  
718 Employment and Society*, 28(5), 718–734. <https://doi.org/10.1177/0950017013501956>
- 719 Calnan, M., & Richardson, K. (1977). Speech problems among children in a national  
720 survey—associations with reading, general ability, mathematics and syntactic  
721 maturity. *Educational Studies*, 3(1), 55–66. <https://doi.org/10.1080/0305569770030105>

- 722 Critical Appraisal Skills Programme (2018). CASP Qualitative Studies Checklist.  
723 <https://casp-uk.net/images/checklist/documents/archive/CASP-Randomised-Controlled->  
724 [Trial-Checklist-2018.pdf](https://casp-uk.net/images/checklist/documents/archive/CASP-Randomised-Controlled-Trial-Checklist-2018.pdf)
- 725 Chang, S. E., Garnett, E. O., Etschell, A., & Chow, H. M. (2018). Functional and  
726 neuroanatomical bases of developmental stuttering: Current insights. *The Neuroscientist*,  
727 *25*(6), 566–582. <https://doi.org/10.1177/1073858418803594>
- 728 Coalson, G. A., Crawford, A., Treleaven, S. B., Byrd, C. T., Davis, L., Dang, L., Edgerly, J.,  
729 & Turk, A. (2022). Microaggression and the adult stuttering experience. *Journal of*  
730 *Communication Disorders*, *95*, Article 106180.  
731 <https://doi.org/10.1016/j.jcomdis.2021.106180>
- 732 Connery, A., McCurtin, A., & Robinson, K. (2020). The lived experience of stuttering: a  
733 synthesis of qualitative studies with implications for rehabilitation. *Disability and*  
734 *rehabilitation*, *42*(16), 2232–2242. <https://doi.org/10.1080/09638288.2018.1555623>
- 735 Constantino, C. D. (2016). *Stuttering gain*. International Stuttering Association, International  
736 Stuttering Awareness Day Online Conference.
- 737 Constantino, C. D., Leslie, P., Quesal, R. W., & Yaruss, J. S. (2016). A preliminary  
738 investigation of daily variability of stuttering in adults. *Journal of Communication*  
739 *Disorders*, *60*, 39–50. <https://doi.org/10.1016/j.jcomdis.2016.02.001>
- 740 Craig, A., Blumgart, E., & Tran, Y. (2009). The impact of stuttering on the quality of life in  
741 adults who stutter. *Journal of Fluency Disorders*, *34*(2), 61–71.  
742 <https://doi.org/10.1016/j.jfludis.2009.05.002>
- 743 Craig, A. R., & Calver, P. (1991). Following up on treated stutterers: Studies of perceptions  
744 of fluency and job status. *Journal of Speech, Language, and Hearing Research*, *34*(2),  
745 279–284. <https://doi.org/10.1044/jshr.3402.279>



- 746 Craig, A., & Tran, Y. (2014). Trait and social anxiety in adults with chronic stuttering:  
747 Conclusions following meta-analysis. *Journal of Fluency Disorders*, 40, 35–43.  
748 <https://doi.org/10.1016/j.jfludis.2014.01.001>
- 749 Crichton-Smith, I. (2002). Communicating in the real world: Accounts from people who  
750 stammer. *Journal of Fluency Disorders*, 27(4), 333–352. [https://doi.org/10.1016/S0094-](https://doi.org/10.1016/S0094-730X(02)00161-4)  
751 [730X\(02\)00161-4](https://doi.org/10.1016/S0094-730X(02)00161-4)
- 752 Dakin, H., Abel, L., Burns, R., & Yang, Y. (2018). Review and critical appraisal of studies  
753 mapping from quality of life or clinical measures to EQ-5D: An online database and  
754 application of the MAPS statement. *Health and Quality of Life Outcomes*, 16(1), 1–9.  
755 <https://doi.org/10.1186/s12955-018-0857-3>
- 756 Daniels, D. E., Gabel, R. M., & Hughes, S. (2012). Recounting the K-12 school experiences  
757 of adults who stutter: A qualitative analysis. *Journal of Fluency Disorders*, 37(2), 71–82.  
758 <https://doi.org/10.1016/j.jfludis.2011.12.001>
- 759 de Sonnevile-Koedoot, C., Bouwmans, C., Franken, M. -C., & Stolk, E. (2015). Economic  
760 evaluation of stuttering treatment in preschool children: The RESTART-study. *Journal of*  
761 *Communication Disorders*, 58, 106–118. <https://doi.org/10.1016/j.jcomdis.2015.10.006>
- 762 Doody, I., Kalinowski, J., Armson, J., & Stuart, A. (1993). Stereotypes of stutterers and  
763 nonstutterers in three rural communities in Newfoundland. *Journal of Fluency Disorders*,  
764 18(4), 363–373. [https://doi.org/10.1016/0094-730X\(93\)90015-V](https://doi.org/10.1016/0094-730X(93)90015-V)
- 765 Douglass, J. E., Schwab, M., & Alvarado, J. (2018). Covert stuttering: Investigation of the  
766 paradigm shift from covertly stuttering to overtly stuttering. *American Journal of Speech-*  
767 *Language Pathology*, 27(3S), 1235–1243. [https://doi.org/10.1044/2018\\_AJSLP-ODC11-](https://doi.org/10.1044/2018_AJSLP-ODC11-17-0190)  
768 [17-0190](https://doi.org/10.1044/2018_AJSLP-ODC11-17-0190)

- 769 Drummond, M., Sculpher, M., Torrance, G., O'Brien, B., & Stoddart, G. L. (2005). *Methods*  
770 *for the economic evaluation of health care programmes* (3rd ed.). Oxford University  
771 Press.
- 772 Erickson, S., & Block, S. (2013). The social and communication impact of stuttering on  
773 adolescents and their families. *Journal of Fluency Disorders*, 38(4), 311–324.  
774 <https://doi.org/10.1016/j.jfludis.2013.09.003>
- 775 Franic, D. M., Bothe, A. K., & Bramlett, R. E. (2012). A welfare economic approach to  
776 measure outcomes in stuttering: comparing willingness to pay and quality adjusted life  
777 years. *Journal of Fluency Disorders*, 37(4), 300–313.  
778 <https://doi.org/10.1016/j.jfludis.2012.04.005>
- 779 Franken, M. -C. (2016, March 18). Personal communication.
- 780 Georges, N. A. (2017). *Perceptions, thoughts, and feelings, of women who stutter in four life*  
781 *domains* (Doctoral dissertation, Walden University).  
782 <https://scholarworks.waldenu.edu/dissertations/4060/>
- 783 Gerlach, H., Totty, E., Subramanian, A., & Zebrowski, P. (2018). Stuttering and labor market  
784 outcomes in the United States. *Journal of Speech, Language, and Hearing*  
785 *Research*, 61(7), 1649–1663. [https://doi.org/10.1044/2018\\_JSLHR-S-17-0353](https://doi.org/10.1044/2018_JSLHR-S-17-0353)
- 786 Gunn, A., Menzies, R. G., O'Brian, S., Onslow, M., Packman, A., Lowe, R., Iverach, L.,  
787 Heard, R., & Block, S. (2014). Axis I anxiety and mental health disorders among  
788 stuttering adolescents. *Journal of Fluency Disorders*, 40, 58–68.  
789 <https://doi.org/https://doi.org/10.1016/j.jfludis.2013.09.002>
- 790 Ham, R. E. (1990). What is stuttering: Variations and stereotypes. *Journal of Fluency*  
791 *Disorders*, 15(5–6), 259–273. [https://doi.org/10.1016/0094-730X\(90\)90041-P](https://doi.org/10.1016/0094-730X(90)90041-P)

- 792 Hayhow, R., Cray, A. M., & Enderby, P. (2002). Stammering and therapy views of people  
793 who stammer. *Journal of Fluency Disorders*, 27(1), 1–17. [https://doi.org/10.1016/S0094-](https://doi.org/10.1016/S0094-730X(01)00102-4)  
794 730X(01)00102-4
- 795 Helgadóttir, F. D., Menzies, R. G., Onslow, M., Packman, A., & O’Brian, S. (2014). A  
796 standalone internet cognitive behavior therapy treatment for social anxiety in adults who  
797 stutter: CBTpsych. *Journal of Fluency Disorders*, 41, 47–54.  
798 <https://doi.org/10.1016/j.jfludis.2014.04.001>
- 799 Horsley, I. A., & Fitzgibbon, C. T. (1987). Stuttering children: Investigation of a stereotype.  
800 *British Journal of Disorders of Communication*, 22(1), 19–35.  
801 <https://doi.org/10.3109/13682828709088686>
- 802 Hugh-Jones, S., & Smith, P. K. (1999). Self-reports of short-and long-term effects of bullying  
803 on children who stammer. *British Journal of Educational Psychology*, 69(2), 141–158.  
804 <https://doi.org/10.1348/000709999157626>
- 805 Iimura, D., & Miyamoto, S. (2022). The influence of stuttering and co-occurring disorders on  
806 job difficulties among adults who stutter. *Speech, Language and Hearing*, 25(2), 235–244.  
807 <https://doi.org/10.1080/2050571X.2020.1852494>
- 808 Iverach, L., Jones, M., McLellan, L., Lyneham, H., Menzies, R., Onslow, M., & Rapee, R.  
809 (2016). Prevalence of anxiety disorders among children who stutter. *Journal of Fluency*  
810 *Disorders*, 49, 13–28. <https://doi.org/10.1016/j.jfludis.2016.07.002>
- 811 Iverach, L., & Rapee, R. M. (2014). Social anxiety disorder and stuttering: Current status and  
812 future directions. *Journal of Fluency Disorders*, 40, 69–82.  
813 <https://doi.org/10.1016/j.jfludis.2013.08.003>
- 814 Jackson, E. S., Yaruss, J. S., Quesal, R. W., Terranova, V., & Whalen, D. H. (2015).  
815 Responses of adults who stutter to the anticipation of stuttering. *Journal of Fluency*  
816 *Disorders*, 45, 38–51. <https://doi.org/10.1016/j.jfludis.2015.05.002>

- 817 Johnson, W. (1934). Stutterers' attitudes toward stuttering. *The Journal of Abnormal and*  
818 *Social Psychology*, 29(1), 32–44. <https://doi.org/10.1037/h0073371>
- 819 Johnson, W. (1961). Measurements of oral reading and speaking rate and disfluency of adult  
820 male and female stutterers and nonstutterers. *Journal of Speech and Hearing Disorders*,  
821 *Monograph Supplement*, 7, 1–20.
- 822 Johnson, W., & Solomon, A. (1937). Studies in the psychology of stuttering: IV. A  
823 quantitative study of expectation of stuttering as a process involving a low degree of  
824 consciousness. *Journal of Speech Disorders*, 2(2), 95–97.  
825 <https://doi.org/10.1044/jshd.0202.95>
- 826 Karimi, H., O'Brian, S., Onslow, M., Jones, M., Menzies, R., & Packman, A. (2013a).  
827 Unscheduled telephone calls to measure percent syllables stuttered during clinical trials.  
828 *Journal of Speech, Language, and Hearing Research*, 56(5), 1455–1461.  
829 [https://doi.org/10.1044/1092-4388\(2013/12-0264\)](https://doi.org/10.1044/1092-4388(2013/12-0264))
- 830 Karimi, H., O'Brian, S., Onslow, M., Jones, M., Menzies, R., & Packman, A. (2013b). Using  
831 statistical process control charts to study stuttering frequency variability during a single  
832 day. *Journal of Speech, Language, and Hearing Research*, 56(6), 1789–1799.  
833 [https://doi.org/10.1044/1092-4388\(2013/12-0328\)](https://doi.org/10.1044/1092-4388(2013/12-0328))
- 834 Klein, J. F., & Hood, S. B. (2004). The impact of stuttering on employment opportunities  
835 and job performance. *Journal of Fluency Disorders*, 29(4), 255–273.  
836 <https://doi.org/10.1016/j.jfludis.2004.08.001>
- 837 Klompas, M., & Ross, E. (2004). Life experiences of people who stutter, and the perceived  
838 impact of stuttering on quality of life: Personal accounts of South African individuals.  
839 *Journal of Fluency Disorders*, 29(4), 275–305.  
840 <https://doi.org/10.1016/j.jfludis.2004.10.001>

- 841 Langevin, M., Packman, A., & Onslow, M. (2010). Parent perceptions of the impact of  
842 stuttering on their preschoolers and themselves. *Journal of Communication Disorders*,  
843 *43*(5), 407–423. <https://doi.org/10.1016/j.jcomdis.2010.05.003>
- 844 Lee A., Van Dulm, O, Robb, M. P., & Ormond, T. (2015). Communication restriction in  
845 adults who stutter. *Clinical Linguistics and Phonetics*, *29*(7), 536–556.  
846 <https://doi.org/10.3109/02699206.2015.1030039>
- 847 Leko Krhen, A., Milić Babić, M., & Horovskij, M. (2021). Everyday challenges and  
848 perspectives of women who stutter. *Hrvatska revija za rehabilitacijska istraživanja*, *57*(1),  
849 85–115. <https://doi.org/10.31299/hrri.57.1.5>
- 850 Li, T., Higgins, J. P. T., & Deeks, J. J. (2021). Chapter 5: Collecting data. 2021 February  
851 2021. In J. P. T. Higgins, J. Thomas, J. Chandler, M. Cumpston, T. Li, M. J., Page, & V.  
852 A. Welch (Eds.). *Cochrane Handbook for Systematic Reviews of Interventions* (version  
853 6.2). Cochrane. [www.training.cochrane.org/handbook](http://www.training.cochrane.org/handbook).
- 854 Martens, C. F., & Engel, D. C. (1986). Measurement of the sound-based word avoidance of  
855 persons who stutter. *Journal of Fluency Disorders*, *11*(3), 241–250.  
856 [https://doi.org/10.1016/0094-730X\(86\)90011-2](https://doi.org/10.1016/0094-730X(86)90011-2)
- 857 Martin, R. R., & Haroldson, S. K. (1967). The relationship between anticipation and  
858 consistency of stuttered words. *Journal of Speech and Hearing Research*, *10*(2), 323–327.  
859 <https://doi.org/10.1044/jshr.1002.323>
- 860 McAllister, J. (2016). Behavioural, emotional and social development of children who stutter.  
861 *Journal of Fluency Disorders*, *50*, 23–32. <https://doi.org/10.1016/j.jfludis.2016.09.003>
- 862 McAllister, J., Collier, J., & Shepstone, L. (2012). The impact of adolescent stuttering on  
863 educational and employment outcomes: Evidence from a birth cohort study. *Journal of*  
864 *Fluency Disorders*, *37*(2), 106–121. <https://doi.org/10.1016/j.jfludis.2012.01.002>

- 865 McAllister, J., Gascoine, S., Carroll, A., Humby, K., Kingston, M., Shepstone, L., Risebro,  
866 H., Mackintosh, B., Davidson Thompson, T., & Hodgekins, J. (2017). Cognitive bias  
867 modification for social anxiety in adults who stutter: a feasibility study of a randomised  
868 controlled trial. *BMJ Open*, 7(10), Article e015601. [http://dx.doi.org/10.1136/bmjopen-](http://dx.doi.org/10.1136/bmjopen-2016-015601)  
869 2016-015601
- 870 McClure, J. A., & Yaruss, J. S. (2003). Stuttering survey suggests success of attitude-  
871 changing treatment. *ASHA Leader*, 8(9). <https://doi.org/10.1044/leader.FTR2.08092003.3>
- 872 Menzies, R. G., O'Brian, S., Onslow, M., Packman, A., St Clare, T., & Block, S. (2008). An  
873 experimental clinical trial of a cognitive-behavior therapy package for chronic stuttering.  
874 *Journal of Speech, Language and Hearing Research*, 51(6), 1451–1464.  
875 [https://doi.org/10.1044/1092-4388\(2008/07-0070](https://doi.org/10.1044/1092-4388(2008/07-0070)
- 876 Menzies, R. G., Onslow, M., Packman, A., & O'Brian, S. (2009). Cognitive behavior therapy  
877 for adults who stutter: A tutorial for speech-language pathologists. *Journal of Fluency*  
878 *Disorders*, 34(3), 187–200. <https://doi.org/10.1016/j.jfludis.2009.09.002>
- 879 Menzies, R. G., Packman, A., Onslow, M., O'Brian, S., Jones, M., & Helgadóttir, F. D.  
880 (2019). In-clinic and standalone Internet cognitive behavior therapy treatment for social  
881 anxiety in stuttering: A randomized trial of iGlebe. *Journal of Speech, Language, and*  
882 *Hearing Research*, 62(6), 1614–1624. [https://doi.org/10.1044/2019\\_JSLHR-S-18-0340](https://doi.org/10.1044/2019_JSLHR-S-18-0340)
- 883 Mihalopoulos, C., Chen, G., Iezzi, A., Khan, M. A., & Richardson, J. (2014). Assessing  
884 outcomes for cost-utility analysis in depression: comparison of five multi-attribute utility  
885 instruments with two depression-specific outcome measures. *The British Journal of*  
886 *Psychiatry*, 205(5), 390–397. <https://doi.org/10.1192/bjp.bp.113.136036>
- 887 Milisen, R. (1938). Frequency of stuttering with anticipation of stuttering controlled. *Journal*  
888 *of Speech Disorders*, 3, 207–214.

- 889 Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G.; PRISMA Group. (2009). Preferred  
890 reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS*  
891 *Medicine*, 6(7), Article e1000097. <https://doi.org/10.1371/journal.pmed.1000097>
- 892 Murphy, B., Quesal, R. W., & Gulker, H. (2007). Covert stuttering. *Perspectives on Fluency*  
893 *and Fluency Disorders*, 17(2), 4–9. <https://doi.org/10.1044/ffd17.2.4>
- 894 Nang, C., Hersh, D., Milton, K., & Lau, S. R. (2018). The impact of stuttering on  
895 development of self-identity, relationships, and quality of life in women who  
896 stutter. *American Journal of Speech-Language Pathology*, 27(3S), 1244–1258.  
897 [https://doi.org/10.1044/2018\\_AJSLP-ODC11-17-0201](https://doi.org/10.1044/2018_AJSLP-ODC11-17-0201)
- 898 Nippold, M. (2019). Language development in children who stutter: A review of recent  
899 research. *International Journal of Speech-Language Pathology*, 21(4), 368–376.  
900 <https://doi.org/10.1080/17549507.2018.1457721>
- 901 Noyes, J., Booth, A., Flemming, K., Garside, R., Harden, A., Lewin, S., Pontoja, T., Hannes,  
902 K., Cargo, M. & Thomas, J. (2018). Cochrane Qualitative and Implementation Methods  
903 Group guidance series–Paper 3: Methods for assessing methodological limitations, data  
904 extraction and synthesis, and confidence in synthesized qualitative findings. *Journal of*  
905 *Clinical Epidemiology*, 97, 49–58. <https://doi.org/10.1016/j.jclinepi.2017.06.020>
- 906 O’Brian, S., Jones, M., Packman, A., Menzies, R., & Onslow, M. (2011). Stuttering severity  
907 and educational attainment. *Journal of Fluency Disorders*, 36(2), 86–92.  
908 <https://doi.org/10.1016/j.jfludis.2011.02.006>
- 909 O’Brian, S., Jones, M., Packman, A., Onslow, M., Menzies, R., Lowe, R., Cream, A., Hearne,  
910 A., Hewat, S., Harrison, E., Block, S., & Briem, A. (2022). The complexity of stuttering  
911 behavior in adults and adolescents: Relationship to age, severity, mental health, impact of  
912 stuttering, and behavioral treatment outcome. *Journal of Speech, Language, and Hearing*  
913 *Research*, 65(7), 2446–2458. [https://doi.org/10.1044/2022\\_JSLHR-21-00452](https://doi.org/10.1044/2022_JSLHR-21-00452)

- 914 Omori, F., Nakagawa, A., Hosogaya, R., Suzuki, M., & Shimizu, E. (2021). A pilot study of a  
915 two-arm, non-randomized controlled trial of guided self-help cognitive behavioral therapy  
916 for stuttering. *Chiba Medical Journal*, 97E(4), 57–70. [https://doi.org/10.20776/S03035476-](https://doi.org/10.20776/S03035476-97E-4-P57)  
917 97E-4-P57
- 918 Packman, A. (2012). Theory and therapy in stuttering: A complex relationship. *Journal of*  
919 *Fluency Disorders*, 37(4), 225–233. <https://doi.org/10.1016/j.jfludis.2012.05.004>
- 920 Palasik, S., Gable, R., Hughes, C., & Rusnak, E. (2012). Perceptions about occupational  
921 experiences by people who stutter. *Perspectives on Fluency and Fluency Disorders*, 22(1),  
922 22–33. <https://doi.org/10.1044/ffd22.1.22>
- 923 Perkins, W. H. (1983). The problem of definition: Commentary on “stuttering.” *Journal of*  
924 *Speech and Hearing Disorders*, 48(3), 246–249. <https://doi.org/10.1044/jshd.4803.246b>
- 925 Plexico, L. W., Hamilton, M. B., Hawkins, H., & Erath, S. (2019). The influence of  
926 workplace discrimination and vigilance on job satisfaction with people who  
927 stutter. *Journal of Fluency Disorders*, 62, Article 105725.  
928 <https://doi.org/10.1016/j.jfludis.2019.105725>
- 929 Porter, H. V. K. (1939). Studies in the psychology of stuttering. XIV. Stuttering phenomena  
930 in relation to size and personnel of audience. *Journal of Speech Disorders*, 4(4), 323–333.  
931 <https://doi.org/10.1044/jshd.0404.323>
- 932 Rees, D. I., & Sabia, J. J. (2014). The kid's speech: The effect of stuttering on human capital  
933 acquisition. *Economics of Education Review*, 38(C), 76–88.  
934 <https://doi.org/10.1016/j.econedurev.2013.07.007>
- 935 Reilly, S., Onslow, M., Packman, A., Wake, M., Bavin, E. L., Prior, M., Eadie, P., Cini, E.,  
936 Bolzonello, C., & Ukoumunne, O. C. (2009). Predicting stuttering onset by the age of 3  
937 years: A prospective, community cohort study. *Pediatrics*, 123(1), 270–277.  
938 <https://doi.org/10.1542/peds.2007-3219>



- 939 Rice, M., & Kroll, R. (1994). A survey of stutterers' perceptions of challenges and  
940 discrimination in the workplace. *Journal of Fluency Disorders*, 19(3), 203.  
941 [https://doi.org/10.1016/0094-730X\(94\)90165-1](https://doi.org/10.1016/0094-730X(94)90165-1)
- 942 Rice, M., & Kroll, R. (2006, October 22). *The impact of stuttering at work: Challenges and*  
943 *discrimination* [Online poster presentation]. International Stuttering Awareness Day  
944 Online Conference 2006, Mankato, USA.  
945 <https://web.mnsu.edu/comdis/isad9/papers/rice9.html>
- 946 Rosenbaum, J. E. (2018). Disabilities and degrees: Identifying health impairments that  
947 predict lower chances of college enrollment and graduation in a nationally representative  
948 sample. *Community College Review*, 46(2), 145–175.  
949 <https://doi.org/10.1177/009155211876263>
- 950 Rowen, D., Rivero-Arias, O., Devlin, N., & Ratcliffe, J. (2020). Review of valuation methods  
951 of preference-based measures of health for economic evaluation in child and adolescent  
952 populations: Where are we now and where are we going? *Pharmacoeconomics*, 38(4),  
953 325–340. <https://doi.org/10.1007/s40273-019-00873-7>
- 954 Siegel, G. M., & Haugen, D. (1964). Audience size and variations in stuttering behavior.  
955 *Journal of Speech and Hearing Research*, 7(4), 381–388.  
956 <https://doi.org/10.1044/jshr.0704.381>
- 957 Silverman, E. M., & Zimmer, C. H. (1982). Demographic characteristics and treatment  
958 experiences of women and men who stutter. *Journal of Fluency Disorders*, 7(2), 273–285.  
959 [https://doi.org/10.1016/0094-730X\(82\)90013-4](https://doi.org/10.1016/0094-730X(82)90013-4)
- 960 Sjøstrand, Å., Kefalianos, E., Hofslundsengen, H., Guttormsen, L. S., Kirmess, M., Lervåg,  
961 A., Hulme, C., & Bottegaard Naess, K. -A. (2021). Non-pharmacological interventions for  
962 stuttering in children six years and younger. *Cochrane Database of Systematic*  
963 *Reviews*, 9(9), Article CD013489. <https://doi.org/10.1002/14651858.CD013489.pub2>

- 964 Sommer, M., Waltersbacher, A., Schlotmann, A., Schröder, H., & Strzelczyk, A. (2021).  
965 Prevalence and therapy rates for stuttering, cluttering, and developmental disorders of  
966 speech and language: evaluation of German health insurance data. *Frontiers in Human*  
967 *Neuroscience*, 15, Article 645292. <https://doi.org/10.3389/fnhum.2021.645292>
- 968 Sønsterud, H., Howells, K., & Ward, D. (2022). Covert and overt stuttering: Concepts and  
969 comparative findings. *Journal of Communication Disorders*, 99, Article 106246.  
970 <https://doi.org/10.1016/j.jcomdis.2022.106246>
- 971 Spencer, E., Packman, A., Onslow, M., & Ferguson, A. (2005). A preliminary investigation  
972 of the impact of stuttering on language use. *Clinical Linguistics and Phonetics*, 19(3),  
973 191–201. <https://doi.org/10.1080/02699200410001698625>
- 974 Spencer, E., Packman, A., Onslow, M., & Ferguson, A. (2009). The effect of stuttering on  
975 communication: A preliminary investigation. *Clinical Linguistics and Phonetics*, 23(7),  
976 473–488. <https://doi.org/10.1080/02699200902833373>
- 977 Steer, M. D., & Johnson, W. (1936). An objective study of the relationship between  
978 psychological factors and the severity of stuttering. *Journal of Abnormal and Social*  
979 *Psychology*, 31(1), 36–46. <https://doi.org/10.1037/h0060643>
- 980 Stein, M. B., Baird, A., & Walker, J. R. (1996). Social phobia in adults with stuttering.  
981 *American Journal of Psychiatry*, 153(2), 278–280. <https://doi.org/10.1176/ajp.153.2.278>
- 982 Sterne, J. A., Hernán, M. A., Reeves, B. C., Savović, J., Berkman, N. D., Viswanathan, M.,  
983 Henry, D., Altman, D. G., Ansari, M. T., Boutron, I., Carpenter, J. R., Chan, A. -W.,  
984 Churchill, R., Deeks, J. J., Hróbjartsson, A., Kirkham, J., Jüni, P., Loke, Y. K., Pigott, T.  
985 D., . . . Higgins, J. P. (2016). ROBINS-I: a tool for assessing risk of bias in non-  
986 randomised studies of interventions. *BMJ*, 355:i4919 <https://doi.org/10.1136/bmj.i4919>
- 987 Sterne, J. A. C., Savović, J., Page, M. J., Elbers, R. G., Blencowe, N. S., Boutron, I., Cates,  
988 C. J., Cheng, H. -Y., Corbett, M. S., Eldridge, S. M., Emberson, J. R., Hernán, M. A.,

- 989 Hopewell, S., Hróbjartsson, A., Junqueira, D. R., Jüni, P., Kirkham, J. J., Lasserson, T., Li,  
990 T., . . . Higgins, J. P. (2019). RoB 2: A revised tool for assessing risk of bias in  
991 randomised trials. *BMJ*, *366*:14898. <https://doi.org/10.1136/bmj.14898>
- 992 Teesson, K., Packman, A., & Onslow, M. (2003). The Lidcombe Behavioral Data Language  
993 of stuttering. *Journal of Speech, Language, and Hearing Research*, *46*(4), 1009–1015.  
994 [https://doi.org/10.1044/1092-4388\(2003/078\)](https://doi.org/10.1044/1092-4388(2003/078))
- 995 Tichenor, S. E., & Yaruss, J. S. (2019). Group experiences and individual differences in  
996 stuttering. *Journal of Speech, Language, and Hearing Research*, *62*(12), 4335–4350.  
997 [https://doi.org/10.1044/2019\\_JSLHR-19-00138](https://doi.org/10.1044/2019_JSLHR-19-00138)
- 998 Tichenor, S. E., & Yaruss, J. S. (2021). Variability of stuttering: Behavior and  
999 impact. *American Journal of Speech-Language Pathology*, *30*(1), 75–88.  
1000 [https://doi.org/10.1044/2020\\_AJSLP-20-00112](https://doi.org/10.1044/2020_AJSLP-20-00112)
- 1001 Ulliana, L., & Ingham, R. J. (1984). Behavioral and nonbehavioral variables in the  
1002 measurement of stutterers' communication attitudes. *Journal of Speech and Hearing*  
1003 *Disorders*, *49*(1), 83–93. <https://doi.org/10.1044/jshd.4901.83>
- 1004 Van Borsel, J., Brepoels, M., & De Coene, J. (2011). Stuttering, attractiveness and romantic  
1005 relationships: The perception of adolescents and young adults. *Journal of Fluency*  
1006 *Disorders*, *36*(1), 41–50. <https://doi.org/10.1016/j.jfludis.2011.01.002>
- 1007 Vanryckeghem, M., Brutton, G. J., Uddin, N., & Van Borsel, J. (2004). A comparative  
1008 investigation of the speech-associated coping responses reported by adults who do and do  
1009 not stutter. *Journal of Fluency Disorders*, *29*(3), 237–250.  
1010 <https://doi.org/10.1016/j.jfludis.2004.07.001>
- 1011 Weiss, A. L., & Zebrowski, P. M. (1994). The narrative productions of children who stutter:  
1012 A preliminary view. *Journal of Fluency Disorders*, *19*(1), 39–63.  
1013 [https://doi.org/10.1016/0094-730X\(94\)90013-2](https://doi.org/10.1016/0094-730X(94)90013-2)

- 1014 Werle, D., & Byrd, C. T. (2022). College professors' perceptions of students who stutter and  
1015 the impact on comfort approaching professors. *Journal of Fluency Disorders*, 67, Article  
1016 105826. <https://doi.org/10.1016/j.jfludis.2020.105826>
- 1017 White, P. A., & Collins, S. R. (1984). Stereotype formation by inference: A possible  
1018 explanation for the "stutterer" stereotype. *Journal of Speech and Hearing Research*, 27(4),  
1019 567–570. <https://doi.org/10.1044/jshr.2704.567>
- 1020 Williams, D. E., Melrose, B. M., & Woods, C. L. (1969). The relationship between stuttering  
1021 and academic achievement in children. *Journal of Communication Disorders*, 2(2), 87–98.  
1022 [https://doi.org/10.1016/0021-9924\(69\)90032-X](https://doi.org/10.1016/0021-9924(69)90032-X)
- 1023 Wilson, I. B., & Cleary, P. D. (1995). Linking clinical variables with health-related quality of  
1024 life. A conceptual model of patient outcomes, *JAMA*, 273(1), 59–65.  
1025 <https://doi:10.1001/jama.1995.03520250075037>
- 1026 Woods, C. L., & Williams, D. E. (1976). Traits attributed to stuttering and normally fluent  
1027 males. *Journal of Speech, Language, and Hearing Research*, 19(2), 267–278.  
1028 <https://doi.org/10.1044/jshr.1902.267>
- 1029 Wu, X. Y., Ohinmaa, A., Johnson, J. A., & Veugelers, P. J. (2014). Assessment of children's  
1030 own health status using visual analogue scale and descriptive system of the EQ-5D-Y:  
1031 Linkage between two systems. *Quality of Life Research*, 23(2), 393–402.  
1032 <https://www.jstor.org/stable/24726538>
- 1033 Yairi, E., & Ambrose, N. (2013). Epidemiology of stuttering: 21st century advances. *Journal*  
1034 *of Fluency Disorders*, 38(2), 66–87. <https://doi.org/10.1016/j.jfludis.2012.11.002>
- 1035 Yairi, E., & Lewis, B. (1984). Disfluencies at the onset of stuttering. *Journal of Speech,*  
1036 *Language, and Hearing Research*, 27(1), 154–159. <https://doi.org/10.1044/jshr.2701.154>

- 1037 Yairi, E., & Williams, D. E. (1970). Speech clinicians' stereotypes of elementary-school boys  
1038 who stutter. *Journal of Communication Disorders*, 3(3), 161–170.  
1039 [https://doi.org/10.1016/0021-9924\(70\)90012-2](https://doi.org/10.1016/0021-9924(70)90012-2)
- 1040 Zhou, T., Guan, H., Wang, L., Zhang, Y., Rui, M., & Ma, A. (2021). Health-related quality of  
1041 life in patients with different diseases measured with the EQ-5D-5L: A systematic  
1042 review. *Frontiers in Public Health*, 9, Article 675523.  
1043 <https://doi.org/10.3389/fpubh.2021.675523>

Figure 2. Results of the quality of life search strategy.




Figure 2. Results of the cost of illness search strategy.




Table 1. Instruments excluded from the review because they do not generate a utility measure

Instrument	Pertinence to quality of life
Rosenberg Self-Esteem Scale	N
Modified Erickson Scale of Communication Attitude	N
Self-Efficacy Scaling for Adult Stutterers	N
Revised Communication Attitude Inventory (S24)	N
Stuttering Severity Instrument (SSI-3 & SSI-4)	N
Perception of Stuttering Inventory	N
Beck Anxiety Inventory	N
Subjective Screening of Stuttering	N
Clinician-based Global Impression	N
Overall Assessment of the Speakers Experience of Stuttering–Quality of Life subscale	Y
Assessment of the Child’s Experience of Stuttering	Y
Self-Assessment Protocol–version for adults	Y
Satisfaction with Life Scale	Y
EuroQol-visual analogue scale (EQ-VAS)	Y
Therapy Outcome Measure	Y
Wright and Ayre Stuttering Self-Rating Profile	Y
Strength and Difficulties Questionnaire	Y
36-Item Short Form Survey (SF-36)	Y
Quality of Life Assessment Schedule	Y
Visual Analog Rating Scale of Health-Related Quality of Life (VAS)	Y



Table 2. Summary of studies included in the quality of life analysis

Author (Year)	Region	Treatment Setting	Study design	Size (N)	Cohort	Male (%)	Population characteristics (stuttering diagnosis, severity, treatment)	Intervention & Comparator	Outcomes (utility measure)	Risk of bias
de Sonnevile-Koedoot et al. (2015)	The Netherlands	Outpatient and home-based	Multi-centre parallel group RCT	199	Children	Lidcombe Program: 69.4 RESTART-DCM: 70.0	Stuttering severity rating $\geq 2$ ('mild') and stuttered at least 3% of syllables.	<u>Two stuttering interventions:</u> Lidcombe Program and RESTART-DCM	HUI3	High
McAllister et al. (2017)	Britain	Online	Two-group parallel design (treatment vs placebo), double-blinded feasibility study.	31	Adults	Treatment: 94 Placebo: 67	Individuals who stutter and have social anxiety disorder.	Cognitive bias modification using neutral/disgusted faces and letter prompts. <u>Intervention:</u> letter replaces neutral face. <u>Comparator:</u> letter replaces neutral or disgusted face	EQ-5D-3L	Low
Omori et al. (2021)	Japan	Outpatient and home-based	Single-centre parallel design (CBT + speech therapy vs speech therapy), non-randomised pilot study.	12	Mixed (age 12 to 65 years)	Treatment: 71.4 Control: 60.0	Meet criteria of childhood-onset fluency disorder (DSM-5) after being diagnosed by an otolaryngologist	<u>Intervention:</u> 7 guided self-hep CBT sessions + 4 speech therapy sessions (fluency shaping method) <u>Control:</u> 4 speech therapy sessions (fluency shaping method)	EQ-5D-5L	High

CBT = Cognitive Behavior Therapy; DCM = Demands and Capacities Model; DSM-5 = Diagnostic and Statistical Manual of Mental Disorders, 5<sup>th</sup> Edition; EQ-5D-3L = EuroQol-5 Dimension, 3 level; HUI3 = Health Utilities Index 3; RCT = Randomised Controlled Trial.

## Supplemental Material S1 – Search Terms

### Search term mapping

Embase subject headings were mapped to subject headings or topics in CINAHL, ProQuest, and Web of Science (Tables S1 and S2).

Title/abstract/keyword searches were used in place of subject heading searches in Scopus, PEDE, NHS EED, and the Cochrane Library, which do not use subject headings or topics to index records. Word variations were searched on all lines of the Cochrane Library search. PEDE, NHS EED, ANZCTR, ClinicalTrials.gov, and OpenGrey had limited search functionality and a small number of results. Searches of these databases were truncated after combining the population search terms and results exported for title/abstract review.

Table S1: Quality of life search term mapping from Embase to CINAHL, ProQuest and Web of Science

Embase 'subject heading'	CINAHL 'subject heading'	ProQuest 'mainsubject'	Web of Science 'topic'
Fluency disorder	Fluency Disorders	No subject heading	Fluency disorder*
Stuttering	No subject heading	Stuttering	Stuttering
Therapy	Therapeutics+	Therapy	Therapy
Clinical outcome	No subject heading	Clinical outcomes	Clinical outcome*
Outcome variable	Dependent Variable	Clinical outcomes	Outcome variable*
Outcome assessment	Outcome Assessment	Clinical outcomes	Outcome assessment*
Treatment outcome	Treatment Outcome+	Clinical outcomes	Treatment outcome*
Patient-reported outcome	Patient-Reported Outcomes+	No subject heading	Patient-reported outcome* OR Patient reported outcome*
Quality of life	Quality of Life+	Quality of life	Quality of life
Quality adjusted life year	Quality-Adjusted Life Years	No subject heading	Quality-adjusted life year* or Quality adjusted life year*

Table S2: Quality of life search term mapping from Embase to CINAHL, ProQuest and Web of Science

Embase 'subject heading'	CINAHL 'subject heading'	ProQuest 'mainsubject'	Web of Science 'topic'
Fluency disorder	Fluency Disorders	No subject heading	Fluency disorder*
Stuttering	No subject heading	Stuttering	Stuttering
Disease burden	No subject heading	No subject heading	Disease burden
Caregiver burden	Caregiver Burden	No subject heading	Caregiver burden
Health care cost	Health Resource Utilization OR Health Care Costs+	Health care expenditures	Health care cost* OR Healthcare cost*
Cost	No subject heading	Cost	Not used <sup>c</sup>
Cost of illness	Economic Aspects of Illness	Cost of illness	Cost of illness
Education	Outcomes of Education <sup>a</sup>	Academic achievement OR Academic underachievement OR Education attainment <sup>b</sup>	Outcomes of education OR Education outcomes OR Academic achievement OR Academic underachievement OR Educational attainment <sup>d</sup>
Occupation	Occupations and Professions+	Occupations	Occupation
Employment	Employment+	Employment	Employment
Attitude	Attitude+	Attitudes	Not used <sup>c</sup>
Personnel management	Personnel Management+	Personnel management	Not used <sup>c</sup>
Mental Health	Mental Health	Mental health	Mental health
Human relation	No subject heading	Human relations	Human relation*
Productivity	Productivity	Productivity	Productivity
Social aspect	No subject heading	No subject heading	Social aspect*

<sup>a</sup> Education+ subject heading expanded; outcomes of education selected from list of sub-headings

<sup>b</sup> Education subject heading expanded; academic achievement, academic underachievement, and education attainment selected from list of sub-headings

<sup>c</sup> Topic returned many irrelevant results. Title/abstract used instead.

<sup>d</sup> Education topic expanded; outcomes of education, education outcomes, academic achievement, academic underachievement, and educational attainment selected as topics, consistent with CINAHL and ProQuest subject headings.

## Quality of Life search terms

Tables S3 to S8 detail the quality of life search terms used for Embase, CINAHL, ProQuest, Web of Science, Scopus, and the Cochrane Library.

Table S3: Quality of Life search terms used in Embase

<b>Databases:</b> Embase (1974 to 2021 January 15), Ovid MEDLINE ® ALL (1946 to January 15, 2021) & APA PsycInfo 1806 to January Week 2 2021		
<b>Search date:</b> 20-Jan-21		
#	Search terms	Results
1	fluency disorder/ or stuttering/	13,680
2	(stuttering or stammering or disfluency).kw. or (stuttering or stammering or disfluency).tw.	14,504
3	1 or 2	17,350
4	intervention.kw. or intervention.tw.	1,844,627
5	treat*.kw. or treat*.tw.	14,205,344
6	therapy/	1,328,181
7	4 or 5 or 6	16,037,827
8	clinical outcome/ or outcome variable/ or outcome assessment/ or treatment outcome/ or patient-reported outcome/	2,602,676
9	quality of life/	737,668
10	quality adjusted life year/	40,878
11	health related outcome*.tw. or health related outcome*.kw.	5,267
12	8 or 9 or 10 or 11	3,235,601
13	3 and 7 and 12	751
14	remove duplicates from 13	516

Explanatory notes: / = subject heading; \* = unlimited truncation (i.e., unlimited suffix variations); (...) = limiter; kw = keyword heading; tw = text word (includes title, abstract, and drug trade name).

Table S4: Quality of Life search terms used in CINAHL

<b>Databases:</b> CINAHL Complete		
<b>Search date:</b> 22-Feb-21		
#	Search terms	Results

1	(MH "Fluency Disorders")	3,166
2	TI stuttering or fluency disorder or stammering or dysfluency	1,486
3	AB stuttering or fluency disorder or stammering or dysfluency	1,487
4	1 or 2 or 3	3,424
5	TI intervention or AB intervention	424,759
6	TI treat* or AB treat*	1,080,074
7	(MH "Therapeutics+")	1,585,326
8	5 or 6 or 7	2,541,569
9	TI clinical outcome or AB clinical outcome	75,289
10	(MH "Dependent Variable")	2,957
11	(MH "Outcome Assessment") or (MH "Treatment Outcomes+") or (MH "Patient-Reported Outcomes+")	428,083
12	(MH "Quality of Life+")	125,305
13	(MH "Quality-Adjusted Life Years")	5,025
14	TI health related outcome* or AB health related outcome*	5,087
15	9 or 10 or 11 or 12 or 13 or 14	587,035
16	4 and 8 and 15	314

Explanatory notes: \* = truncation wildcard (i.e., unlimited suffix variations); + = explode subject heading; (...) = limiter; "... " = exact phrase; AB = abstract; MH = exact subject heading; TI = title.

Table S5: Quality of Life search terms used in ProQuest

<b>Databases:</b> ProQuest		
<b>Search date:</b> 26-Feb-21		
#	Search terms	Results
1	MAINSUBJECT.EXACT("stuttering")	6,150
2	AB(stuttering or stammering or disfluency) or TI(stuttering or stammering or disfluency)	13,878
3	1 or 2	15,128
4	AB(intervention) or TI(intervention)	2,054,834
5	AB(treat*) or ti(treat*)	12,202,146
6	MAINSUBJECT.EXACT("therapy")	77,618
7	4 or 5 or 6	13,696,514
8	MAINSUBJECT.EXACT("Clinical outcomes")	47,463
9	AB(patient reported outcome) or TI(patient reported outcome)	240,347

10	MAINSUBJECT.EXACT("Quality of life")	354,124
11	AB(quality adjusted life year) or TI(quality adjusted life year)	28,738
12	AB(health related outcome) or TI(health related outcome)	152,245
13	8 or 9 or 10 or 11 or 12	744,537
14	3 and 7 and 13	123

Explanatory notes: \* = truncation (up to 500 word variations); (...) = limiter; “...” = exact phrase; AB = abstract; MAINSUBJECT.EXACT = exact subject heading; TI = title.

Table S6: Quality of Life search terms used in Web of Science

<b>Database:</b> Web of Science (All years, 1864-2021)		
<b>Search date:</b> 27-Feb-21		
#	Search terms	Results
1	TS=("fluency disorder*") or TS=(stuttering)	9,146
2	TI=(stuttering or stammering or disfluency) or AB=(stuttering or stammering or disfluency)	9,293
3	1 or 2	10,434
4	TI=(intervention) or AB=(intervention)	1,441,805
5	TI=(treat*) or AB=(treat*)	9,534,163
6	TS=(therapy)	11,596,190
7	4 or 5 or 6	17,618,885
8	TS=("clinical outcome*" or "outcome variable*" or "outcome* assessment*" or "treatment outcome*" or "patient-reported outcome*" or "patient reported outcome*")	1,382,075
9	TS=("quality of life")	604,326
10	TS=("quality adjusted life year*" or "quality-adjusted life year*")	21,227
11	TI=("health related outcome*") or AB=("health related outcome*")	2,350
12	8 or 9 or 10 or 11	1,895,176
13	3 and 7 and 12	496

Explanatory notes: \* = wildcard for right hand truncation (any group of suffix characters, including no character); (...) = limiter to override operator precedence; “...” = exact phrase; AB = abstract; TI = title; TS = topic.

Table S7: Quality of Life search terms used in Scopus

<b>Database:</b> Scopus (All years)
-------------------------------------

<b>Search date:</b> 02-Mar-21		
#	<b>Search terms</b>	<b>Results</b>
1	TITLE-ABS-KEY ("fluency disorder*" or stuttering)	8,230
2	TITLE-ABS (stuttering OR stammering or disfluency)	7,421
3	1 or 2	9,469
4	TITLE-ABS (intervention)	1,360,735
5	TITLE-ABS (treat*)	8,598,971
6	TITLE-ABS-KEY (therapy)	4,659,707
7	4 or 5 or 6	11,885,688
8	TITLE-ABS-KEY ("clinical outcome*" or "outcome variable*" or "outcome* assessment*" or "treatment outcome*" or "patient-reported outcome*" or "patient reported outcome*")	2,015,330
9	TITLE-ABS-KEY ("quality of life")	539,489
10	TITLE-ABS-KEY ("quality adjusted life year" or "quality-adjusted life year")	25,642
11	TITLE-ABS ("health related outcome*")	2,207
12	8 or 9 or 10 or 11	2,245,159
13	3 and 7 and 12	616

Explanatory notes: \* = wildcard; (...) = limiter; "... " = loose phrase (exact words, ignores punctuation); TITLE-ABS = article title or abstract; TITLE-ABS-KEY = article title or abstract or keywords.

Table S8: Quality of Life search terms used in the Cochrane Library

<b>Database:</b> Cochrane Library (including Cochrane Database of Systematic Reviews and Cochrane Central Register of Controlled Trials)		
<b>Search date:</b> 20-May-21		
#	<b>Search terms</b>	<b>Results</b>
1	("fluency disorder" or stuttering):ti,ab,kw	276
2	(stuttering or stammering or disfluency):ti or (stuttering or stammering or disfluency):ab	271
3	1 or 2	291
4	intervention:ti,ab	422,259
5	treat*:ti,ab	806,531
6	therapy:ti,ab,kw	710,888
7	4 or 5 or 6	1,212,754



8	("clinical outcome*" or "outcome variable*" or "outcome* assessment*" or "treatment outcome*" or "patient-reported outcome*" or "patient reported outcome*"):ti,ab,kw	267,756
9	"quality of life":ti,ab,kw	115,018
10	("quality adjusted life year" or "quality-adjusted life year"):ti,ab,kw	0
11	("health related outcome*"):ti,ab	429
12	8 or 9 or 10 or 11	0
13	8 or 9 or 11	348,994
14	3 and 7 and 12	0
15	3 and 7 and 13	69

Explanatory notes: \* = wildcard to search zero or more characters; (...) = nesting to change search precedence; “...” = phrase search; ab = abstract; kw = keyword; ti = title.

### Cost of illness search terms

Tables S9 to S14 detail the cost search terms used for Embase, CINAHL, ProQuest, Web of Science, Scopus, and the Cochrane Library.

Table S9: Cost search terms used in Embase

<b>Databases:</b> Embase (1974 to 2021 January 15), Ovid MEDLINE ® ALL (1946 to January 15, 2021) & APA PsycInfo 1806 to January Week 2 2021		
<b>Search date:</b> 19-Jan-21		
<b>#</b>	<b>Search terms</b>	<b>Results</b>
1	fluency disorder/ or stuttering/	13,680
2	(stuttering or stammering or disfluency).kw. or (stuttering or stammering or disfluency).tw.	14,504
3	1 or 2	17,350
4	economic.kw. or economic.tw.	614,353
5	health resource*.kw. or health resource*.tw.	18,082
6	disease burden/ or caregiver burden/	36,056
7	economic burden.kw. or economic burden.tw.	32,765
8	emotional burden.kw. or emotional burden.tw.	2,482
9	“health care cost”/	245,531
10	"cost”/	108,319

11	cost.kw. or cost.tw.	1,126,791
12	"cost of illness"/	47,767
13	education/	480,763
14	occupation/ or employment/	187,403
15	attitude/ or personnel management/	199,338
16	mental health/	256,431
17	human relation/	89,318
18	productivity/	57,123
19	social aspect/	78,028
20	4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19	3,045,646
21	3 and 20	662
22	Remove duplicates from 21	518

Explanatory notes: / = subject heading; \* = unlimited truncation (i.e., unlimited suffix variations); (...) = limiter; "... " = literal string (i.e., an exact phrase); kw = keyword heading; tw = text word (includes title, abstract, and drug trade name).

Table S10: Cost search terms used in CINAHL

<b>Database:</b> CINAHL		
<b>Search date:</b> 22-Feb-21		
#	Search terms	Results
1	(MH "Fluency Disorders")	3,166
2	TI stuttering or fluency disorder or stammering or dysfluency	1,486
3	AB stuttering or fluency disorder or stammering or dysfluency	1,487
4	1 or 2 or 3	3,424
5	TI economic or AB economic	70,268
6	(MH "Health Resource Utilization")	19,592
7	TI health resource* or AB health resource*	18,217
8	TI disease burden or AB disease burden	14,159
9	(MH "Caregiver Burden")	10,172
10	TI caregiver burden or AB caregiver burden	4,247
11	TI economic burden or AB economic burden	4,958
12	TI emotional burden or AB emotional burden	987
13	(MH "Health Care Costs+")	61,717
14	TI cost or AB cost	163,546

15	(MH "Economic Aspects of Illness")	10,072
16	(MH "Education+")	948,231
17	(MH "Outcomes of Education")	14,151
18	(MH "Occupations and Professions+") or (MH "Employment+")	143,243
19	(MH "Attitude+") or (MH "Personnel Management+")	715,705
20	(MH "Mental Health")	41,151
21	TI human relation or AB human relation	969
22	(MH "Productivity")	11,248
23	TI social aspect OR AB social aspect	4,499
24	5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 18 or 19 or 20 or 21 or 22 or 23	1,741,435
25	4 and 24	1,392
26	5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 17 or 18 or 19 or 20 or 21 or 22 or 23	1,103,452
27	4 and 26	534

Explanatory notes: \* = truncation wildcard (i.e., unlimited suffix variations); + = explode subject heading; (...) = limiter; "...” = exact phrase; AB = abstract; MH = exact subject heading; TI = title.

Table S11: Cost search terms used in ProQuest

<b>Database:</b> ProQuest		
<b>Search date:</b> 26-Feb-21		
#	Search terms	Results
1	MAINSUBJECT.EXACT("stuttering")	6,150
2	AB(stuttering or stammering or disfluency) or TI(stuttering or stammering or disfluency)	13,878
3	1 or 2	15,128
4	AB(economics) or TI(economics)	368,891
5	AB(health resource) or TI(health resource)	307,910
6	AB(disease burden) or TI(disease burden) or AB(caregiver burden) or TI(caregiver burden)	185,051
7	AB(economic burden) or TI(economic burden)	56,629
8	AB(emotional burden) or TI(emotional burden)	9,914
9	MAINSUBJECT.EXACT("health care expenditures")	23,951
10	MAINSUBJECT.EXACT("cost")	17,285
11	AB(cost) or TI(cost)	4,449,850

12	MAINSUBJECT.EXACT("cost of illness")	30,643
13	MAINSUBJECT(education)	3,101,730
14	MAINSUBJECT.EXACT("Academic achievement") or MAINSUBJECT.EXACT("Academic underachievement") or MAINSUBJECT.EXACT("Educational attainment")	156,203
15	MAINSUBJECT(Occupations) or MAINSUBJECT(Employment)	617,811
16	MAINSUBJECT.EXACT("attitudes") or MAINSUBJECT.EXACT("personnel management")	163,449
17	MAINSUBJECT.EXACT("mental health")	246,652
18	MAINSUBJECT.EXACT("human relations")	15,322
19	MAINSUBJECT.EXACT("productivity")	234,137
20	AB(social aspect) or TI(social aspect)	72,028
21	4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 15 or 16 or 17 or 18 or 19 or 20	10,258,536
22	3 and 21	818
23	4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 16 or 17 or 18 or 19 or 20	7,603,501
24	3 and 23	449

Explanatory notes: \* = truncation (up to 500 word variations); (...) = limiter; “...” = exact phrase; AB = abstract; MAINSUBJECT.EXACT = exact subject heading; TI = title.

Table S12: Cost search terms used in Web of Science

<b>Database:</b> Web of Science (All years, 1864-2021)		
<b>Search date:</b> 27-Feb-21		
<b>#</b>	<b>Search terms</b>	<b>Results</b>
1	TS=("fluency disorder*") or TS=(stuttering)	9,146
2	TI=(stuttering or stammering or disfluency) or AB=(stuttering or stammering or disfluency)	9,293
3	1 or 2	10,434
4	TI=(economic) or AB=(economic)	1,314,049
5	TI=("health resource*") or AB=("health resource*")	8,302
6	TS=("disease burden" or "caregiver burden")	29,506
7	TI=("economic burden") or AB=("economic burden")	16,123
8	TI=("emotional burden") or AB=("emotional burden")	1,046
9	TS=("health care cost*" or "healthcare cost*")	80,388
10	TI=("cost") or AB=("cost")	1,921,726

11	TS=("cost of illness")	30,415
12	TS=("outcomes of education" or "education outcomes" or "academic achievement" or "academic underachievement" or "educational attainment")	48,068
13	TS=(occupation or employment)	460,611
14	TI=(attitude or "personnel management") or AB=(attitude or "personnel management")	493,853
15	TS=(mental health)	580,975
16	TS=("human relation*")	9,504
17	TS=(productivity)	548,342
18	TS=("social aspect*")	60,479
19	4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18	5,046,284
20	3 and 19	807

Explanatory notes: \* = wildcard for right hand truncation (any group of suffix characters, including no character); (...) = limiter to override operator precedence; “...” = exact phrase; AB = abstract; TI = title; TS = topic.

Table S13: Cost search terms used in Scopus

<b>Database:</b> Scopus (All years)		
<b>Search date:</b> 02-Mar-21		
#	Search terms	Results
1	TITLE-ABS-KEY ("fluency disorder*" or stuttering)	8,230
2	TITLE-ABS (stuttering OR stammering or disfluency)	7,421
3	1 or 2	9,469
4	TITLE-ABS (economic)	1,335,483
5	TITLE-ABS ("health resource*")	7,926
6	TITLE-ABS-KEY ("disease burden" or "caregiver burden")	43,876
7	TITLE-ABS ("economic burden")	14,870
8	TITLE-ABS ("emotional burden")	1,168
9	TITLE-ABS-KEY ("health care cost*" or "healthcare cost*")	209,210
10	TITLE-ABS ("cost")	2,482,855
11	TITLE-ABS-KEY ("cost of illness")	34,851
12	TITLE-ABS-KEY ("outcomes of education" or "education outcomes" or "academic achievement" or "academic underachievement" or "educational attainment")	73,056

13	TITLE-ABS-KEY (occupation or employment)	443,479
14	TITLE-ABS (attitude or "personnel management")	460,999
15	TITLE-ABS-KEY (mental and health)	484,748
16	TITLE-ABS ("human relation*")	7,505
17	TITLE-ABS-KEY (productivity)	412,547
18	TITLE-ABS-KEY ("social aspect*")	115,313
19	4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18	5,364,509
20	3 and 19	761

Explanatory notes: \* = wildcard; (...) = limiter; "... " = loose phrase (exact words, ignores punctuation); TITLE-ABS = abstract title or abstract; TITLE-ABS-KEY = article title or abstract or keywords.

Table S14: Cost search terms used in the Cochrane Library

<b>Database:</b> Cochrane Library (including Cochrane Database of Systematic Reviews and Cochrane Central Register of Controlled Trials)		
<b>Search date:</b> 12-Mar-21		
#	Search terms	Results
1	("fluency disorder" or stuttering):ti,ab,kw	272
2	(stuttering or stammering or disfluency):ti or (stuttering or stammering or disfluency):ab	268
3	1 or 2	287
4	(economic):ti or (economic):ab	18,767
5	("health resource"):ti or ("health resource"):ab	762
6	("disease burden" or "caregiver burden"):ti,ab,kw	3,110
7	("economic burden"):ti or ("economic burden"):ab	1,050
8	("emotional burden"):ti or ("emotional burden"):ab	100
9	("health care cost" or "healthcare cost"):ti,ab,kw	9,037
10	("cost"):ti,ab	69,665
11	("cost of illness"):ti,ab,kw	1,115
12	("outcomes of education" or "education outcomes" or "academic achievement" or "academic underachievement" or "educational attainment"):ti,ab,kw	2,231
13	(occupation or employment):ti,ab,kw	34,867
14	(attitude or "personnel management"):ti,ab	12,689
15	("mental health"):ti,ab,kw	20,867

16	("human relation"):ti,ab	30
17	(productivity):ti,ab,kw	82,265
18	("social aspect"):ti,ab,kw	762
19	4 or 5 or 6 or 7 or 8 or 9 or 10 11 or 12 or 13 or 14 or 15 or 16 or 17 18	212,907
20	3 and 19	77

Explanatory notes: (...) = nesting to change search precedence; “...” = phrase search; ab = abstract; kw = keyword; ti = title.

### Search terms using population only

Tables S15 to S19 detail the truncated search terms used for the Paediatric Economic Database Evaluation (PEDE), the National Health Service Economic Evaluation Database (NHS EED), the Australian New Zealand clinical trials registry (ANZCTR), United States National Institutes of Health trial register, and Open Grey.

Table S15: Search terms used in the Paediatric Economic Database Evaluation (PEDE) database

<b>Database:</b> PEDE (All years, 1980 - 2019)		
<b>Search date:</b> 02-Mar-21		
#	Search terms	Results
1	TITLE_ABSTRACT_KEYWORDS (stuttering)	1
2	TITLE_ABSTRACT_KEYWORDS (stammering)	0
3	TITLE_ABSTRACT_KEYWORDS (disfluency)	0
4	TITLE_ABSTRACT_KEYWORDS ("fluency disorder")	0
5	1 or 2 or 3 or 4	1

Explanatory notes: (...) = limiter; “...” = phrase search.

Table S16: Search terms used in the National Health Service Economic Evaluation Database (NHS EED)

<b>Database:</b> NHS EED (All years, 1994 - 2015)		
<b>Search date:</b> 02-Mar-21		

#	Search terms	Results
1	Any field (stuttering or stammering or disfluency)	13
2	Any field ("fluency disorder")	0
3	1 or 2	13

Explanatory notes: (...) = limiter; "... " = phrase search.

Table S17: Search terms used in the Australian New Zealand clinical trials registry (ANZCTR)

<b>Database:</b> ANZCTR		
<b>Search date:</b> 20-May-21		
#	Search terms	Results
1	Health condition(s) or problem(s) studied: (stuttering)	10
2	Health condition(s) or problem(s) studied: (stammering)	0
3	Health condition(s) or problem(s) studied: (disfluency)	1
4	Health condition(s) or problem(s) studied: ("fluency disorder")	0
5	1 or 2 or 3 or 4	10

Explanatory notes: (...) = limiter; "... " = phrase search.

Table S18: Search terms used in the United States National Institutes of Health trial register (ClinicalTrials.gov)

<b>Database:</b> United States National Institutes of Health trial register (ClinicalTrials.gov)		
<b>Search date:</b> 20-May-21		
#	Search terms	Results
1	Condition or disease: (stuttering or stutters or stammering)	34
2	Condition or disease: (disfluency)	3
3	Condition or disease: ("fluency disorder")	14
4	1 or 2 or 3	45

Explanatory notes: (...) = limiter; "... " = phrase search.

Table S19: Search terms used in OpenGrey

<b>Database:</b> OpenGrey
---------------------------



<b>Search date: 20-May-21</b>		
<b>#</b>	<b>Search terms</b>	<b>Results</b>
1	stuttering	6
2	stammering	6
3	disfluency	12
4	fluency disorder	5
5	1 or 2 or 3 or 4	28

## Supplemental Material S2.

### *Data extraction form*

The data extraction form was developed, based on the Cochrane checklist of items to consider in data collection (Li et al., 2021). Relevant data from each of the included studies was extracted and recorded in Microsoft Access.

<b>1. Study design</b>
Study aim
Study design (e.g., cluster, cross-over)
Recruitment and sampling
Was there randomization? If yes, how was it achieved? If no, how were groups allocated?
Was there blinding?
Sequence generation and concealment
Incomplete outcome data/selective outcome reporting?
Enrolment start/end
Duration of follow-up
<b>2. Population</b>
Inclusion criteria
Exclusion criteria
Total number (N)
Population description (including diagnostic characteristics & disease severity)
Setting
Region/countries involved
Baseline characteristics Age Sex Comorbidities Socio-economic status Ethnicity
<b>3. Intervention and comparator</b>
Number of intervention groups
Intervention/comparator description Components Delivery method Timing/intensity/frequency Duration of treatment
Implementation description Format and content Staffing and equipment
Intervention/comparator integrity/fidelity/compliance
Co-interventions (for intervention and comparator arm)
<b>4. Utility measures outcomes</b>
<b>Overview</b>
Measurement instrument (e.g., CHU9D) Name of instrument Unit of measurement Upper/lower limits

Whether high or low is favourable
Clinical thresholds
Relevance to study aim
Metric (e.g., change pre/post intervention)
Timing of outcome measurements
Method of collection
Method of aggregation (e.g., mean & SD; proportion with condition)
Between group estimate (e.g., RR, OR, mean difference)
<b>At each time point</b>
Description (e.g., baseline, 6mo, 12mo)
Number of participants:
In each intervention
Censored (withdrawn/lost to follow-up)
Result - summary data (e.g., mean & SD)
Estimate of effect (effect size, 95%CI, <i>p</i> value)
<b>Summary/conclusion</b>
Is there any variation in measurement or reporting of the outcome?
Content available/appropriate for meta-analysis?
Were subgroup analyses conducted?
<b>5. Outcomes - cost</b>
Analytic perspective (e.g., health system)
Time horizon
Setting (i.e., study country)
Context and relevance
Cost items included, grouped as:
Health sector
Other sector (e.g., education)
Patient and family
Productivity impacts
Resource use (e.g., number of speech path visits)
Unit costs (currency, price year) (e.g., cost per speech path visit)
Discount rate
<b>6. Miscellaneous</b>
Key conclusions & limitations noted by the study authors
Miscellaneous comments from study authors
References to relevant studies
Generalisability of the results
Funding source
Conflicts of interest
Correspondence with study authors required?
Study quality notes (ethical approval, sample size calculation)

### Supplemental Material S3.

Details of the risk of bias for the quality of life and cost studies using (i) the Rob 2 tool (Sterne et al., 2019) for randomized trials, (ii) the ROBINS-I tool (Sterne et al., 2016) for non-randomized studies, and (iii) the CASP qualitative studies checklist (Critical Appraisal Skills Programme, 2018) for qualitative research.

#### (i) Randomized trials

	Randomization	Deviations from intended interventions	Missing outcome data	Measurement of outcomes	Selection of reported result	Overall score
de Sonneville-Koedoot et al. (2015)	High	Low	Low	Some concerns	Low	High
McAllister et al. (2017)	Low	Low	Low	Low	Low	Low

#### (ii) Non-randomized trials

	Confounding	Participant selection	Classification of interventions	Deviations from intended intervention	Missing data	Measurement of outcomes	Selection of reported result	Overall score
Berchiatti et al. (2020)	High	Low	Not Applicable	Not Applicable	Low	Low	Low	High
Blumgart et al. (2010a)	Moderate	Low	Not Applicable	Not Applicable	Low	Moderate	Low	Moderate
Boulet et al. (2009)	Moderate	Low	Not Applicable	Not Applicable	Moderate	Low	Low	Moderate
Boyle (2018)	High	Low	Not Applicable	Not Applicable	High	Low	Low	High

Boyle et al. (1994)	High	Low	Not Applicable	Not Applicable	Moderate	Low	Low	High
Calnan and Richardson (1977)	High	Low	Not Applicable	Not Applicable	Low	Low	Low	High
Craig and Calver (1991)	High	Low	Low	Low	High	Low	Low	High
Erickson and Block (2013)	High	Low	Not Applicable	Not Applicable	Low	Low	Low	High
Franic et al. (2012)	High	Low	Not Applicable	Not Applicable	Moderate	Low	Low	High
Gerlach et al. (2018)	Moderate	Low	Not Applicable	Not Applicable	No Information	Low	Low	Moderate
Hayhow et al. (2002)	High	Low	Not Applicable	Not Applicable	Low	Low	Low	High
Hugh-Jones and Smith (1999)	High	Low	Not Applicable	Not Applicable	Low	Low	Low	High
Imura and Miyamoto (2022)	High	Low	Not Applicable	Not Applicable	Moderate	Low	Low	High
Klein and Hood (2004)	High	Low	Not Applicable	Not Applicable	No Information	Low	Low	High
McAllister et al. (2012)	Moderate	Low	Not Applicable	Not Applicable	Moderate	Low	Low	Moderate
McClure and Yarus (2003)	High	Low	Not Applicable	Not Applicable	No Information	Low	High	Critical
O'Brian et al. (2011)	High	Low	Not Applicable	Not Applicable	Low	Low	Low	High

Omori et al. (2021)	High	Low	Low	Moderate	High	High	Low	High
Palasik (2012)	High	Low	Not Applicable	Not Applicable	No Information	Low	Low	High
Plexico et al. (2019)	High	Low	Not Applicable	Not Applicable	Low	Low	Low	High
Rees and Sabia (2014)	Moderate	Low	Not Applicable	Not Applicable	Moderate	Low	Low	Moderate
Rice (1994)	High	No Information	Not Applicable	Not Applicable	No Information	Low	High	Critical
Rice and Kroll (2006)	High	Low	Not Applicable	Not Applicable	No Information	Low	Low	High
Rosenbaum (2018)	Moderate	Low	Not Applicable	Not Applicable	Moderate	Low	Low	Moderate
Sommer et al. (2021)	High	Low	Not Applicable	Not Applicable	Low	Low	Low	High
Werle and Byrd (2022)	Moderate	Low	Not Applicable	Not Applicable	Moderate	Low	Low	Moderate
Williams et al. (1969)	Moderate	Low	Not Applicable	Not Applicable	No Information	Low	Low	Moderate

(iii) Qualitative research

	Clear research aim	Appropriate methodology	Appropriate design	Appropriate recruitment	Appropriate data selection	Researcher-participant relationship considered	Ethical issues considered	Rigorous data analysis	Statement of findings	Valuable research	Overall
Beilby et al. (2013)	Yes	Yes	Yes	Unclear	Yes	Unclear	Yes	Yes	Yes	Yes	Lower

Boberg and Boberg (1990)	Yes	Yes	Unclear	Unclear	Yes	Unclear	No	Unclear	Yes	Yes	Yes	Yes	Higher
Bricker-Katz et al. (2013)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Lower
Butler (2013)	Yes	Yes	Yes	Yes	Yes	Unclear	Yes	Unclear	Yes	Yes	Yes	Yes	Lower
Butler (2014)	Yes	Yes	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Lower
Crichton-Smith (2002)	Yes	Yes	Yes	Yes	Yes	Unclear	Yes	Unclear	Yes	Yes	Yes	Yes	Lower
Georges (2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Lower
Johnson (1934)	Yes	Yes	Yes	Unclear	Unclear	Unclear	No	No	Unclear	Yes	Yes	Yes	Higher
Leko Krhen et al. (2021)	Yes	Yes	Unclear	No	Yes	No	Unclear	No	Yes	Yes	Yes	Yes	Higher
Nang et al. (2018)	Yes	Yes	Unclear	Yes	Yes	Unclear	Yes	Unclear	Yes	Yes	Yes	Yes	Lower
Silverman and Zimmer (1982)	Yes	Yes	Unclear	No	Unclear	No	No	No	Unclear	Yes	Yes	Yes	Higher

## Supplemental Material S4.

Summary of studies included in the cost of illness analysis.

Author (Year)	Country	Study design	Size (N)	Cohort	Male (%)	Population characteristics (stuttering diagnosis, severity, treatment)	Cost outcomes	Risk of bias
<b>Randomized controlled trials</b>								
de Sonnevile-Koedoot et al. (2015)	The Netherlands	Multi-centre parallel group RCT	199	Children	Lidcombe Program: 69.4 RESTART-DCM: 70.0	Stuttering severity rating $\geq 2$ ('mild') and stuttered at least 3% of syllables.	Health care utilisation, direct and indirect costs.	High
McAllister et al. (2017)	Britain	Two-group parallel design (treatment vs placebo), double-blinded feasibility study.	31	Adults	Treatment: 94 Placebo: 67	Individuals who stutter and have social anxiety disorder.	Health care utilisation, direct costs	Low
<b>Non-randomized studies</b>								
Berchiatti et al. (2020)	Italy	Case-control	572 (CWS $n = 62$ , CWNS $n = 474$ , Teachers $n = 36$ )	Children	CWS: 58.1 CWNS: 49.2	Stuttering diagnosis made by speech therapist in medical centres. CWS had prior/current formal therapy.	Education (teacher relationships; academic performance), social (friendships)	High
Blumgart et al. (2010a)	Australia	Cross-sectional <sup>a</sup>	200	Adults	75.5	Mean (SD) SS = 3.7 (2.8). Mean (SD) perceived stuttering severity = 4.2 (2). 94% had sought prior treatment.	Direct and indirect costs, employment	Moderate



Author (Year)	Country	Study design	Size (N)	Cohort	Male (%)	Population characteristics (stuttering diagnosis, severity, treatment)	Cost outcomes	Risk of bias
Boulet et al. (2009)	USA	Cross-sectional	95,132 (CWS n = 1,530)	Children	71.4	Parent/ guardian report based on clinical diagnosis	Health care utilisation, education (special services/early intervention)	Moderate
Boyle (2018)	USA	Cross-sectional	324	Adults	67	Self-diagnosed PWS. 95% had prior therapy	Social (discrimination), employment (recruitment)	High
Boyle et al. (1994)	USA	Cross-sectional	2,779 (CWS n = 297)	Children with disabilities	Not reported	Parent/ guardian report	Health care utilisation, education (school attendance and performance)	High
Calnan and Richardson (1977)	England, Scotland, and Wales	Cross-sectional	11,455 (CWS n = 65)	Children	Not reported	Stuttering 'diagnosed' in three ways: clinician assessment, teacher assessment, speech test. Results based on different diagnosis methods	Education (performance)	High
Craig and Calver (1991)	Australia	Study II: 10-month follow-up of 2 non-randomized experimental trials	62	Adults	Not reported	Craig 1984: AWS Craig 1985: AWS, mean %SS = 12.9 All participants were successfully treated with smooth speech prior to 10-month follow-up.	Employment (promotion, career improvement)	High
Erickson and Block (2013)	Australia	Cross-sectional	36	Adolescents	77.8	Mean (SD) stuttering frequency 6.9% (5.7%) SS, range = 2.1–26.8% Mean (SD) onset: 8.5 (3.0). Previously received treatment: 30 (84%). Major treatment type: speech restructuring (63% of those treated).	Education (avoid school, affected schoolwork), social (teasing/bullying, exclusion), indirect costs	High

Author (Year)	Country	Study design	Size (N)	Cohort	Male (%)	Population characteristics (stuttering diagnosis, severity, treatment)	Cost outcomes	Risk of bias
Francic et al. (2012)	USA	Cross-sectional	80 (included <i>n</i> = 78, history of stuttering <i>n</i> = 4)	Adults	17.9%	Not discussed	Direct and indirect costs (willingness to pay)	High
Gerlach et al. (2018)	USA	Cohort	20,745 (included <i>n</i> = 13,564, PWS <i>n</i> = 261)	Stuttered in adolescence, outcomes as adults	PWS: 64.8 PWNS: 47.5	PWS: 84% of respondents described their stuttering as "mild"; 15% as "moderate"; and less than 1% as "severe."	Employment (earnings, employment status)	Moderate
Hayhow et al. (2002)	UK	Cross-sectional	332	Adolescents and adults (age range 16–86 years)	71	Self-reported severity (10-point scale, 1=mild and 10=severe). 264 (80%) rating their stammering as "mild" (1–3) on a good day. Bad day 8% (1–3), 15% (4–5), 32% (6–7), 45% (8–10)	Education (performance), employment (occupation choice, promotion), social (friendships, romantic partners)	High
Hugh-Jones and Smith (1999)	UK	Cross-sectional	276	Adolescents or adults	75.7	Not discussed	Education (attendance, performance), employment (performance), social (relationships)	High
Imura and Miyamoto (2022)	Japan	Cross-sectional	112 (included <i>n</i> = 110, AWS without comorbidity <i>n</i> = 52)	Adults	82.7	Participates in self-help group: 5% no, 18% rarely, 30% sometimes, 35% often, 12% almost always.	Employment (self-rated job difficulties)	High
Klein and Hood (2004)	USA	Cross-sectional	232	Adults	71.1	Self-rated severity: 33 (14%) very mild, 72 (31%) mild, 103 (44%) moderate, 20 (9%) severe, 4 (<2%) very severe. 91% had been enrolled in speech therapy at one time.	Employment (performance, promotion)	High

Author (Year)	Country	Study design	Size (N)	Cohort	Male (%)	Population characteristics (stuttering diagnosis, severity, treatment)	Cost outcomes	Risk of bias
McAllister et al. (2012)	UK	Cohort	18,558 (no stutter at age 16 and no history of speech problems $n = 15,694$ , stutter at age 16 $n = 217$ )	Stuttered in adolescence, outcomes as adults	PWS: 78.7 to 86.6 <sup>b</sup> PWNS: 47.1 to 67.6 <sup>b</sup>	CWS identified by parent report	Educational (attainment), employment (status and income)	Moderate
McClure and Yaruss (2003)	USA	Cross-sectional	642 (AWS $n = 544$ , parents of CWS $n = 98$ )	Children, adolescents, and adults	Not reported	Prior treatment: of those who had received treatment, 85% had $\geq 2$ experiences. All survey respondents were NSA members, <50% had attended convention, workshop, or meeting	Education (performance), employment (performance, promotion, recruitment)	Critical
O'Brian et al. (2011)	Australia and New Zealand	Cross-sectional	147	Adults	78.9	Stuttering confirmed by SLP. 123 (81.6%) had previously received treatment for stuttering.	Education (achievement)	High
Palasik (2012)	USA	Cross-sectional	184	Adults	72.8	Self-rated stuttering severity: 51% mild, 41% moderate, 8% severe. 21% currently in therapy, 90% have had therapy at any stage	Employment (performance, career development)	High
Plexico et al. (2019)	USA (Geographic locations not equally distributed between PWS and PWNS groups $p = .000$ )	Cross-sectional	164 (PWS $n = 72$ , PWNS $n = 92$ )	Adults	PWS: 63.4 PWNS: 34.1	On a 1-10 (worst) scale, mean (SD) severity: 5.17 (2.37) 86% had some past treatment for stuttering. 40% had received treatment $\geq 4$ times. 74% of those receiving prior treatment experienced relapse. 60% were members of NSA.	Employment (roles, career progression, income, discrimination)	High

Author (Year)	Country	Study design	Size (N)	Cohort	Male (%)	Population characteristics (stuttering diagnosis, severity, treatment)	Cost outcomes	Risk of bias
Rees and Sabia (2014)	USA	Cohort	15,170 (analysed $n = 13,549$ )	Stuttered in adolescence, outcomes as adults	PWS: 57.8 PWNS: 45.5	Self-reported stutterer. 7% answered affirmative in Wave III ( $n \approx 948$ ).	Education (achievement attainment)	Moderate
Rice (1994)	Canada	Cross-sectional	>250	Not reported	Not reported	Not discussed	Employment	Critical
Rice and Kroll (2006)	USA ( $n = 239$ ), Britain ( $n = 64$ ), Canada ( $n = 32$ ), Australia ( $n = 24$ ), India ( $n = 13$ ), other ( $n = 40$ ). 32 countries total	Cross-sectional	412	Adults	71.6	Stuttering severity: mild (39%), moderate (54%) severe (7%).	Employment	High
Rosenbaum (2018)	USA	Case-control	9,909 enrolled in community college $n = 1494$ , enrolled in 4-year college $n = 2,721$ )	Adults	39.6	Self-reported stutterer: 7.2% overall ( $n = 714$ ), 8.1% not attending college, 7.0% attending community college, 5.5% attending 4-yr college	Education (attainment)	Moderate
Sommer et al. (2021)	Germany	Cross-sectional	27,977	Children, adolescents, and adults	75	Confirmed outpatient diagnosis (ICD-10 code recorded in insurer database)	Healthcare utilisation	High
Werle and Byrd (2022)	USA	Cross-sectional	158	College instructors	Not reported	Instructors who teach or evaluate oral presentations in university foundational oral communications courses within last 5 yrs.	Education (performance)	Moderate

Author (Year)	Country	Study design	Size (N)	Cohort	Male (%)	Population characteristics (stuttering diagnosis, severity, treatment)	Cost outcomes	Risk of bias
Williams et al. (1969)	USA	Case-control	1. 400 (CWS <i>n</i> = 100, CWNS <i>n</i> = 300). 2. 200 (male CWS <i>n</i> = 50, male CWNS <i>n</i> = 150)	Children	87	Names of CWS submitted by SLP	Education (performance)	Moderate
<b>Qualitative research</b>								
Beilby et al. (2013)	Unclear	Mixed method (questionnaires, interviews).	20	Adults	PWS: 90 Partner: 10	Clinical diagnosis of stuttering confirmed by a SLP with ≥10 years of experience in assessment and treatment of fluency disorders;	Social (romantic partner)	Lower
Boberg and Boberg (1990)	Canada	Interviews	15	Adults	0	Non-stuttering wives of PWS. 7/15 husbands completed 3wk intensive stuttering program, 3 completed precision fluency program, 3 avoidance reduction, 2 no therapy	Social (family)	Higher
Bricker-Katz et al. (2013)	Australia	Interviews	9	Adults	66.6	Stuttering severity: 2 severe, 3 moderate, 1 mod/mild, 3 mild. Treatment history: 2 current, 3 intermittent/not current, 3 never, 1 early childhood	Employment (occupational progression)	Lower
Butler (2013)	UK	Interviews, focus groups	38	Adults	Just over 80	Typically experiencing dysfluency from age 5yrs	Education (academic achievement, progression)	Lower

Author (Year)	Country	Study design	Size (N)	Cohort	Male (%)	Population characteristics (stuttering diagnosis, severity, treatment)	Cost outcomes	Risk of bias
Butler (2014)	UK	Interviews, focus groups, conversations	36	Adults	100	Diagnosis/ severity not discussed.	Education (achievement), employment (aspirations, recruitment)	Lower
Crichton-Smith (2002)	UK	Interviews	14	Adults	78.6	13 persistent developmental stuttering, 1 acquired stammer. Previous therapy: 2/14 never, 3/14 as child, 4/14 as adult, 5/14 as child + adult.	Education (attainment), employment	Lower
Georges (2017)	USA (n = 9) NZ (n = 1)	Interviews	10	Adults	0	Not described	Employment, social, direct and indirect costs	Lower
Johnson (1934)	USA	Mixed method (interviews, questionnaires, clinical examination, autobiographies).	80	Children, adolescents, and adults (age range 7–42 years)	76.3	Greater proportion of severe cases among boys than girls.	Education (attainment), employment (occupation choice)	Higher
Leko Krhen et al. (2021)	Croatia	Interviews	6	Adults	0	Self-report. PWS >5 yrs. Half perceived stuttering moderate at onset, remainder severe. Stutter onset: half <6rs, remainder 6–10 yrs.	Education, employment, social	Higher
Nang et al. (2018)	Australia	Interviews	9	Adults	0	Stuttering severity: 5 mild, 2 mild-moderate, 2 moderate. All started stuttering in childhood and had received some SLP intervention. Support group membership: 3 < 10yrs, 4 10-20 years, 2 > 30 years.	Social (romantic relationships), employment (recruitment)	Lower

Author (Year)	Country	Study design	Size (N)	Cohort	Male (%)	Population characteristics (stuttering diagnosis, severity, treatment)	Cost outcomes	Risk of bias
Silverman and Zimmer (1982)	US	Interviews	20 (Group 1 n = 10, Group 2 n = 10)	Adults	Group 1: 0 Group 2: 100	Some parent, self, teachers, relative, or friend-diagnosed (no mention of clinical diagnosis). Stutter onset: 4yrs women, 6.2yrs men. Treatment onset: 11.4yrs w, 9.8yr m.	Education, employment, social, healthcare utilisation	Higher

AWS = adults who stutter; BSA = British Stuttering Association; CWNS = children who do not stutter; CWS = children who stutter; DCM = Demands and Capacities Model; LP = Lidcombe Program; NSA = National Stuttering Association; NZ = New Zealand; PWNS = people who do not stutter; PWS = people who stutter; RCT = Randomized Controlled Trial; SD = standard deviation; SLP = speech-language pathologist; SS = sample size; UK = United Kingdom; USA = United States of America; yrs. = years.

<sup>a</sup> This study is described by the authors as a population group cohort study. However, it specifically recruited PWS so has been classified as a cross-sectional study in this review.

<sup>b</sup> Proportion of males was presented by outcome variable (e.g., highest qualification at 50 years, unemployment at 23 years, pay at age 23)