








PERSPECTIVE

Selecting coral species for reef restoration

Joshua S. Madin¹  | Michael McWilliam¹  | Kate Quigley²  | Line K. Bay³ |
David Bellwood⁴  | Christopher Doropoulos⁵  | Leanne Fernandes⁶ | Peter Harrison⁷ |
Andrew S. Hoey⁴  | Peter J. Mumby⁸ | Juan C. Ortiz³ | Zoe T. Richards⁹ |
Cynthia Riginos⁸  | Nina M. D. Schiettekatte¹ | David J. Suggett¹⁰ |
Madeleine J. H. van Oppen^{3,11}

¹Hawai'i Institute of Marine Biology, University of Hawai'i at Mānoa, Kāne'ohe, Hawai'i, USA; ²Minderoo Foundation, Perth, Western Australia, Australia; ³Australian Institute of Marine Science, Townsville, Queensland, Australia; ⁴College of Science and Engineering, James Cook University, Townsville, Queensland, Australia; ⁵CSIRO Environment, Brisbane, Queensland, Australia; ⁶Great Barrier Reef Marine Park Authority, Townsville, Queensland, Australia; ⁷Marine Ecology Research Centre at Southern Cross University, Lismore, New South Wales, Australia; ⁸School of Biological Sciences, The University of Queensland, St. Lucia, Queensland, Australia; ⁹Coral Conservation and Research Group, Trace and Environmental DNA Laboratory, School of Molecular and Life Sciences, Curtin University, Bentley, Western Australia, Australia; ¹⁰University of Technology Sydney, Climate Change Cluster, Sydney, New South Wales, Australia and ¹¹School of BioSciences, The University of Melbourne, Parkville, Victoria, Australia

Correspondence

Joshua S. Madin

Email: jmadin@hawaii.edu

Funding information

Australian Research Council, Grant/Award Number: FL180100036, FL190100062 and LP160101508; Defense Advanced Research Projects Agency, Grant/Award Number: BAA HR001121S0012; National Science Foundation, Grant/Award Number: 1948946; Australian Governments Reef Trust and the Great Barrier Reef Foundation

Handling Editor: Fraser Andrew

Januchowski-Hartley

Abstract

1. Humans have long sought to restore species but little attention has been directed at how to best select a subset of foundation species for maintaining rich assemblages that support ecosystems, like coral reefs and rainforests, which are increasingly threatened by environmental change.
2. We propose a two-part hedging approach that selects optimized sets of species for restoration. The first part acknowledges that biodiversity supports ecosystem functions and services, and so it ensures precaution against loss by allocating an even spread of phenotypic traits. The second part maximizes species and ecosystem persistence by weighting species based on characteristics that are known to improve ecological persistence—for example abundance, species range and tolerance to environmental change.
3. Using existing phenotypic-trait and ecological data for reef building corals, we identified sets of ecologically persistent species by examining marginal returns in occupancy of phenotypic trait space. We compared optimal sets of species with those from the world's southern-most coral reef, which naturally harbours low coral diversity, to show these occupy much of the trait space. Comparison with an existing coral restoration program indicated that current corals used for restoration only cover part of the desired trait space and programs may be improved by including species with different traits.
4. *Synthesis and applications.* While there are many possible criteria for selecting species for restoration, the approach proposed here addresses the need to insure

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2023 The Authors. *Journal of Applied Ecology* published by John Wiley & Sons Ltd on behalf of British Ecological Society.

against unpredictable losses of ecosystem services by focusing on a wide range of phenotypic traits and ecological characteristics. Furthermore, the flexibility of the approach enables the functional goals of restoration to vary depending on environmental context, stakeholder values, and the spatial and temporal scales at which meaningful impacts can be achieved.

KEYWORDS

ecosystem services, hedging, marginal returns, phenotypic traits, reef corals, restoration, species selection

1 | INTRODUCTION

The rate and extent of environmental change experienced by contemporary ecosystems have resulted in major deviations from their historical state (Hobbs et al., 2011). Protection alone may no longer suffice to preserve biodiversity and related ecosystem functions and services, and active restoration is increasingly considered essential. The objective of ecosystem restoration is, through human intervention, to recover a disturbed or degraded ecosystem as far as possible towards some preferred previous state. Interventions can be direct, such as propagation and field deployment of habitat builders through seeds (Orth et al., 2020), propagules (Vanderklift et al., 2020), early recruits (Fredriksen et al., 2020) or parts of adult tissues (Page et al., 2018; Rinkevich, 2014). Indirect interventions, such as physical stabilization of degraded reef structures and removal of macroalgae are also possible (Ceccarelli et al., 2020).

To date, the augmentation or reintroduction of one or few species has been the most common approach, such as the restoration of the endangered Caribbean coral species *Acropora cervicornis* and *A. palmata* (Ladd et al., 2019), the reintroduction of the grey wolf across parts of Europe and North America (Ripple et al., 2014) and assisted colonization of the Tasmanian Devil to the Australian mainland (Brainard, 2020). However, climate change is now affecting many assemblages of foundation species in most if not all the world's ecosystems, including forests, kelp beds and coral reefs. Therefore, broadening the focus of restoration activities to encompass more species and their contributions to ecosystem functioning is required (Coleman & Bragg, 2020; Laughlin, 2014).

2 | SELECTING SPECIES FOR RESTORATION

Prioritizing sets of species for the restoration of biodiverse ecosystems is a challenging task. Some approaches focus on the roles that species play in providing ecosystem goods or services, including carbon storage in rainforests (Strassburg et al., 2020) or reef accretion on coral reefs for coastal protection (Bellwood et al., 2019). Another common focus is on keystone species: species that maintain the organization and stability of their communities and have disproportionately large, positive impacts on their ecosystems (Hale &

Koprowski, 2018). Alternatively, weedy pioneer species may quickly restore habitat functions such as providing shelter or stabilizing substratum; this is exemplified by the emphasis on fast growing acroporids in coral gardening and larval-based restoration initiatives (Boström-Einarsson et al., 2020). However, approaches for species selection that consider multiple ecological, functional and logistical criteria are rare (Lamb, 2018; Suding et al., 2004). Some examples exist for forest restoration (Meli et al., 2013), and some have used linkages between phenotypic traits and ecosystem services to select species (Giannini et al., 2017). Given the rapid growth in ecological and phenotypic trait databases across taxa globally (Gallagher et al., 2020), we propose a hedging approach that maximizes success in the context of a changing environment and uncertain future. When applied to restoration, hedging is the process of selecting species to balance added value to the ecosystem against the risk of future extinction. We contend that hedging approaches should become an integral part of selecting sets of species for restoration.

3 | ANTICIPATING FUTURE ECOSYSTEMS

An overarching challenge is that restoration initiatives need to anticipate future ecosystem states, which are expected to be very different due to the escalating impacts of climate change (Gaitán-Espitia & Hobday, 2021; Rogers et al., 2015). Faced with complex ecosystems, multiple threats to biodiversity and limited funding, conservation practitioners must prioritize investment into different management options, including restoration actions, and difficult decisions are to be made on which sets of species to allocate resources (Game et al., 2018). Strategic decisions must be made about supporting sets of species most likely to do better to improve future persistence and resilience, or helping species that will struggle through a period of elevated and prolonged stress (Coles & Brown, 2003). Protecting habitat-forming species such as corals is imperative for securing the ecosystem services they provide, such as reef building, habitat and food provisioning for commercially important species, coastal protection, and biomedical, social, cultural and recreational opportunities. Given the sheer number of interacting ecological and social considerations, as well as the massive diversity of taxa in ecosystems such as coral reefs, deciding which species to select for restoration is complex, and the answer can change with each new consideration. Hedging strategies may offer a way to

reduce risk of future loss of functions and services, although in some cases these strategies are not much better than selecting species at random (e.g. Nee & May, 1997).

4 | TWO-PART HEDGING APPROACH

The restoration of ecosystems via foundation assemblages should not only consider which species are most likely to persist in the future but also which species provide essential contributions to the current suite of ecosystem services. To balance these considerations, we argue for

a two-part process when selecting sets of foundation species. The first part centers on the maintenance of certain ecosystem services provided by foundation species, ranging from habitat engineering to phylogenetic diversity. However, rather than targeting specific properties that support ecosystem services, we propose to minimize their loss by maximizing phenotypic variation (e.g. selecting species that occupy trait space evenly). To demonstrate, we focus on reef-building corals and used the dataset from McWilliam et al. (2018) containing seven mean phenotypic traits (growth rate, corallite width, rugosity/branch spacing, surface area per unit volume, colony height, maximum colony size/diameter, and skeletal density; Figure 1a) for the




FIGURE 1 Phenotypic trait space as represented by the first two principal component axes for 396 eastern Australia coral species. (a) The 5 × 5 cell grid used to calculate occupancy (see Supporting Information) and the trait loadings: growth rate (GR), corallite width (CW), rugosity/branch spacing (R), surface area per unit volume (SAV), colony height (CH), maximum colony size/diameter (MCS), and skeletal density (SD). (b) A schematic showing that some ecosystem services require species broadly across the space (e.g. reef-building and coastal protection) whereas others only a limited region of the space (e.g. reef stability and habitat provision). The schematic is illustrative only; not based on data. (c) Goreau's (1963) categories of essential reef builders. (d) Darling et al. (2012) life history categories. (e) Veron's (2000) families. Ellipses in (c–e) are 95% confidence for each category's centroid.

396 reef coral species from eastern Australia. These trait data enabled us to capture important dimensions of species life history globally, ranging from fast to slow growth (Darling et al., 2012), fragile to robust morphologies (Zawada et al., 2019) and small to large colonies that drive up colony fecundity (Álvarez-Noriega et al., 2016).

Our reasons for maximizing phenotypic variation are four-fold. First, mechanistic linkages between phenotypic traits of species and the functions and services they provide are poorly understood, especially in marine systems (Bellwood et al., 2019). Therefore, it is difficult to ascertain regions of trait space responsible for particular services. Second, maximizing phenotypic variation increases life history variation, and therefore minimizes the risk of wholesale species loss (i.e. response diversity), because no species is at a selective optimum in all situations and environments (Stearns, 1992). Third, while some services may occupy small regions of trait space, maintaining multiple ecosystem services requires different groups of species covering the entire trait space (Figure 1b). For example, Goreau's (1963) reef building groups—builders, fillers and cementers—span most of coral phenotypic trait space (Figure 1c; see Supporting Information); as do Darling et al.'s (2012) life-history groups (Figure 1d). Finally, hedging strategies also act to increase phylogenetic diversity because many life-history traits are phylogenetically conserved and so occupy limited regions of trait space (Figure 1e) (Westoby et al., 2002). Although mean traits were used here, identifying species with high intraspecific trait variation can enhance the phenotypic variation captured

per species and improve the potential for response diversity, and is recommended where data are available.

The second part of the process is to select species based on ecological characteristics that make them better equipped to avoid depletion and resist or recover from large-scale events, such as marine heatwaves. For example, species with small range sizes and small local populations generally have a higher extinction risk (Stauder et al., 2020), while species with higher local abundances and larger range sizes tend to bounce back faster following disturbance (Halford et al., 2004). Meanwhile, some species are naturally more tolerant to disturbances that are expected to become more frequent and intense, or have greater capacity to adapt or acclimate via intraspecific change. To demonstrate, we used three ecological characteristics from the Coral Trait Database (Madin et al., 2016; ecological abundance, geographic range size, and thermal bleaching susceptibility) to estimate emergent characteristics of populations that are likely to make them more persistent.

5 | EASTERN AUSTRALIA CASE STUDY

While there are many definitions of phenotypic trait diversity (Villéger et al., 2008), our goal under a hedging strategy was to evenly capture the largest area of trait space with the fewest species in a biogeographic region, therefore ensuring a spread of species along important, often orthogonal, trait dimensions. This goal was accomplished by iteratively





FIGURE 2 (a) Proportion occupancy of selected eastern Australia species in trait space as a function of number of species using a 5 × 5 cell grid (see Figure 1a). Regions of high, medium and low marginal returns delineated with dotted vertical lines. Included are symbols that show trait diversity and redundancy for common species at the world's southernmost coral reef, Lord Howe Island (n = 15), and for the Coral Nuture Program (n = 39). The grey shaded region shows 95% CIs for randomized species selection. (b) Lord Howe Island (n = 15) and (c) Coral Nuture Program (n = 39) species are shown in trait space as black points. Red ellipse are Goreau's (1963) reef building categories (centroid 95% confidence regions). Blue crosses are species selected by the two-part hedging process for 15 and 39 species, respectively. Red arrow in (c) is *Galaxea fascicularis*.

removing the species closest to other species in multi-dimensional trait space until some prerequisite number of species remained (see [Supporting Information](#)). Ecological persistence was estimated by multiplying species distances in trait space by standardized ecological characteristics during each iteration (i.e. species with standardized values closer to 1—that is large ranges, ecologically common and resistant to thermal bleaching—were considered ecologically persistent and therefore less likely to be removed during an iteration; [Supporting Information](#)). Occupancy of trait space was measured using a 5 × 5 grid of cells ([Figure 1a](#)). The results are reported in [Figure 2a](#).

The hedging approach highlighted several important points for selecting species ([Figure 2a](#)).

- Selecting species that maximize distances in trait space resulted in superior levels of occupancy ([Figure 2a](#), yellow curve). However, selecting species randomly yields relatively high levels of occupancy (see also Nee & May, 1997), and so might be used when phenotypic trait data are missing or incomplete ([Figure 2a](#), black curve and grey 95% confidence band).
- Selecting species based on ecological persistence alone (i.e. hedging against future loss of restored species) was broadly equivalent to random selection with regard to trait space occupancy ([Figure 2a](#), green curve). This result suggests that species likely to weather future conditions are distributed randomly across the trait space, which is a positive outcome.



- Integrating ecological persistence with trait diversity slightly reduces trait space occupancy compared with using traits diversity only, because species occupying distinct regions of trait space are being removed because of their lower potential to survive future conditions. That is, this two-part hedging approach protects against selecting high-risk species in the effort to hedge against losses of ecosystem service.
- In terms of how many species to select, as species are added, marginal returns in trait space occupancy are high (approximately 5% occupancy per species) up until approximately 11 species; medium (approximately 2% per species) between from 11 to 28 species; and low (<0.3% per species) above $n=28$ (Figure 2a, vertical dotted lines). The general patterns shown in Figure 2a were robust to the cell size of the grid used to calculate occupancy (Figure S2); with the proviso that more species are required to maintain specified levels of occupancy for finer grids. Our hedging approach therefore clearly quantifies “bang for buck”: that is, the amount of future hedging per species to be restored.
- We compared our hedging results with those of the southernmost accreting reef in eastern Australia, Lord Howe Island, and a coral restoration program, the Coral Nurture Program. The 15 common coral species found at Lord Howe Island occupy trait space no differently from randomly selecting species (Figure 2a, asterisk) and occupy broad regions that likely support reef building (Figure 2b, red ellipses). Coral Nurture Program, which selected 39 species primarily based on commonness and ease of out-planting (i.e. mostly branching species), occupies only 40%–50% of the possible trait space (Figure 2a, diamond) and does not occupy regions of trait space with reef “builders” (with the exception of one species, *Galaxea fascicularis*, indicated by the red arrow, Figure 2c). We identify in both examples the species that would have been selected from the total community of corals found based on the two-part hedging process (Figure 2b,c, blue crosses).
- Finally, the two-part approach can be flexible and expanded upon. By way of example, we use the hedging approach to select 20 eastern Australia species (a number selected to represent medium marginal returns in Figure 2a) based on two management scenarios. First, we select species most likely to persist in the future based on high scores for ecological abundance, geographic range size and bleaching resistance (Figure 3a). Second, we select species most vulnerable to thermal bleaching, but with high scores for ecological abundance and geographic range size (Figure 3b). The first columns in each panel are sets of species based on ecological characteristics alone. The second columns are sets of species when additionally considering trait space occupancy. The third columns are sets of species when expanding the approach to also consider a species' ease of restoration (here we rank morphologies by ease of restoration as per Boström-Einarsson et al., 2020; see Supporting Information). Figure 3 shows the impact of switching to species that are less difficult to restore based on our criteria, but not as

ecologically valuable, while retaining an even spread of species in the trait space.

6 | CONCLUSIONS

The two-part hedging approach outlined here, and demonstrated with reef-building corals, identified species for active restoration projects based on both diversity of life-history traits and ecologically beneficial characteristics. Selection based on ecological characteristics are important for hedging against future species loss, whereas trait diversity is important for hedging against the loss of certain ecosystem services, reef-building groups, life-history categories, and phylogenetic diversity. The importance of individual species for ecosystem processes, functions and services are poorly understood, particularly for reef-building corals. Furthermore, selecting species based on ecosystem services only is likely to vary across systems depending on the specific environment, and the values of the local stakeholders (Bellwood et al., 2019). Our tactic was therefore to prioritize an even spread of species across the trait space rather than prioritizing particular phenotypic trait values or targeting regions in the trait space (Figure 1). When selecting species for restoration, a species-oriented focus may favour rare or depleted species with low persistence. However, ecosystem restoration based on trait diversity is more likely to be robust if it favours foundation species that are dominant and persistent (as we advocate here), and therefore more likely to maintain a broad range of ecosystem services. The number of species that can be selected for a restoration project will ultimately depend upon project goals, as well as resource and logistical constraints; however, marginal returns in trait-space occupancy can help managers and practitioners decide where to draw the best line. The flexible species selection process developed here can serve as a framework for such decisions, and can serve an important role even as the goals of restoration are refined based on an improved knowledge of ecosystem services, successful restoration methods, diverse stakeholder values, and the scales at which restoration is most effective.

AUTHOR CONTRIBUTIONS

Joshua S. Madin, Michael McWilliam, Kate Quigley, Line K. Bay, David Bellwood, Christopher Doropoulos, Leanne Fernandes, Peter Harrison, Andrew S. Hoey, Peter J. Mumby, Juan C. Ortiz, Zoe T. Richards, Cynthia Riginos, David J. Suggett and Madeleine J. H. van Oppen conceived the idea during a working group meeting organized by Madeleine J. H. van Oppen and Kate Quigley. Joshua S. Madin and Michael McWilliam developed the idea, gathered data and ran analyses. Joshua Madin, Michael McWilliam, Kate Quigley and Madeleine J. H. van Oppen wrote the first draft. Joshua S. Madin, Michael McWilliam, Kate Quigley, Line K. Bay, David Bellwood, Christopher Doropoulos, Leanne Fernandes, Peter Harrison, Andrew S. Hoey, Peter J. Mumby, Juan C. Ortiz, Zoe T. Richards, Cynthia Riginos, Nina M. D. Schiettekatte, David J. Suggett and Madeleine J. H. van Oppen critically revised drafts and added intellectual content.

ACKNOWLEDGEMENTS

The workshop was funded by the Australian Research Council Laureate Fellowship FL180100036 to Madeleine J. H. van Oppen. Madeleine J. H. van Oppen, Christopher Doropoulos and Line K. Bay acknowledge the Reef Restoration and Adaptation program, which is funded by the partnership between the Australian Governments Reef Trust and the Great Barrier Reef Foundation. We also acknowledge the National Science Foundation (1948946 to Joshua S. Madin), Australian Research Council (LP160101508 to Zoe T. Richards and FL190100062 to David Bellwood) and Defence Advanced Research Projects Agency under the Reefense Program (BAA HR001121S0012 to Joshua S. Madin).

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

DATA AVAILABILITY STATEMENT

No new data were collected for this study. The data sets used are cited and included with the analytical code at Zenodo (<https://www.doi.org/10.5281/zenodo.7949521>, Madin et al., 2016; McWilliam et al., 2018; Swain et al., 2016) and GitHub (<https://github.com/jmadinlab/species-choice>).

ORCID

Joshua S. Madin  <https://orcid.org/0000-0002-5005-6227>
 Michael McWilliam  <https://orcid.org/0000-0001-5748-0859>
 Kate Quigley  <https://orcid.org/0000-0001-5558-1904>
 David Bellwood  <https://orcid.org/0000-0001-8911-1804>
 Christopher Doropoulos  <https://orcid.org/0000-0001-8038-2771>
 Andrew S. Hoey  <https://orcid.org/0000-0002-4261-5594>
 Cynthia Riginos  <https://orcid.org/0000-0002-5485-4197>

REFERENCES

- Álvarez-Noriega, M., Baird, A., Dornelas, M., Madin, J., Cumbo, V., & Connolly, S. (2016). Fecundity and the demographic strategies of coral morphologies. *Ecology*, 97, 3485–3493.
- Bellwood, D. R., Pratchett, M. S., Morrison, T. H., Gurney, G. G., Hughes, T. P., Álvarez-Romero, J. G., Day, J. C., Grantham, R., Grech, A., Hoey, A. S., Jones, G. P., Pandolfi, J. M., Tebbett, S. B., Techera, E., Weeks, R., & Cumming, G. S. (2019). Coral reef conservation in the Anthropocene: Confronting spatial mismatches and prioritizing functions. *Biological Conservation*, 236, 604–615.
- Boström-Einarsson, L., Babcock, R., Bayraktarov, E., Ceccarelli, D., Cook, N., Ferse, S., Hancock, B., Harrison, P., Hein, M., Shaver, E., Smith, A., Suggett, D., Stewart-Sinclair, P., Vardi, T., & McLeod, I. (2020). Coral restoration—A systematic review of current methods, successes, failures and future directions. *PLoS One*, 15, e0226631.
- Brainard, J. (2020). Tasmanian devil reintroduced. *Science*, 370, 268.
- Ceccarelli, D. M., McLeod, I. M., Boström-Einarsson, L., Bryan, S. E., Chartrand, K. M., Emslie, M. J., Gibbs, M. T., Gonzalez Rivero, M., Hein, M. Y., Heyward, A., Kenyon, T. M., Lewis, B. M., Mattocks, N., Newlands, M., Schlappi, M.-L., Suggett, D. J., & Bay, L. K. (2020). Substrate stabilisation and small structures in coral restoration: State of knowledge, and considerations for management and implementation. *PLoS One*, 15, e0240846.
- Coleman, M. A., & Bragg, J. G. (2020). A decision framework for evidence-based climate adaptation interventions. *Global Change Biology*, 27, 472–474.
- Coles, S., & Brown, B. (2003). Coral bleaching-capacity for acclimatization and adaptation. *Advances in Marine Biology*, 46, 183–223.
- Darling, E., Alvarez-Filip, L., Oliver, T., McClanahan, T., & Côté, I. (2012). Evaluating life-history strategies of reef corals from species traits. *Ecology Letters*, 15, 1378–1386.
- Fredriksen, S., Filbee-Dexter, K., Norderhaug, K. M., Steen, H., Bodvin, T., Coleman, M. A., Moy, F., & Wernberg, T. (2020). Green gravel: A novel restoration tool to combat kelp forest decline. *Scientific Reports*, 10, 3983.
- Gaitán-Espitia, J., & Hobday, A. (2021). Evolutionary principles and genetic considerations for guiding conservation interventions under climate change. *Global Change Biology*, 27, 475–488.
- Gallagher, R., Falster, D., Maitner, B., Salguero-Gómez, R., Vandvik, V., Pearse, W., Schneider, F., Kattge, J., Poelen, J., Madin, J., Ankenbrand, M., Penone, C., Feng, X., Adams, V., Alroy, J., Andrew, S., Balk, M., Bland, L., Boyle, B., ... Enquist, B. J. (2020). Open Science principles for accelerating trait-based science across the tree of life. *Nature Ecology & Evolution*, 4, 294–303.
- Game, E., Tallis, H., Olander, L., Alexander, S., Busch, J., Cartwright, N., Kalies, E., Masuda, Y., Mupepele, A.-C., Qiu, J., Rooney, A., Sills, E., & Sutherland, W. (2018). Cross-discipline evidence principles for sustainability policy. *Nat Sustain*, 1, 452–454.
- Giannini, T., Giuletta, A., Harley, R., Viana, P., Jaffe, R., Alves, R., Pinto, C., Mota, N., Caldeira, C., Imperatriz-Fonseca, V., Furtini, A., & Siqueira, J. (2017). Selecting plant species for practical restoration of degraded lands using a multiple-trait approach. *Austral Ecology*, 42, 510–521.
- Goreau, T. (1963). Calcium carbonate deposition by coralline algae and corals in relation to their roles as reef-builders. *Comparative Biology of Calcified Tissue*, 109, 127–167.
- Hale, S., & Koprowski, J. (2018). Ecosystem-level effects of keystone species reintroduction: A literature review: Effects of keystone species reintroduction. *Restoration Ecology*, 26, 439–445.
- Halford, A., Cheal, A., Ryan, D., & Williams, D. (2004). Resilience to large-scale disturbance in coral and fish assemblages on the Great Barrier Reef. *Ecology*, 85, 1892–1905.
- Hobbs, R. J., Hallett, L. M., Ehrlich, P. R., & Mooney, H. A. (2011). Intervention ecology: Applying ecological science in the twenty-first century. *Bioscience*, 61, 442–450.
- Ladd, M. C., Burkepile, D. E., & Shantz, A. A. (2019). Near-term impacts of coral restoration on target species, coral reef community structure, and ecological processes. *Restoration Ecology*, 27, 1166–1176.
- Lamb, D. (2018). Undertaking large-scale forest restoration to generate ecosystem services: Landscape restoration and ecosystem services. *Restoration Ecology*, 26, 657–666.
- Laughlin, D. (2014). Applying trait-based models to achieve functional targets for theory-driven ecological restoration. *Ecology Letters*, 17(7), 771–784.
- Madin, J., Anderson, K., Andreasen, M., Bridge, T., Cairns, S., Connolly, S., Darling, E., Diaz, M., Falster, D., Franklin, E., Gates, R., Hoogenboom, M., Huang, D., Keith, S., Kosnik, M., Kuo, C.-Y., Lough, J., Lovelock, C., Luiz, O., ... Baird, A. (2016). The coral trait database, a curated database of trait information for coral species from the global oceans. *Scientific Data*, 3, 160017.
- McWilliam, M., Hoogenboom, M., Baird, A., Kuo, C.-Y., Madin, J., & Hughes, T. (2018). Biogeographical disparity in the functional diversity and redundancy of corals. *Proceedings of the National Academy of Sciences of the United States of America*, 115, 3084–3089.
- Meli, P., Martínez-Ramos, M., & Rey-Benayas, J. (2013). Selecting species for passive and active riparian restoration in southern Mexico: Selecting species for riparian restoration. *Restoration Ecology*, 21, 163–165.

- Nee, S., & May, R. (1997). Extinction and the loss of evolutionary history. *Science*, 278, 692–694.
- Orth, R. J., Lefcheck, J. S., McGlathery, K. S., Aoki, L., Luckenbach, M. W., Moore, K. A., Oreska, M. P. J., Snyder, R., Wilcox, D. J., & Lusk, B. (2020). Restoration of seagrass habitat leads to rapid recovery of coastal ecosystem services. *Science Advances*, 6, eabc6434.
- Page, C., Muller, E., & Vaughan, D. (2018). Microfragmenting for the successful restoration of slow growing massive corals. *Ecological Engineering*, 123, 86–94.
- Rinkevich, B. (2014). Rebuilding coral reefs: Does active reef restoration lead to sustainable reefs? *Current Opinion in Environmental Sustainability*, 7, 28–36.
- Ripple, W., Estes, J., Beschta, R., Wilmers, C., Ritchie, E., Hebblewhite, M., Berger, J., Elmhagen, B., Letnic, M., Nelson, M., Schmitz, O., Smith, D., Wallach, A., & Wirsing, A. (2014). Status and ecological effects of the World's largest carnivores. *Science*, 343, 1241484.
- Rogers, A., Harborne, A., Brown, C., Bozec, Y.-M., Castro, C., Chollett, I., Hock, K., Knowland, C., Marshall, A., Ortiz, J., Razak, T., Roff, G., Samper-Villarreal, J., Saunders, M., Wolff, N., & Mumby, P. (2015). Anticipative management for coral reef ecosystem services in the 21st century. *Global Change Biology*, 21, 504–514.
- Staude, I., Navarro, L., & Pereira, H. (2020). Range size predicts the risk of local extinction from habitat loss. *Global Ecology and Biogeography*, 29, 16–25.
- Stearns, S. (1992). *The evolution of life histories*. Oxford University Press.
- Strassburg, B., Iribarrem, A., Beyer, H., Cordeiro, C., Crouzeilles, R., Jakovac, C., Braga, J. A., Lacerda, E., Latawiec, A., Balmford, A., Brooks, T., Butchart, S., Chazdon, R., Erb, K.-H., Brancalion, P., Buchanan, G., Cooper, D., Díaz, S., Donald, P., ... Visconti, P. (2020). Global priority areas for ecosystem restoration. *Nature*, 586, 724–729.
- Suding, K., Gross, K., & Houseman, G. (2004). Alternative states and positive feedbacks in restoration ecology. *Trends in Ecology & Evolution*, 19, 46–53.
- Swain, T. D., Vega-Perkins, J. B., Oestreich, W. K., Triebold, C., DuBois, E., Henss, J., Baird, A., Siple, M., Backman, V., & Marcelino, L. (2016). Coral bleaching response index: A new tool to standardize and compare susceptibility to thermal bleaching. *Global Change Biology*, 22, 2475–2488.
- Vanderklift, M. A., Doropoulos, C., Gorman, D., Leal, I., Minne, A. J. P., Statton, J., Steven, A. D. L., & Wernberg, T. (2020). Using propagules to restore coastal marine ecosystems. *Frontiers in Marine Science*, 7, 724.
- Veron, J. (2000). *Corals of the world*. Australian Institute of Marine Science and CCR Qld Pty Ltd.
- Villéger, S., Mason, N., & Mouillot, D. (2008). New multidimensional functional diversity indices for a multifaceted framework in functional ecology. *Ecology*, 89, 2290–2301.
- Westoby, M., Falster, D., Moles, A., Vesk, P., & Wright, I. (2002). Plant ecological strategies: Some leading dimensions of variation between species. *Annual Review of Ecology and Systematics*, 33, 125–159.
- Zawada, K., Dornelas, D., & Madin, J. (2019). Quantifying coral morphology. *Coral Reefs*, 38, 1281–1292.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

Appendix S1. Supplementary methods.

How to cite this article: Madin, J. S., McWilliam, M., Quigley, K., Bay, L. K., Bellwood, D., Doropoulos, C., Fernandes, L., Harrison, P., Hoey, A. S., Mumby, P. J., Ortiz, J. C., Richards, Z. T., Riginos, C., Schiethkatte, N. M. D., Suggett, D. J., & van Oppen, M. J. H. (2023). Selecting coral species for reef restoration. *Journal of Applied Ecology*, 60, 1537–1544. <https://doi.org/10.1111/1365-2664.14447>