

# Mental health consequences of COVID-19 suppression strategies in Victoria, Australia: a narrative review

Journal of International Medical Research

2022, Vol. 50(11) 1–17

© The Author(s) 2022

Article reuse guidelines:

[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)

DOI: 10.1177/03000605221134466

[journals.sagepub.com/home/imr](https://journals.sagepub.com/home/imr)

Jacqueline Jiang<sup>1</sup> , Hamed Akhlaghi<sup>2,3,4</sup>,  
Darren Haywood<sup>5</sup> , Brendan Morrissey<sup>2</sup> and  
Stephen Parnis<sup>2,6</sup>

## Abstract

The COVID-19 pandemic has imposed significant mental health burdens upon the general population worldwide, either directly owing to the disease or indirectly through aggressive public health measures to control spread of the virus that causes COVID-19. In this narrative review, we used a systematic approach to summarize the impact of restrictive lockdown measures on the general mental health of people living in Victoria, Australia during 2020 and to identify the groups with an increased risk of adverse mental health outcomes. A systematic database search (Ovid Medline, PsycINFO, Embase) for articles examining the mental health of Victorians in the context of the COVID-19 pandemic during 2020 yielded 88 articles, of which 15 articles were finally included in this review. We found that the general mental health of Victorians was negatively affected by COVID-19 restrictions during 2020. Although studies reported heterogeneous mental health outcomes, we found that the general population consistently used coping strategies and demonstrated mental health help-seeking behaviors in response to the restrictions. Women, children, young people, carers, people who became unemployed owing to the pandemic, and those with pre-existing psychiatric conditions had a higher risk of adverse mental health consequences during the COVID-19 pandemic in 2020.

<sup>1</sup>Melbourne Medical School, University of Melbourne, Melbourne, VIC, Australia

<sup>2</sup>Department of Emergency Medicine, St Vincent's Hospital Melbourne, Melbourne, VIC, Australia

<sup>3</sup>Faculty of Health, Deakin University, Melbourne, VIC, Australia

<sup>4</sup>Faculty of Medical Education, University of Melbourne, Melbourne, VIC, Australia

<sup>5</sup>St Vincent's Mental Health, St Vincent's Hospital Melbourne, Melbourne, VIC, Australia

<sup>6</sup>Royal Victorian Eye and Ear Hospital, Melbourne, VIC, Australia

## Corresponding author:

Jacqueline Jiang, St Vincent's Clinical School, 23 Regent St, Fitzroy VIC 3065, Australia.

Email: [jacqueline.jiang@svha.org.au](mailto:jacqueline.jiang@svha.org.au); [jacqueline.j.jiang@gmail.com](mailto:jacqueline.j.jiang@gmail.com)



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative

Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

**Keywords**

Mental health, COVID-19, lockdown, Victoria, Australia, public health, social measures, epidemic control

Date received: 3 June 2022; accepted: 5 October 2022

**Introduction**

The novel coronavirus SARS-CoV-2, initially detected in Wuhan, China in December 2019, has spread to nearly 200 countries worldwide.<sup>1</sup> By the end of 2020, the COVID-19 pandemic had led to more than 79 million reported cases and caused over 1.7 million deaths.<sup>2</sup> In an effort to limit the spread of the virus and support health services at risk of being overwhelmed, governments around the world implemented restrictions to limit people's movements, such as physical distancing, restrictions on social gatherings, industry and education closures, stay-at-home orders, and lockdowns. These COVID-19 pandemic suppression strategies were effective at slowing the spread of the virus<sup>3</sup> but were not without profound social, economic, and psychological consequences.<sup>4</sup>

In March 2020, the World Health Organization recognized the global increase in fear, stress, and anxiety in response to the pandemic.<sup>4</sup> An early study in Italy, the first European country to be seriously affected by SARS-CoV-2, showed high rates of post-traumatic stress disorder, depression, anxiety, insomnia, perceived stress, and adjustment disorder symptoms.<sup>5</sup> These mental health presentations were either directly associated with SARS-CoV-2, such as by infection with the virus or death of a loved one, or indirectly related to lockdown measures owing to the pandemic, such as having a lack of certainty about the future, reduced income or unemployment, and social isolation.<sup>5</sup> In Switzerland, higher

levels of stress, anxiety, loneliness, and depression were reported owing to COVID-19 pandemic restrictions, tightened physical and social isolation, and decreased emotional support.<sup>6</sup> Worldwide, the mental health impacts have been profound. However, further research is required to distinguish between direct and indirect consequences of the COVID-19 pandemic.

In the state of Victoria, Australia, the prevalence and transmission of SARS-CoV-2 remained relatively low throughout 2020. Proportional to the number of SARS-CoV-2 cases, the Victorian government imposed some of the strictest restrictions in the world.<sup>7,8</sup> As of 30 November 2020, Victoria had recorded 20,345 confirmed cases of SARS-CoV-2 since the beginning of the pandemic and 820 deaths.<sup>9</sup> The first epidemic wave in Australia began in March 2020, with nationwide implementation of COVID-19 pandemic suppression strategies, including national border closures, 1.5-meter social distancing, stay-at-home orders, work-from-home directives, industry and school closures, and bans on public and household gatherings.<sup>10</sup> After a brief period of staggered lifting of restrictions in late May 2020, a second wave of SARS-CoV-2 emerged in Victoria, prompting the state government to instate the most stringent measures in Australia. New stricter rules were introduced in August 2020, including a maximum 1 hour of outdoor exercise daily, travel restrictions to within 5 km of one's home, and an 8 pm to 5 am curfew, in addition to the

restrictions implemented during the first lockdown.<sup>11</sup> These rules were strictly enforced by Victorian police, who were authorized by the Victoria Department of Health under the Pandemic Declaration to issue on-the-spot fines of AUD 1652 to individuals for infringement of the emergency measures.<sup>12</sup> By 13 May 2020, the Victorian police had completed 43,829 COVID-19 police checks and issued 2894 fines and 833 warnings.<sup>12</sup> These rules were enforced until a gradual staged reopening began in September 2020, with the state entering COVID-Safe Summer rules on 7 December 2020 and most restrictions lifted except for limits on density and gatherings.<sup>11</sup> In 2020, Melbournians in the capital of Victoria spent 154 days in lockdown, 43 days during the first lockdown from 30 March to 12 May (the “first wave”), followed by 111 days during the second lockdown from 8 July to October 27 (“second wave”).<sup>13</sup> This was one of the longest lockdowns in any region during 2020.<sup>14</sup>

Emerging evidence from other countries on the adverse mental health consequences of lockdowns indicates that it is important to specifically investigate these effects in Victoria, Australia, where restrictions were relatively long and onerous.<sup>15</sup> In this narrative review, we aimed to summarize the impact of COVID-19 pandemic restrictions on the mental health of Victoria residents in 2020 and identify the groups with an increased risk of adverse mental health outcomes during the pandemic. The information in this review can assist public health officials and policymakers in decision-making when implementing further COVID-19 pandemic suppression strategies. This review can also provide valuable insight for primary care physicians, psychologists, and other relevant mental health professionals and organizations regarding the impact of COVID-19 pandemic restrictions on the general population of Victoria. This narrative review will also

assist in identifying which groups are most vulnerable and at higher risk of developing adverse mental health consequences during a pandemic.

## Methods

### *Literature search strategy*

On 16 August 2021, we conducted a search of OVID Medline, PsycINFO, and Embase for publications related to the mental health of residents in Victoria, Australia associated with the COVID-19 pandemic.

The first Medline search included the Medical Subject Headings (MeSH) terms “COVID-19” and “pandemics.” These were combined using the Boolean operator OR in a title and abstract search using the terms “coronavirus,” “SARS-CoV-2,” “COVID-19,” and “pandemic.” The second Medline search included “Victoria\*” or “Melbourne” in a title and abstract search. Next, we used the MeSH terms (“Mental Health,” exp “Stress, Psychological,” exp “Suicide,” exp “Anxiety,” “Depression”), combined with title and abstract searches of the terms “mental\*,” “psych\*,” “mood\*,” “psychosocial,” “psycho-social,” “stress\*,” “distress\*,” “burnout,” “burnt-out,” “burnt out,” “burn out,” “burn-out,” “anxiet\*,” “worry,” “worri\*,” “fear\*,” “depress\*,” and “lonel\*.” Subject headings were explored when sub-headings were deemed relevant to this review. The three searches were combined using the AND Boolean operator. Results were limited to studies published from “2019-current” to exclude studies prior to the COVID-19 pandemic. The same search was performed in Embase. In PsycINFO, Thesaurus of Psychological Index Terms (exp “Mental Health,” exp “Psychological Stress,” exp “Stress,” exp “Social Stress,” exp “Stress Reactions,” exp “Stress and Coping Measures,” exp “Depression (emotion),”

exp “Anxiety,” “Fear,” and “Stress”) were used rather than MeSH terms.

### Inclusion and exclusion criteria

The inclusion criteria were all studies that investigated the mental health of Victorians (Australia) during the COVID-19 pandemic.

The exclusion criteria were studies about mental health associated with direct COVID-19 infection, study protocols, clinical correspondence, journal letters, and articles without an available full text. We also excluded studies of psychotic disorders, electroconvulsive therapy, in-patient admissions, and psychiatry treatment practices (e.g., telehealth patterns). However, we included studies reporting emergency department psychiatric presentations.

An ethics review and informed consent were not necessary as the study was a narrative review of existing literature.

## Results

The initial search identified 133 articles, 58 from Medline (OVID–All), 6 from

PsycINFO, and 69 from Embase. These were imported into Covidence software.<sup>16</sup> Duplicates were automatically removed and the retrieved articles were then checked manually, leaving 88 articles. The title/abstract and then full text of these articles were screened by one reviewer (JJ), according to the inclusion and exclusion criteria. An independent reviewer (HA) subsequently verified the suitability of the included and excluded studies, according to the inclusion and exclusion criteria. Any disagreements were solved through discussion. The results of study selection are shown in Figure 1.

Ultimately, 15 articles were included in this review, including 7 cross-sectional surveys, 4 retrospective data analyses, 2 qualitative studies, 1 cohort study, and 1 mixed-methods study. Table 1 provides the details of each study.

The 15 articles included different study populations. Rahman, et al.,<sup>17</sup> Staples et al.,<sup>18</sup> Czeisler et al.,<sup>15</sup> and Fisher et al.<sup>19</sup> reported on the general population. The remaining studies focused on early childhood educators, people living with HIV, pet owners, adults with type 2 diabetes,




Figure 1. PRISMA diagram.

**Table 1.** Summary of selected studies.

Author(s) or Organization/ Publication date	Study design	Sample size	Target population	Study period	Pandemic context	Key results	Limitations
Rahman et al./ 8 October 2020	Cross-sectional survey	587	Patients, frontline health/other essential service workers, Australian community members (88.2% Victorian), aged 18+ years	1–30 June 2020	Early 2nd wave	Higher psychological distress reported by individuals with pre-existing mental health conditions, increased smoking/alcohol use behaviors, and female individuals. Higher fear among individuals who were middle aged, living with family members, born overseas, unemployed, used alcohol.	No comparable pre-pandemic data; selection bias likely toward distressed individuals.
Eadie et al./ 10 May 2021	Cross-sectional survey	232	Early childhood educators (70% Victorian)	July 2020	Early 2nd wave	In total, 85.9% reported the pandemic had a strong negative or slightly negative impact on their well-being.	No comparable pre-pandemic data; selection bias likely for educators with poorer well-being.
Weerasuria et al./ 17 March 2021	Cross-sectional survey	153	People living with HIV	26 August 2020 to 26 November 2020	2nd wave	Doubled proportion of people living with HIV reporting negative impact on relationships since the NAPWHA survey earlier in the pandemic. In total, 66% of participants reported worrying about mental health.	Small sample size; no comparable data to general population or other states; selection bias as participants recruited through infectious disease clinics were likely to be more engaged in health care.
Phillipou et al./ 8 June 2021	Retrospective data analysis—data from cross-sectional survey	263	Pet owners (138) vs. non-pet owners (125)	1–4 September 2020	Mid-2nd wave	Owning a dog or cat associated with reduced quality of life, but not resilience or loneliness.	Inadequate discussion of likely confounding factors; no comparable pre-pandemic data to infer impact of COVID-19 restrictions.

(continued)

**Table 1.** Continued.

Author(s) or Organization/ Publication date	Study design	Sample size	Target population	Study period	Pandemic context	Key results	Limitations
Staples et al./27 February 2021	Retrospective data analysis	14,703	Victorians vs. other Australians	March to October 2020	1st and 2nd waves	Increase in digital mental health clinic service demand in first and second waves among Victorians. No increase in mean standardized measures of psychological distress (K-10), depression (PHQ-9), and anxiety (GAD-7).	Results not generalizable to Victorians owing to selection bias toward participants who actively sought help from the digital mental health service.
Sacre et al./ 26 May 2021	Cohort study	489	Adults 18–80 years with type 2 diabetes, living within 10-km radius of Baker Institute (Melbourne)	30 April 2020 to June/July 2020	Pre-pandemic before 2020, early 2nd wave	No increased anxiety (GAD-7) or depressive (PHQ-9) symptoms compared with pre-COVID-19 levels. No associations between worry levels and age, sex. Leisure activities, feelings about the future, and emotional well-being most common domains impacted by the pandemic. Limited impact of pandemic on within-household relationships and sleep. Greater negative impact on quality of life and leisure activities in younger participants.	Participants not representative of general population—predominantly male (65%), university-educated (55%), older adults (mean age 66 years)—owing to selection of participants within 10 km of Baker Institute.
Atmuri et al./ 10 March 2021	Qualitative descriptive—semi-structured interviews	15	Pregnant patients from secondary outer metropolitan Melbourne hospital	1–19 June 2020	Early 2nd wave	Decreased support owing to physical-distancing restrictions set by the community and hospital. Change in pregnancy service delivery (e.g., to telehealth) caused stress, concern, and uncertainty. Women with children needed additional support.	Small sample size selected by convenience sampling limits generalizability of findings and ability to identify risk factors and confounding factors for mental health outcomes.

(continued)

**Table 1.** Continued.

Author(s) or Organization/ Publication date	Study design	Sample size	Target population	Study period	Pandemic context	Key results	Limitations
Dwyer et al./ 12 July 2021	Mixed-methods— quantitative (interrupted time-series analy- sis) and qualita- tive (content analysis)	N/A	Suicide cases in Victoria	1 January 2015– 31 January 2021	1st and 2nd waves	Suicide frequency did not increase during the pan- demic. No significant change in the demographics of suicide cases compared with pre- pandemic characteristics. COVID-19 pandemic unlikely a discrete stressor but increased stress through negative life events (e.g., job loss, isolation, rela- tionship strain).	Police summaries used to analyze suicide context varied in detail and were not standardized, limiting qualitative analysis. Suicides where COVID-19 pandemic could have been a background or indirect stressor were not deemed to be pandemic-related— possible underestimation of COVID-19-linked suicides.
Digby et al./ 29 September 2020	Qualitative (survey with five free-text questions)	321	Medical, nursing, allied health, non-clinical staff at met- ropolitan Melbourne hospital	16 April–13 May 2020	Easing of 1st wave, prior to 2nd wave	Hospital staff commonly reported subjective anx- iety and concern for the future, emotional isola- tion, and worry about pandemic management. Respondents shared con- cerns about job security, global economic impacts, and loss of control over work and personal situations.	Mental health parameters not formally measured. Themes identified applicable to health care workers but not all applicable to general population of Victoria.
Holton et al./ 9 October 2020	Cross-sectional survey	668	Nurses, midwives, doctors, allied health staff at met- ropolitan Melbourne hospital	15 May–10 June 2020	Between 1st and 2nd waves	Psychological distress symptoms reported by approximately one-quar- ter of respondents. Between 11% (AH staff) and 29% (nurses/midwives) had anxiety scores in the range of mild to extremely severe. Nurses and mid- wives had significantly higher anxiety scores than doctors and AH staff.	No comparison to pre- pandemic data. Study focused on hospital's response to the pandemic causing mental health impacts on staff. No broader discussion about impacts from COVID-19 suppression strategies or confounding factors.

(continued)

**Table 1.** Continued.

Author(s) or Organization/ Publication date	Study design	Sample size	Target population	Study period	Pandemic context	Key results	Limitations
Czeisler et al./ 29 May 2021	Two cross-sectional surveys (cross-sectional & longitudinal analysis)	300 (1st survey) 1157 (2nd survey)	1st survey—Australians aged 18+ years 2nd survey—Victorians aged 18+	2–8 April 2020 15–24 September 2020	1st and 2nd waves	Depression, anxiety and stress levels higher in staff with less clinical expertise, poorer general health, and more COVID-related concerns No significant difference in anxiety, depressive disorder, burnout or sleep measures between April and September 2020. Significant difference in behavioral outcomes (e.g., 1 h increased screen time) between April and September 2020. Adverse mental health and behavioral conditions more common in younger adults, people with disabilities, caregivers, and those with a previous psychiatric diagnosis.	Longitudinal analyses used data from participants recruited in first survey and re-contacted for the second survey, thus limited by selection bias twice, possibly selecting for people who are more proactive about their mental health.
Siette et al./ 30 June 2021	Cross-sectional survey	2990 (253 Victorians)	Australian residents aged 55+ years, no self-reported diagnosis of dementia	10 July–28 September 2020	2nd wave	In total, 47.8% of Victorians versus 40.5% of other Australians reported a negative impact of COVID-19 on their mental health ( $p < 0.0001$ ). No significant difference in prevalence of depression or anxiety (health status) between Victorians and other Australians. Mean quality of life scores were higher in Victorians (0.84) than in other Australians (0.80).	Mental health parameters not measured in detail or with standardized validated criteria. No comparable demographic mental health characteristics. No confounding factors specifically identified or addressed.

(continued)



**Table 1. Continued.**

Author(s) or Organization/ Publication date	Study design	Sample size	Target population	Study period	Pandemic context	Key results	Limitations
Fisher et al./ 27 June 2021	Two cross-sectional surveys	23,749 1st survey: 13,829 (6105 Victorians) 2nd survey: 9220 (4844 Victorians)	Australian residents aged 18+ years	3 April–2 May 2020 1 July–31 August 2020	1st and 2nd waves	Victoria's stage 4 restrictions associated with increased odds of clinically significant depression and anxiety symptoms, becoming easily annoyed or irritable, and lower optimism about the future.  No significant effect on suicidal ideation or self-harm. In both surveys, poor mental health outcomes were more common in people who had low socioeconomic status, lived alone or in poorly resourced areas, lost their job, students with courses suspended, members of marginalized minority groups, carers, women, and young people aged 18–29 years.	Study did not discuss how survey participants were recruited or possible biases from the methods.
Kratochvil et al./ November 2020	Retrospective data analysis (Audit)	N/A	Psychiatric ED presentations in regional tertiary hospital (Geelong)	27 February to 25 March 2020 26 March to 22 April 2020	Before and during 1st wave	A 32% decrease in psychiatric presentations (316 vs. 214) in the 4 weeks post-stage 3 restrictions compared with the 4 weeks prior (pre = 316, post = 214).  Increased reporting of COVID-19 as a stressor (pre = 2%, post = 23%, $p < 0.01$ ), and reduction in reported suicidality (pre = 71%, post = 61%, $p = 0.01$ ).	No causal inferences or qualitative analysis possible.  No discussion of what identifying "COVID-19 as a stressor" referred to (e.g., direct impact of the virus or pandemic restrictions).

(continued)

**Table 1. Continued.**

Author(s) or Organization/ Publication date	Study design	Sample size	Target population	Study period	Pandemic context	Key results	Limitations
Cheek et al./23 July 2020	Retrospective data analysis	N/A	Pediatric ED presentations in two tertiary children's hospitals and two urban district hospitals, including attendance for mental health	Mid-March to mid-May 2020 compared with the same period in 2019	1st wave	<p>No differences in demographics (e.g., female sex, age) or other clinical characteristics.</p> <p>A 47.2% decrease in total ED presentations in 2020 compared with 2019, but a 35% (485 vs. 656) increase in mental health patients (daily means 8.0 vs. 10.8, difference 2.8, 95% CI 1.7–3.9, <math>p &lt; 0.001</math>).</p>	<p>No causal inferences or qualitative analysis possible.</p> <p>No data available regarding demographics or reasons for mental health presentation in relation to the pandemic.</p>

ED, emergency department; CI, confidence interval; N/A, not available; AH, Allied Health; NAPWHA, National Association of People with HIV Australia; K-10, 10-item Kessler Psychological Distress; GAD-7, Generalized Anxiety Disorder Scale; PHQ-9, Patient Health Questionnaire 9.

pregnant women, health care workers, older Australians, and children.

The articles included different study periods during 2020. Kratochvil et al.<sup>20</sup> and Cheek et al.<sup>21</sup> focused on the first epidemic wave of SARS-CoV-2 in March to May 2020. Rahman et al.,<sup>17</sup> Eadie et al.,<sup>22</sup> Weerasuria et al.,<sup>23</sup> Phillipou et al.,<sup>24</sup> Atmuri et al.,<sup>25</sup> and Siette et al.<sup>26</sup> studied the second wave during July to October 2020. Staples et al.,<sup>18</sup> Sacre et al.,<sup>27</sup> Dwyer et al.,<sup>28</sup> Czeisler et al.,<sup>15</sup> and Fisher et al.<sup>19</sup> included both the first and second waves. Finally, Digby et al.<sup>29</sup> and Holton et al.<sup>30</sup> conducted their studies between the first and second COVID-19 epidemic waves in 2020.

Generally, the included studies reported that the mental health of Victorians was negatively affected by the COVID-19 pandemic in 2020. However, variability and nuance were apparent amongst studies. In the discussion, we detail the general trends and variation in the findings of the included studies.

## Discussion

In this review, we found that during the COVID-19 pandemic in 2020, the general mental health and well-being of residents in Victoria decreased. However, the prevalence and degree of negative effects varied across the included studies. The evidence regarding the impact of pandemic restrictions on anxiety and depression levels was not consistent. Fisher et al. reported a significant increase in clinical depression (to 44.1%) and anxiety (to 32.3%) among the adult population during the lockdown implemented in the second wave of the COVID-19 pandemic in Victoria. These findings were nearly double those reported in other Australian states with less restrictive measures,<sup>19</sup> indicating a negative effect of restriction and social isolation on mental health. Two cross-sectional surveys

compared mental health among Victorians with residents of other Australian states and assessed the change in adult behaviors between the first and second waves in Victoria. Depressive and anxiety symptoms were assessed using the Patient Health Questionnaire 9 (PHQ-9) and Generalized Anxiety Disorder Scale (GAD-7), respectively, both of which are well-validated and reliable scales.<sup>31,32</sup> Czeisler et al.<sup>15</sup> found no significant difference in the prevalence of anxiety and depressive symptoms among adults in Victoria, based on two separate surveys in April and September 2020.<sup>15</sup> A potential reason for the differing findings of the studies by Czeisler and Fisher and their colleagues<sup>15,19</sup> is that the former study relied on longitudinal data from participants in a multi-time-point study, compared with Fisher's cross-sectional work.<sup>19</sup> It is possible that participants in the multi-time-point study were more pro-active about their mental health compared with participants in a single time-point project. This difference could also be explained by the findings of a cross-country study that found higher levels of anxiety, depression, and stress in Polish participants compared with Chinese participants.<sup>33</sup> The studies by Fisher and Czeisler and their colleagues<sup>15,19</sup> did not compare the prevalence of mental health parameters according to the specific cultural and ethnic backgrounds of participants. The population of Victoria is multicultural and ethnically diverse, which could contribute to the variability between these studies.

Poor mental health is associated with a variety of biopsychosocial variables including social isolation, alcohol use, and deficits in cognitive ability.<sup>34-37</sup> Czeisler et al.<sup>15</sup> demonstrated significant changes in the behavior of adults during the lockdown, such as having a considerably increased screen time and circumventing COVID-19-related information.<sup>15</sup> Additionally, 10% of participants reported new or

increased substance use (e.g., alcohol, legal/illegal drugs, and prescription drugs) to cope during the second wave of SARS-CoV-2 in Victoria.<sup>15</sup> This is consistent with studies in other countries that found increased alcohol use associated with social isolation and lockdown.<sup>38,39</sup> Further studies on alcohol consumption post-lockdown during the COVID-19 pandemic are needed to further quantify the extent of drug and alcohol use in the community.

In a retrospective study by Staples et al.,<sup>18</sup> the trends in mental health symptoms and use of the digital Mental Health service platform (MindSpot Clinic digital mental health service) were compared with pre-pandemic access to similar services.<sup>18</sup> Although those authors found a significant increase in the use of the digital mental health service during the first and second epidemic waves in Victoria, they did not observe any significant increase in psychological distress (10-item Kessler Psychological Distress, K-10), depression (PHQ-9), or anxiety (GAD-7) among users compared with pre-pandemic levels.<sup>18</sup> The authors argued that an increase in service demand was owing to Victorians with mild or moderate mental health symptoms accessing the service.<sup>18</sup> Therefore, whereas the number of people experiencing and seeking help for mental health symptoms increased, the average severity of symptoms did not. That study supports an earlier systematic review<sup>40</sup> examining global sub-syndromal anxiety, depression, and stress responses to the COVID-19 pandemic in Victoria.<sup>40</sup>

Our review highlights the importance of alternative service delivery, such as digital mental health services, to assist with high demands on mental health access during and after the strict COVID-19 pandemic suppression measures in Victoria. Staples et al.<sup>18</sup> also reported a significant decrease in the mean age of users of its digital mental health service, from age 35 years in the early weeks of the pandemic to age 32 years by

weeks 29 to 32.<sup>18</sup> Although it can be argued that the negative effect of SARS-CoV-2 on mental health could be a result of the disease itself, the literature in the Australian context shows that adverse mental health outcomes were minimally associated with direct SARS-CoV-2 exposure and strongly associated with restrictions that disrupted work and social functioning.<sup>41</sup>

More severe adverse mental health outcomes were commonly observed in women, carers, young people aged 18 to 29 years, people from low socioeconomic backgrounds or those with reduced work or unemployment owing to the pandemic, people with disabilities, and people with a previous psychiatric diagnosis.<sup>15,18,19</sup> The vulnerability of these populations to poor mental health outcomes is consistent with other studies in Australia<sup>42,43</sup> and other countries.<sup>44-48</sup> Children and adolescents were not immune to the negative impact of the COVID-19 pandemic on mental health. An audit of two tertiary children's hospitals and two urban district hospitals found a 35% increase in mental health-related presentations during the pandemic compared with the previous year whereas total presentations to the emergency department decreased by 47.2%.<sup>21</sup> Additionally, an Australian cross-sectional survey showed worse mental health deterioration in adolescents with a history of depression or anxiety during the pandemic.<sup>49</sup> Two systematic reviews confirmed the findings of studies conducted in Victoria and reported that children are more vulnerable to the negative effects of containment measures such as social distancing, isolation, and school closures, even though they may be less vulnerable to SARS-CoV-2 itself.<sup>50,51</sup> These studies revealed higher rates of anxiety, depression, and sleep and appetite disturbance among children and adolescents during the pandemic period. These findings are important for primary health care providers, policy makers, and governments to provide extra

training, education, and support for identifying and managing adolescents and younger adults at risk of adverse mental health outcomes. Further research is required to quantify the impacts of COVID-19 pandemic suppression strategies on children in Victoria.

The restrictive measures enforced during the second wave of the COVID-19 pandemic in Victoria have adversely affected other specific groups and populations. Increased psychological distress symptoms have been reported in 25% of frontline health care workers in hospitals throughout Victoria.<sup>17,29,30</sup> This could be explained by the relatively higher exposure rates of health care staff working in hospitals with suspected and confirmed cases of COVID-19.<sup>52</sup> Improving understanding of the virus as well as high vaccination rates among health care staff may reshape the initial response of frontline health care workers to the pandemic. Further research is required to evaluate the longer-term mental health challenges among frontline workers owing to the COVID-19 pandemic. People with specific diseases such as diabetes were concerned about engaging in leisure activities, uncertainty about the future, and emotional well-being,<sup>27</sup> and people living with HIV reported a negative impact on their relationships owing to the pandemic.<sup>23</sup> The impact of COVID-19 pandemic restrictions on personal relationships is a global problem,<sup>53–55</sup> highlighting the need for strategies to improve social connectedness, optimism, and quality of life.

Several strengths and limitations of this review should be considered. Of the 15 articles found using the search strategy above, only three studies (those of Fisher et al.,<sup>19</sup> Czeisler et al.,<sup>15</sup> and Staples et al.<sup>18</sup>) were generalizable to adults in Victoria without being significantly biased toward a specific population group. Many studies were cross-sectional analyses that lacked comparative baseline data;

therefore, causal inferences could not be made. The study by Fisher et al.<sup>19</sup> has several advantages over other similar studies. That study had the largest sample size ( $n = 23,749$ ) and the sample was representative of the general adult population in Victoria. Their study also used national baseline data from the first survey, allowing for a measurable difference to be determined in comparison with the second survey, which was initiated during the second COVID-19 wave in Victoria. However, the study by Fisher et al.<sup>19</sup> also has some limitations, including a lack of discussion regarding the recruitment process (e.g., how the survey was advertised or accessed), which could limit its generalizability. Similar to that research, the design of the study by Czeisler et al.<sup>15</sup> could not demonstrate causality regarding mental health findings and relied on self-reported outcomes rather than diagnostic interviews. The work by Staples et al.<sup>18</sup> had inherent population biases and could only provide information about Victorians who actively chose to engage with its service, who were probably more psychologically-minded and mentally well enough to be motivated to seek help. Lastly, we did not use a measure of bias or study quality when reviewing the included articles.

## Conclusion

In the present review, we found that the general mental health of residents in Victoria, Australia was negatively affected by the COVID-19 pandemic restrictions during 2020. Victorians demonstrated mental health help-seeking behavior and used coping strategies in response to the restrictions, such as increased alcohol use. Although the literature is sparse, and in particular, there are insufficient data during the second wave of SARS-CoV-2, the study findings are consistent regarding the groups that were most vulnerable to

adverse mental health outcomes as a result of Victoria's COVID-19 pandemic suppression strategies.

This review identified several populations that are at risk of poorer mental health outcomes, including women, children and young people, carers, and people with disabilities. Demographic risk factors also include pre-existing psychiatric health conditions, low socioeconomic status, unemployment, and marginalized minority groups. These groups were mentioned consistently in the included studies in relation to Victoria, as well as globally. This highlights the importance of future research, policy interventions, and strategies to address poor mental health in the population, particularly focusing on these specific groups in Victoria, as they are most vulnerable to the negative effects of pandemic restrictions.

This review encompasses studies focusing on mental health outcomes in the 2020 COVID-19 pandemic context. During 2021, Victoria experienced recurrent lockdowns and became the city with the most days of lockdown globally (262 days).<sup>56</sup> 2021 presented new challenges such as “pandemic fatigue,” with one review finding decreasing adherence to some COVID-19 pandemic suppression strategies, such as physical distancing, as the pandemic continued.<sup>57</sup> Another study suggested that poorer personal resilience and coping skills are associated with lockdown fatigue.<sup>58</sup> Studies conducted in the context of the 2021 pandemic are lacking, with ongoing enforcement of COVID-19 pandemic suppression strategies, increased knowledge of the virus, and the roll-out of COVID-19 vaccines. As the pandemic continues to unfold in Victoria, in Australia, and globally, alongside ever-changing COVID-19 pandemic suppression strategies and vaccination strategies, ongoing assessment of the impact on population mental health will

continue to be important issues in future research.

### Acknowledgements

The authors would like to thank Anna Lovang (St Vincent's Hospital Library Service) for her assistance in refining the literature search.

### Declaration of conflicting interests

The authors declare that there is no potential conflict of interest with respect to the research, authorship, and/or publication of this article.

### Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

### ORCID iDs

Jacqueline Jiang  <https://orcid.org/0000-0003-1682-7785>

Darren Haywood  <https://orcid.org/0000-0002-9317-4135>

### References

1. WHO Coronavirus (COVID-19) Dashboard. Geneva: World Health Organization; 2021. Available from: <https://covid19.who.int/table>.
2. World Health Organization. *Weekly epidemiological update – 29 December 2020*. Geneva: World Health Organization, Response E; 2020.
3. Alfano V and Ercolano S. The Efficacy of Lockdown Against COVID-19: A Cross-Country Panel Analysis. *Appl Health Econ Health Policy* 2020; 18: 509–517.
4. World Health Organization. *Mental health and psychosocial considerations during the COVID-19 outbreak*. Geneva: World Health Organization; 2020. Available from: <https://www.who.int/docs/default-source/coronavirus/mental-health-considerations.pdf>.
5. Rossi R, Soggi V, Talevi D, et al. COVID-19 Pandemic and Lockdown Measures Impact on Mental Health Among the General Population in Italy. *Front Psychiatry* 2020; 11: 790.

6. Elmer T, Mepham K and Stadtfeld C. Students under lockdown: Comparisons of students' social networks and mental health before and during the COVID-19 crisis in Switzerland. *PLoS One* 2020; 15: e0236337.
7. Chaudhry R, Dranitsaris G, Mubashir T, et al. A country level analysis measuring the impact of government actions, country preparedness and socioeconomic factors on COVID-19 mortality and related health outcomes. *EClinicalMedicine* 2020; 25: 100464.
8. Saul A, Scott N, Crabb BS, et al. Impact of Victoria's Stage 3 lockdown on COVID-19 case numbers. *Med J Aust* 2020; 213: 494–496.e1.
9. Health and Human Services. Coronavirus update for Victoria – 30 November. Melbourne: Department of Health and Human Services, 30 November 2020. Available from: <https://www.dhhs.vic.gov.au/coronavirus-update-victoria-30-november-2020>.
10. Storen RC and Corrigan N. COVID-19: a chronology of state and territory government announcements (up until 30 June 2020). In: Services DoP, editor. Parliamentary Library & Information Service, Parliament of Victoria. 22 October 2020. Available from: [https://www.aph.gov.au/About\\_Parliament/Parliamentary\\_departments/Parliamentary\\_Library/pubs/rp/rp2021/Chronologies/COVID-19StateTerritoryGovernmentAnnouncements](https://www.aph.gov.au/About_Parliament/Parliamentary_departments/Parliamentary_Library/pubs/rp/rp2021/Chronologies/COVID-19StateTerritoryGovernmentAnnouncements)
11. Health and Human Services. *Media hub – coronavirus (COVID-19)*. Melbourne: Department of Health and Human Services, 20 September 2021.
12. McLean H and Huf B. Emergency Powers, Public Health and COVID-19. In: Services DoP, editor. Parliamentary Library & Information Service, Parliament of Victoria; 2020. p. 37.
13. Health and Human Services. Victorian COVID-19 data. Melbourne: Department of Health and Human Services; 20 September 2021. Available from: <https://www.coronavirus.vic.gov.au/victorian-coronavirus-covid-19-data>.
14. Zhuang Y and Cave D. The Lockdown That Felt Like It Might Last Forever Has Finally Ended 22 October 2020; updated 22 October 2021. Available from: <https://www.nytimes.com/2020/10/28/world/australia/melbourne-lockdown-ends.html>.
15. Czeisler MÉ, Wiley JF, Facer-Childs ER, et al. Mental health, substance use, and suicidal ideation during a prolonged COVID-19-related lockdown in a region with low SARS-CoV-2 prevalence. *J Psychiatr Res* 2021; 140: 533–544.
16. Veritas Health Innovation. *Covidence systematic review software*. Melbourne, Australia: Covidence; 2021.
17. Rahman MA, Hoque N, Alif SM, et al. Factors associated with psychological distress, fear and coping strategies during the COVID-19 pandemic in Australia. *Global Health* 2020; 16: 95.
18. Staples L, Nielssen O, Kayrouz R, et al. Rapid Report 3: Mental health symptoms, characteristics, and regional variation, for users of an Australian digital mental health service during the first 8 months of COVID-19. *Internet Interv* 2021; 24: 100378.
19. Fisher J, Tran T, Hammarberg K, et al. Quantifying the mental health burden of the most severe covid-19 restrictions: A natural experiment. *J Affect Disord* 2021; 293: 406–414.
20. Kratochvil D, Hill H, Moylan S. The impact of Stage 3 COVID-19 lockdown on psychiatric presentations at a regional Victorian emergency department. *Australasian psychiatry: Bulletin of Royal Australian and New Zealand College of Psychiatrists*. 2021; 29(1): 105–106. DOI: 10.1177/1039856220968390
21. Cheek JA, Craig SS, West A, et al. Emergency department utilisation by vulnerable paediatric populations during the COVID-19 pandemic. *Emerg Med Australas* 2020; 32: 870–871.
22. Eadie P, Levickis P, Murray L, Page J, Elek C, Church A. Early Childhood Educators' Wellbeing During the COVID-19 Pandemic. *Early Childhood Education Journal*. 2021: 1–11. DOI: 10.1007/s10643-021-01203-3
23. Weerasuria M, Ko C, Ehm A, et al. The Impact of the COVID-19 Pandemic on People Living with HIV in Victoria, Australia. *AIDS Res Hum Retroviruses* 2021; 37: 322–328.

24. Phillipou A, Tan EJ, Toh WL, Van Rheenen TE, Meyer D, Neill E, et al. Pet ownership and mental health during COVID-19 lockdown. *Australian Veterinary Journal*. 2021. DOI: 10.1111/avj.13102
25. Atmuri K, Sarkar M, Obudu E, Kumar A. Perspectives of pregnant women during the COVID-19 pandemic: A qualitative study. *Women and Birth: Journal of the Australian College of Midwives*. 2021. DOI: 10.1016/j.wombi.2021.03.008
26. Siette J, Seaman K, Dodds L, Ludlow K, Johnco C, Wuthrich V, et al. A national survey on COVID-19 second-wave lockdowns on older adults' mental wellbeing, health-seeking behaviours and social outcomes across Australia. *BMC Geriatrics*. 2021; 21(1): 400. DOI: 10.1186/s12877-021-02352-1
27. Sacre JW, Holmes-Truscott E, Salim A, et al. Impact of the COVID-19 pandemic and lockdown restrictions on psychosocial and behavioural outcomes among Australian adults with type 2 diabetes: Findings from the PREDICT cohort study. *Diabet Med* 2021; 38: e14611.
28. Dwyer J, Dwyer J, Hiscock R, O'Callaghan C, Taylor K, Millar C, et al. COVID-19 as a context in suicide: early insights from Victoria, Australia. *Australian and New Zealand Journal of Public Health*. 2021. DOI: 10.1111/1753-6405.13132
29. Digby R, Winton-Brown T, Finlayson F, et al. Hospital staff well-being during the first wave of COVID-19: Staff perspectives. *Int J Ment Health Nurs* 2021; 30: 440–450.
30. Holton S, Wynter K, Trueman M, et al. Psychological well-being of Australian hospital clinical staff during the COVID-19 pandemic. *Aust Health Rev* 2021; 45: 297–305.
31. Kroenke K, Spitzer RL and Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001; 16: 606–613.
32. Löwe B, Decker O, Müller S, et al. Validation and Standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the General Population. *Med Care* 2008; 46: 266–274.
33. Wang C, Chudzicka-Czupala A, Grabowski D, et al. The Association Between Physical and Mental Health and Face Mask Use During the COVID-19 Pandemic: A Comparison of Two Countries With Different Views and Practices. *Front Psychiatry* 2020; 11: 569981.
34. Haywood D and Baughman F. Multidimensionality in Executive Function Profiles in Schizophrenia: a Computational Approach Using the Wisconsin Card Sorting Task. *Computational Brain & Behavior* 2021; 4: 381–394.
35. Chou KL, Liang K and Sareen J. The association between social isolation and DSM-IV mood, anxiety, and substance use disorders: wave 2 of the National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry* 2011; 72: 1468–1476.
36. Haywood D, Baughman FD, Mullan BA, et al. What Accounts for the Factors of Psychopathology? An Investigation of the Neurocognitive Correlates of Internalising, Externalising, and the p-Factor. *Brain Sci* 2022; 12: 421.
37. Haywood D, Baughman FD, Mullan BA, et al. Neurocognitive Artificial Neural Network Models Are Superior to Linear Models at Accounting for Dimensional Psychopathology. *Brain Sci* 2022; 12: 1060.
38. Arora T and Grey I. Health behaviour changes during COVID-19 and the potential consequences: A mini-review. *J Health Psychol* 2020; 25: 1155–1163.
39. Naughton F, Ward E, Khondoker M, et al. Health behaviour change during the UK COVID-19 lockdown: Findings from the first wave of the C-19 health behaviour and well-being daily tracker study. *Br J Health Psychol* 2021; 26: 624–643.
40. Rajkumar RP. COVID-19 and mental health: A review of the existing literature. *Asian J Psychiatr* 2020; 52: 102066.
41. Dawel A, Shou Y, Smithson M, et al. The Effect of COVID-19 on Mental Health and Wellbeing in a Representative Sample of Australian Adults. *Front Psychiatry* 2020; 11: 579985.
42. Stanton R, To QG, Khalesi S, et al. Depression, Anxiety and Stress during



- COVID-19: Associations with Changes in Physical Activity, Sleep, Tobacco and Alcohol Use in Australian Adults. *Int J Environ Res Public Health* 2020; 17: 4065.
43. Newby JM, O'Moore K, Tang S, et al. Acute mental health responses during the COVID-19 pandemic in Australia. *PLoS One* 2020; 15: e0236562.
  44. Bäuerle A, Teufel M, Musche V, et al. Increased generalized anxiety, depression and distress during the COVID-19 pandemic: a cross-sectional study in Germany. *J Public Health (Oxf)* 2020; 42: 672–678.
  45. Gasteiger N, Vedhara K, Massey A, et al. Depression, anxiety and stress during the COVID-19 pandemic: results from a New Zealand cohort study on mental well-being. *BMJ Open* 2021; 11: e045325.
  46. Varma P, Junge M, Meaklim H, et al. Younger people are more vulnerable to stress, anxiety and depression during COVID-19 pandemic: A global cross-sectional survey. *Prog Neuropsychopharmacol Biol Psychiatry* 2021; 109: 110236.
  47. Fancourt D, Steptoe A and Bu F. Trajectories of anxiety and depressive symptoms during enforced isolation due to COVID-19 in England: a longitudinal observational study. *Lancet Psychiatry* 2021; 8: 141–149.
  48. El-Zoghby SM, Soltan EM and Salama HM. Impact of the COVID-19 Pandemic on Mental Health and Social Support among Adult Egyptians. *J Community Health* 2020; 45: 689–695.
  49. Li SH, Beames JR, Newby JM, et al. The impact of COVID-19 on the lives and mental health of Australian adolescents. *Eur Child Adolesc Psychiatry* 2021; 31: 1–13.
  50. Meherali S, Punjani N, Louie-Poon S, et al. Mental Health of Children and Adolescents Amidst COVID-19 and Past Pandemics: A Rapid Systematic Review. *Int J Environ Res Public Health* 2021; 18: 3432.
  51. Nearchou F, Flinn C, Niland R, et al. Exploring the Impact of COVID-19 on Mental Health Outcomes in Children and Adolescents: A Systematic Review. *Int J Environ Res Public Health* 2020; 17: 8479.
  52. O'Reilly GM, Mitchell RD, Mitra B, et al. Epidemiology and clinical features of emergency department patients with suspected COVID-19: Insights from Australia's 'second wave' (COVED-4). *Emerg Med Australas* 2021; 33: 331–342.
  53. Goodwin R, Hou WK, Sun S, et al. Quarantine, distress and interpersonal relationships during COVID-19. *Gen Psychiatr* 2020; 33: e100385.
  54. Sheffler JL, Joiner TE and Sachs-Ericsson NJ. The Interpersonal and Psychological Impacts of COVID-19 on Risk for Late-Life Suicide. *Gerontologist* 2021; 61: 23–29.
  55. Pieh C, O'Rourke T, Budimir S, et al. Relationship quality and mental health during COVID-19 lockdown. *PLoS One* 2020; 15: e0238906.
  56. Zhuang Y. Melbourne, after 262 days in lockdown, celebrates a reopening. *The New York Times* 22 October 2021.
  57. Petherick A, Goldszmidt R, Andrade EB, et al. A worldwide assessment of changes in adherence to COVID-19 protective behaviours and hypothesized pandemic fatigue. *Nat Hum Behav* 2021; 5: 1145–1160.
  58. Labrague LJ and Ballad CA. Lockdown fatigue among college students during the COVID-19 pandemic: Predictive role of personal resilience, coping behaviors, and health. *Perspect Psychiatr Care* 2021; 57: 1905–1912.