

GENERALIZATIONS OF SOME SIMPLE CONGRUENCES

R. S. Melham and A. G. Shannon

University of Technology, Sydney 2007, Australia
(Submitted July 1993)

1. INTRODUCTION

Over many years in this journal there have appeared results concerning congruence and divisibility in relation to the Fibonacci and Lucas numbers. Here we take four such results and translate them to sequences which generalize the Fibonacci and Lucas sequences.

We hope that the nature of our results will demonstrate to the beginning Fibonacci enthusiast that there is scope to obtain further generalizations of a similar nature.

2. THE SEQUENCES

In the notation of Horadam [7] write

$$W_n = W_n(a, b; p, q), \quad (2.1)$$

meaning that

$$W_n = pW_{n-1} - qW_{n-2}, \quad W_0 = a, W_1 = b, \quad n \geq 2. \quad (2.2)$$

We assume throughout that a, b, p, q are integers.

The auxiliary equation associated with (2.2) is

$$x^2 - px + q = 0, \quad (2.3)$$

whose roots

$$\alpha = \frac{p + \sqrt{p^2 - 4q}}{2}, \quad \beta = \frac{p - \sqrt{p^2 - 4q}}{2} \quad (2.4)$$

are assumed distinct. We write

$$\Delta = (\alpha - \beta)^2 = p^2 - 4q. \quad (2.5)$$

We shall be concerned with specializations of the following two sequences:

$$\begin{cases} U_n = W_n(0, 1; p, q), \\ V_n = W_n(2, p; p, q). \end{cases} \quad (2.6)$$

The sequences $\{U_n\}$ and $\{V_n\}$ are the fundamental and primordial sequences, respectively, generated by (2.2). They are natural generalizations of the Fibonacci and Lucas sequences and have been studied extensively, particularly by Lucas [11]. Further information can be found, for example, in [1], [7], and [10].

The Binet forms for U_n and V_n are

$$U_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad (2.7)$$

$$V_n = \alpha^n + \beta^n. \quad (2.8)$$

These sequences can be extended to negative subscripts by the use of the recurrence (2.2) or the Binet forms.

We will make use of the following well-known results which we state for easy reference:

$$q^n U_{-n} = -U_n, \tag{2.9}$$

$$q^n V_{-n} = V_n, \tag{2.10}$$

$$U_{2n} = U_n V_n, \tag{2.11}$$

$$\text{if } m|n \text{ then } U_m | U_n. \tag{2.12}$$

The following identities, which occur in Bergum and Hoggatt [1], will also be needed:

$$U_{n+k} + q^k U_{n-k} = U_n V_k, \tag{2.13}$$

$$U_{n+k} - q^k U_{n-k} = U_k V_n, \tag{2.14}$$

$$V_{n+k} + q^k V_{n-k} = V_n V_k, \tag{2.15}$$

$$V_{n+k} - q^k V_{n-k} = \Delta U_n U_k. \tag{2.16}$$

The sequences

$$\begin{cases} U_n = W_n(0, 1; p, -1), \\ V_n = W_n(2, p; p, -1), \end{cases} \tag{2.17}$$

are an important subclass of the sequences (2.6) and can be looked upon as an intermediate level of generalization of the Fibonacci and Lucas numbers in which $p = 1$. The specializations $p = 2$ and $p = 2x$ also yield cases of interest. For $p = 2$ see [4], [8], [15] and for $p = 2x$ see [9], [12], [13].

We use the $U_n - V_n$ notation throughout to refer to the sequences (2.6) and to the sequences (2.17). There will be no ambiguity since we shall always indicate the set to which we are referring.

3. CONGRUENCE RESULT I

Singh [17] gives the following:

$$L_{2^n} \equiv 7 \pmod{40} \text{ for } n \geq 2. \tag{3.1}$$

Generalization: Let $\{U_n\}$ and $\{V_n\}$ be the sequences defined in (2.17). Then

$$V_{2^n} \equiv V_4 \pmod{\Delta U_6 U_{10}}, \quad n = 2, 4, 6, \dots, \tag{3.2}$$

$$V_{2^n} \equiv V_8 \pmod{\Delta U_6 U_{20} V_2 (V_4 - 1)}, \quad n = 3, 5, 7, \dots, \tag{3.3}$$

and so

$$V_{2^n} \equiv V_4 \pmod{\Delta U_2 U_6} \text{ for } n \geq 2. \tag{3.4}$$

Proof: We shall use the following, all of which can be proved using Binet forms:

$$V_{2^{k+1}} = V_{2^k}^2 - 2, \tag{3.5}$$

$$p(V_4 + 1) = U_6, \tag{3.6}$$

$$V_4 - 2 = \Delta p^2, \quad (3.7)$$

$$p(V_4^2 + V_4 - 1) = U_{10}, \quad (3.8)$$

$$p(V_8 + 1) = U_6(V_4 - 1), \quad (3.9)$$

$$\Delta p^2 V_2^2 = V_8 - 2, \quad (3.10)$$

$$pV_2(V_8^2 + V_8 - 1) = U_{20}. \quad (3.11)$$

Using (3.5) twice, we also obtain

$$V_{2^{k+2}} = V_{2^k}^4 - 4V_{2^k}^2 + 2. \quad (3.12)$$

Now (3.2) is true for $n = 2$ and if it is true for $n = k$ (even) then by (3.12) and the induction hypothesis

$$V_{2^{k+2}} = V_{2^k}^4 - 4V_{2^k}^2 + 2 \equiv V_4^4 - 4V_4^2 + 2 \pmod{\Delta U_6 U_{10}}.$$

But by (3.6)-(3.8),

$$(V_4^4 - 4V_4^2 + 2) - V_4 = (V_4 + 1)(V_4 - 2)(V_4^2 + V_4 - 1) = \Delta U_6 U_{10}.$$

This proves (3.2). Congruence (3.3) can be proved similarly by making use of (3.9)-(3.12).

From (2.16) we see that $V_8 - V_4 = \Delta U_2 U_6$, and (2.12) shows that $\Delta U_2 U_6$ divides both moduli in (3.2) and (3.3). This proves (3.4). \square

Putting $V_n = L_n$ so that $U_n = F_n$, we see that (3.4) reduces to (3.1).

4. CONGRUENCE RESULT II

Berzsényi [2] states that

$$F_{6n+1}^2 \equiv 1 \pmod{24}, \quad n \text{ an integer.} \quad (4.1)$$

Generalization: Let $\{U_n\}$ and $\{V_n\}$ be the sequences defined in (2.17). Then

$$U_{6n+1}^2 \equiv 1 \pmod{U_4 U_6}, \quad n \text{ an integer.} \quad (4.2)$$

Proof:

$$\begin{aligned} U_{6n+1}^2 - 1 &= (U_{6n+1} - U_1)(U_{6n+1} + U_1) \\ &= (U_{3n+1+3n} - U_{3n+1-3n})(U_{3n+1+3n} + U_{3n+1-3n}) \\ &= U_{3n} V_{3n} U_{3n+1} V_{3n+1}, \end{aligned} \quad (4.3)$$

where we have used (2.13) and (2.14) with $q = -1$.

Taking m to be an integer, we consider two cases:

Case 1. $n = 2m + 1$. Using (2.11), the right side of (4.3) becomes $U_{12m+6} U_{12m+8}$. Then by (2.12), $U_4 | U_{12m+8}$ and $U_6 | U_{12m+6}$ and (4.2) follows.

Case 2. $n = 2m$. Using (2.11), the right side of (4.3) becomes $U_{12m} U_{12m+2}$. Since $U_4 | U_{12m}$, $U_6 | U_{12m}$, and $(U_4, U_6) = U_2 | U_{12m+2}$, then $U_4 U_6 | U_{12m} U_{12m+2}$ and (4.2) follows.

This completes the proof of (4.2). \square

5. CONGRUENCE RESULT III

Freitag [5] gives the following:

$$L_{2p^k} \equiv 3 \pmod{10} \tag{5.1}$$

for all primes $p \geq 5$ and natural numbers k . We caution against confusing the prime p with the parameter p .

Generalization: Let $\{U_n\}$ and $\{V_n\}$ be the sequences defined in (2.17). Then

$$V_{n+12} \equiv V_n \pmod{40}. \tag{5.2}$$

Proof: Using (2.16), we see that

$$V_{n+12} - V_n = (p^2 + 4)U_6U_{n+6} = p(p^2 + 1)(p^2 + 3)(p^2 + 4)U_{n+6}. \tag{5.3}$$

The right side of (5.3) is divisible by 40 since 5 divides either p , $p^2 + 1$, or $p^2 + 4$ and $8|p(p^2 + 4)$ if p is even while $8|(p^2 + 1)(p^2 + 3)$ if p is odd.

To see that (5.2) generalizes (5.1) we note that, as observed in Bruckman [3],

$$2p^k \equiv 2 \text{ or } -2 \pmod{12} \tag{5.4}$$

for all primes $p \geq 5$. Now, since $L_2 = L_{-2} = 3$, (5.1) follows from (5.2) and (5.4). \square

6. A DIVISIBILITY RESULT

Grassi [6] gives the following:

$$12|(F_{4n-2} + F_{4n} + F_{4n+2}), \tag{6.1}$$

$$168|(F_{8n-4} + F_{8n} + F_{8n+4}). \tag{6.2}$$

Generalization: Let $\{U_n\}$ and $\{V_n\}$ be the sequences defined in (2.6). Then for $n \geq 0, k \geq 1$,

$$U_{2k-1}V_{4k-2}V_{6k-3}|(q^{4k-2}U_{(4k-2)(2n-1)} - q^{2k-1}U_{(8k-4)n} + U_{(4k-2)(2n+1)}), \tag{6.3}$$

$$V_{2k}V_{4k}U_{6k}|(q^{4k}U_{4k(2n-1)} + q^{2k}U_{8kn} + U_{4k(2n+1)}). \tag{6.4}$$

Proof: We prove (6.3) by using reasoning similar to Mana [14]. Fixing k and denoting the dividend by $G_n^{(k)}$ we have, by (2.9),

$$\begin{aligned} G_0^{(k)} &= q^{4k-2}U_{-(4k-2)} + U_{4k-2} \\ &= -U_{4k-2} + U_{4k-2} = 0. \end{aligned}$$

Also

$$\begin{aligned} G_1^{(k)} &= q^{4k-2}U_{4k-2} - q^{2k-1}U_{8k-4} + U_{12k-6} \\ &= q^{4k-2}U_{2k-1}V_{2k-1} + U_{2k-1}V_{10k-5} \quad [\text{by (2.11) and (2.14)}] \\ &= U_{2k-1}V_{6k-3}V_{4k-2} \quad [\text{by (2.15)}]. \end{aligned}$$

Now $\{G_n^{(k)}\}$ can be regarded as the sum of three sequences each satisfying the same homogeneous linear second-order recurrence relation with integer coefficients (see Shannon and Horadam [16]).

Hence, $\{G_n^{(k)}\}$ also satisfies this second-order recurrence. Therefore, since $U_{2k-1}V_{4k-2}V_{6k-3}|G_0^{(k)}$ and $U_{2k-1}V_{4k-2}V_{6k-3}|G_1^{(k)}$, then $U_{2k-1}V_{4k-2}V_{6k-3}|G_n^{(k)}$ for all $n \geq 0$. Since k was arbitrary, the proof of (6.3) is complete. The proof of (6.4) is similar. \square

Taking $\{U_n\} = \{F_n\}$, $\{V_n\} = \{L_n\}$ and putting $k = 1$, we see that (6.3) and (6.4) reduce to (6.1) and (6.2), respectively.

7. CONCLUDING COMMENTS

We have chosen an assortment of results requiring essentially different methods of proof. For the most part, the moduli or divisors in question are products of terms from the relevant sequences. We feel that with this observation there is scope for the beginner to discover generalizations of a similar nature.

ACKNOWLEDGMENT

We gratefully acknowledge the comments of an anonymous referee whose suggestions led to the final form of most of our results and provided other ways of streamlining this paper.

REFERENCES

1. G. E. Bergum & V. E. Hoggatt, Jr. "Sums and Products for Recurring Sequences." *The Fibonacci Quarterly* **13.2** (1975):115-20.
2. G. Berzsenyi. Problem B-331. *The Fibonacci Quarterly* **14.2** (1976):188.
3. P. S. Bruckman. Solution to Problem B-314. *The Fibonacci Quarterly* **14.3** (1976):288.
4. J. Ercolano. "Matrix Generators of Pell Sequences." *The Fibonacci Quarterly* **17.1** (1979): 71-77.
5. H. T. Freitag. Problem B-314. *The Fibonacci Quarterly* **13.3** (1975):285.
6. R. M. Grassi. Problems B-202 and B-203. *The Fibonacci Quarterly* **9.1** (1971):106.
7. A. F. Horadam. "Basic Properties of a Certain Generalized Sequence of Numbers." *The Fibonacci Quarterly* **3.2** (1965):161-76.
8. A. F. Horadam. "Pell Identities." *The Fibonacci Quarterly* **9.3** (1971):245-52, 263.
9. A. F. Horadam & Bro. J. M. Mahon. "Pell and Pell-Lucas Polynomials." *The Fibonacci Quarterly* **23.1** (1985):7-20.
10. D. Jarden. *Recurring Sequences*. Jerusalem: Riveon Lematematika, 1966.
11. E. Lucas. *Théorie des Nombres*. Paris: Albert Blanchard, 1961.
12. Bro. J. M. Mahon & A. F. Horadam. "Matrix and Other Summation Techniques for Pell Polynomials." *The Fibonacci Quarterly* **24.3** (1986):290-309.
13. Bro. J. M. Mahon & A. F. Horadam. "Pell Polynomial Matrices." *The Fibonacci Quarterly* **25.1** (1987):21-28.
14. P. Mana. Solution to Problem B-202. *The Fibonacci Quarterly* **9.5** (1971):547.
15. C. Serkland. "Generating identities for Pell Triples." *The Fibonacci Quarterly* **12.2** (1974): 121-28.
16. A. G. Shannon & A. F. Horadam. "Special Recurrence Relations Associated with the Sequences $\{W_n(a, b, p, q)\}$." *The Fibonacci Quarterly* **17.4** (1979):294-99.
17. S. Singh. Problem B-694. *The Fibonacci Quarterly* **29.3** (1991):277.

AMS Classification Numbers: 11B37, 11B39



The Fibonacci Quarterly

Official Publication of The Fibonacci Association



[Journal Home](#) | [Editorial Board](#) | [List of Issues](#)
[How to Subscribe](#) | [General Index](#) | [Fibonacci Association](#)

Volume 33

Number 2

May 1995

CONTENTS

[Cover Page](#)

- G. F. C. de Bruyn
Formulas $a + a^2 2^p + a^3 3^p + \dots + a^n n^p$ 98
[Full text](#)
- Wai-fong Chuan
Generating Fibonacci Words 104
[Full text](#)
- Wai-fong Chuan
Extraction Property of the Golden Sequence 113
[Full text](#)
- A. Rotkiewicz and K. Ziemak
On Even Pseudoprimes 123
[Full text](#)
- R. S. Melham and A. G. Shannon
Generalizations of Some Simple Congruences 126
[Full text](#)
- P. R. Subramanian
Nonzero Zeros of the Hermite Polynomials are Irrational 131
[Full text](#)
- R. S. Melham and A. G. Shannon
A Generalization of a Result of D'Ocagne 135
[Full text](#)
- Helmut Prodinger
Geometric Distributions and Forbidden Subwords 139
[Full text](#)
- Ray Melham and Derek Jennings
On the General Linear Recurrence Relation 142
[Full text](#)
- N. G. Gamkrelidze
On a Probabilistic Property of the Fibonacci Sequence 147

[Full text](#)

Arnold Knopfmacher and M. E. Mays
Pierce Expansions of Ratios and Fibonacci and Lucas Numbers and Polynomials 153

[Full text](#)

Peter Hope
Exponential Growth of Random Fibonacci Sequences 164

[Full text](#)

L. C. Hsu
A Difference-Operational Approach to the Möbius Inversion Formulas 169

[Full text](#)

Jun Wang
On the k^{th} Derivative Sequences of Fibonacci and Lucas Polynomials 174

[Full text](#)

Krystyna Bialek
A Note on Choudhry's Results 179

[Full text](#)

Edited by Stanley Rabinowitz
Elementary Problems and Solutions 181

[Full text](#)

Edited by Raymond E. Whitney
Advanced Problems and Solutions 187

[Full text](#)

[Back Cover](#)

Copyright © 2010 [The Fibonacci Association](http://www.fq.math.ca/). All rights reserved.