

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

How simple can a treatment for early stuttering be? A proposed two-factor early intervention

Mark Onslow,<sup>a</sup> Anna Hearne, Kun Yu,<sup>b</sup> Susan O’Brian,<sup>a</sup> Ross Menzies,<sup>a</sup> and Robyn Lowe<sup>a</sup>

<sup>a</sup> University of Technology Sydney, Australian Stuttering Research Centre, New South Wales, Australia

<sup>b</sup> University of Technology Sydney, Data Science Institute, New South Wales, Australia

\* Corresponding Author: Mark Onslow, Australian Stuttering Research Centre, University of Technology Sydney, Building 1, 15 Broadway, Ultimo, NSW, 2007, AUSTRALIA  
e-mail: Mark.Onslow@uts.edu.au

*Keywords:* stuttering, treatment, preschool

**Declaration of Interest Statement**

The author reports no conflict of interest.

**Abstract**

26

27     *Purpose:* In this paper we draw attention to the need for immediate intervention shortly after  
28 stuttering onset. More than half of stuttering onsets occur before 3 years of age. We argue that  
29 existing interventions for preschool children who stutter require varying levels of cognitive  
30 engagement from children; hence, they are not suitable for children of that age. For this reason, we  
31 argue, there is no clinical trials evidence for treatment efficacy with children younger than 3 years  
32 of age.

33     *Conclusion:* There are many recommended parent strategies that do not require any active  
34 participation from children, which, therefore, may be suitable for immediate stuttering intervention.  
35 Two of these have laboratory support in their favor: parent speech rate reduction and increased  
36 interturn speaker latency. Therefore, we developed a clinical protocol based on those two parent  
37 strategies and showed the clinical viability of the protocol with three children who stuttered. We  
38 argue that automation of our proposed treatment with lifelike AI-generated avatar clinicians will  
39 make it globally viable, and a suitable target for future Phase I–IV clinical trials.

40 **Quality of life and illness burden of stuttering**

41 Stuttering occurs in all cultures and languages. Based on predominantly Western cultures,  
42 Bloodstein et al. (2021) conclude that lifetime incidence is 10% with a prevalence of 1–2%. At the  
43 time of writing, this prevalence suggests that 80–160 million people stutter globally. The condition  
44 is associated with significant quality-of-life impairment in the range of that reported for diabetes,  
45 cardiovascular disease, and HIV (Norman et al., 2024). Presumably, this is because, unlike those  
46 conditions, stuttering begins during early childhood and its effects are experienced throughout the  
47 lifespan. Stuttering is associated with substantive cost of illness. Norman et al. (2024) showed that  
48 in direct terms, willingness to pay for effective treatment can be 2–4 times the family annual  
49 income, and 61% of families report a financial strain because of an adolescent who stutters. Indirect  
50 costs of illness are social, involving failure of those who stutter to attain their full occupational  
51 potential, which has been shown in several cohort studies (Gerlach et al., 2018; Jacobs et al., 2025;  
52 McAllister et al., 2012). Failure to attain occupational potential may be partly related to childhood  
53 educational disadvantage because of stuttering (Berchiatti et al., 2020; Boyle et al., 1994; Williams  
54 et al., 1969).

55 The lifetime effects of stuttering include speech-related anxiety. Social anxiety scores of those  
56 who stutter are nearly a standard deviation above nonstuttering controls (Craig & Tran, 2014).  
57 Compared with controls, those who stutter have greatly increased odds for social anxiety disorder  
58 (Iverach et al., 2009). During childhood, 24% of 7–12-year-old children who stutter may be  
59 diagnosed with social anxiety disorder in comparison to 4.6% of control children (Iverach et al.  
60 2016). There is evidence that this anxiety originates early in life during the pre-school years (Briley  
61 et al., 2019; McAllister, 2016; Tığrak et al., 2020). Possibly, this is a reason why quality-of-life  
62 impairment has been measured during the preschool years (Norman et al., 2024).

63 **A Consensus About Early Stuttering Intervention**

64 The quality-of-life and illness burden of stuttering, and its beginnings during childhood, seem to  
65 have prompted a consensus in the field that early intervention is an optimal management strategy. A  
66 report of 264 speech-language pathologists (SLPs) from 10 countries indicated that they gave  
67 waiting list priority to children who stutter; they prioritized childhood stuttering above all other  
68 developmental speech and language disorders (McGill et al., 2021). Eighteen Australian SLPs  
69 conveyed a similar perspective (Erickson et al., 2022), indicating that stuttering is “more  
70 debilitating than other communication disorders with the potential for long-term consequences to be  
71 more significant for clients” (p. 5). A consensus statement from participants at a conference with  
72 researchers and clinicians from 29 countries established that “current evidence ... clearly tells us the  
73 risks of early stuttering are certain and that they can be serious and potentially lifelong” (Lowe et  
74 al., 2021, p. 9).

75 **Neuroplasticity and Early Stuttering Intervention**

76 A large body of brain imaging research shows one causal factor for stuttering to be anomalies in  
77 the networking of brain regions involved with spoken language (Neef & Chang, 2024). Hence, the  
78 potential for neuroplastic change to produce lasting clinical benefits has been recognized (Chang et  
79 al., 2025). From a neuroscience perspective, it seems that there is a window of clinical opportunity  
80 during early childhood for stuttering to resolve. That neuroscience perspective is supported by  
81 randomized trials of early stuttering intervention (Sjøstrand et al., 2019). Although that evidence  
82 has limitations, it documents a large effect size from early stuttering intervention (with the  
83 Lidcombe Program) in relation to no-treatment controls. That effect size is associated with  
84 improvements in quality of life, attitude to communication, and psychological health (De  
85 Sonnevile-Koedoot et al., 2015; Woods; 2002). The clinical window during early stuttering begins  
86 to close as speech-related brain networking becomes clinically intractable with advancing age  
87 during childhood (Chang 2014; Neef & Chang, 2024; Shenker et al., 2023; Venkatagiri, 2005).  
88 Neuroplastic change during early intervention is a commonly accepted explanation for its reported

89 success (Chang, 2014). Indeed, in the case of the Lidcombe Program, it seems that effectiveness  
90 decreases markedly during late childhood (Johnson et al., 2024). Hence, the international consensus  
91 about the need for early intervention is well justified in terms of a limited opportunity to prevent  
92 lifetime quality-of-life and illness burden.

93 **Barriers to Intervention Immediately After Onset**

94 *Participation of children in the treatment process*

95 At present, various stages of the treatment processes of early interventions that have been studied  
96 with randomized trials—the Westmead Program (Andrews et al., 2023), the Lidcombe Program  
97 (Onslow et al., 2025), and RESTART-DCM (Franken et al., 2025)—require children to be involved  
98 and to participate in treatment activities. The Westmead Program requires children to speak in  
99 syllable-timed speech. The Lidcombe Program treatment process requires children to attend to  
100 parent feedback about their speech. Likewise, for some of its treatment components, RESTART-  
101 DCM requires cognitive development and cooperation from children. For example, during Phase II  
102 of the treatment, procedures include Riley and Riley’s (1985) 14-step speech motor training  
103 program. Children also learn variations of speech restructuring: “relaxed versus tense stuttering,  
104 repetitions versus prolongations/blocks, single repetitions versus multiple, glide sounds versus  
105 multiple repetitions” (Franken et al., 2025, p. 31). The Franken et al. (2025) manual also describes  
106 procedures where children intentionally stutter. Other recommended early interventions that are not  
107 supported by randomized trials commonly incorporate variants of speech restructuring (Kelman &  
108 Nicholas, 2020; Waelkens, 2018; Yaruss et al., 2006).

109 The procedures described above require cognitive engagement from children, yet the majority of  
110 stuttering onset occurs at a time when such cognitive involvement is clinically unrealistic for many  
111 children. The mean age of stuttering onset has been established from two prospective cohort studies  
112 to be in the range of 25–30 months (Ambrose & Yairi, 2015; Yairi, 1983; Yairi & Ambrose, 1992),  
113 with a median of 31 months (Reilly et al., 2013). This means that more than half, and perhaps even

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

114 the majority, of stuttering onset will occur before 3 years of age. A substantial number of those  
115 onsets are as early as 18 months (Reilly et al., 2013). Considering the need for children to actively  
116 participate in current treatment, it is not surprising, then, that there has been no randomized trial of  
117 children younger than 3 years of age (Onslow & Lowe, 2019; Sjøstrand et al., 2019).

### 118 *The challenges of existing early interventions*

119 Ideally, an immediate intervention would be as simple as possible to maximize its global uptake.  
120 Yet, the Lidcombe Program and RESTART-DCM are challenging treatments to administer.  
121 Although the Lidcombe Program’s clinical principal is simple—parent verbal contingencies applied  
122 to stutter-free speech and stuttering—the application of those principles needs to be designed  
123 individually for every child and family. This presents many clinical challenges, and problem-  
124 solving is a routine part of the treatment process. One publication (Van Eerdenbrugh et al., 2018)  
125 detailed 124 clinical challenges that occur with the Lidcombe Program and outlined clinical  
126 responses to them. RESTART-DCM is based on the demands and capacities model of early  
127 stuttering (Packman et al., 2004). As such, there are more than 60 clinical procedures for the  
128 clinician to consider for use with children (Onslow & Lowe, 2019). These include not interrupting  
129 the children when talking, reducing the number of questions asked, speaking with a calm voice, not  
130 imposing linguistic demands, and having an unhurried lifestyle.

### 131 *Limited speech-language pathology resources*

132 Regardless of the need and the potential benefits of immediate intervention after stuttering onset,  
133 the international SLP workforce is not equipped to deal with a disorder that has a prevalence of 1–  
134 2%. This has been confirmed by international government reports in the United Kingdom and  
135 Australia (I CAN, & Royal College of Speech and Language Therapists, 2018; Senate Community  
136 Affairs Committee Secretariat, 2014). The extent of this problem worldwide can be derived from  
137 the American Speech-Language Hearing Association, the world’s largest professional SLP group,  
138 which indicates that, at the time of writing, there are around 2,000 SLPs in the United States

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

139 currently accepting referrals for fluency (<https://www.asha.org/profind/>; Select: “Find a SLP”;  
140 search term: “fluency”; 2025, September). Although that figure is likely to be an underestimate, it  
141 illustrates the extent of the problem, equating to around one SLP for every 1,700 people who stutter  
142 in that country, based on a population of 340 million. Those data are for a country with well-  
143 developed SLP services, but most children who stutter live in countries where the profession is  
144 comparatively under-resourced. For example, Caldera et al. (2023) reported that the United States,  
145 United Kingdom, and Australia have, respectively, 57.7, 35.9, and 23.7 SLPs per 10,000  
146 population, but the figures for India, Malaysia, and Sri Lanka are 0.17, 0.95, and 0.44. In short, no  
147 country appears to have adequate SLP resources to deal with the public health issue of treating very  
148 young children who stutter.

### 149 **Searching for a new early intervention**

150 In summary, immediate intervention after stuttering onset has potential to influence the global  
151 health issue of chronic stuttering. Yet, empirically supported interventions are not suitable for  
152 children shortly after onset, who are mostly younger than 3 years, because those interventions  
153 require involvement from the children and because they are clinically challenging. These problems  
154 are compounded by limited SLP resources for stuttering globally. The search for a solution to these  
155 problems may be found in treatments such as RESTART-DCM, which incorporate many indirect  
156 strategies, as suggested by the demands and capacities model. These strategies recur in modern  
157 publicly available sources of advice to parents from professional associations and health services  
158 (Whelan, 2019; Whittington Health NHS Trust, 2020). These include many of the 60-plus  
159 procedures in RESTART-DCM treatment, as described earlier. Most of those strategies have no  
160 empirical clinical support, and one of them— reducing the number of questions asked to children—  
161 has been empirically refuted (Garbarino & Bernstein Ratner, 2022). However, there is direct  
162 experimental laboratory evidence that two of them may reduce early stuttering, as outlined below.

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

### 163 *Speech rate reduction*

164 A review of five laboratory experiments where parents reduced their speech rate with children  
165 who stutter concluded that reductions in stuttering severity of around 50% were observed overall  
166 (Davidow et al., 2016). For example, three single-subject experiments with preschoolers confirmed  
167 this effect (LaSalle, 2015), as did a study of 27 children in three laboratory sessions (Sawyer et al.,  
168 2017).

### 169 *Increased interturn speaker latency (pausing between speaking turns)*

170 Data from experimental laboratory studies show that when parents increase their interturn  
171 speaker latencies it reduces childhood stuttering. Interturn speaker latency is when a child says  
172 something to a parent, the parent pauses briefly before responding (Livingston et al., 2000;  
173 Winslow & Guitar, 1994). One of these experimental studies with a preschooler involved increased  
174 interturn speaker latency for 15 sessions at the family dinner table over a 7-week period, with  
175 results indicating that the technique was responsible for a 40–50% reduction in stuttering (Winslow  
176 & Guitar, 1994).

### 177 **A two-factor treatment protocol**

178 Based on the data above for parent speech rate reduction and pausing between speaking turns,  
179 we have designed the treatment protocol described below. In designing the protocol, we sought  
180 input from an advisory group comprising a representative from Speech Pathology Australia and  
181 treatment end-users (SLPs and parents of children who stutter). That input involved a 2-hour  
182 informal meeting, providing user and clinician perspectives on the clinical viability of the treatment  
183 and perspectives about ways to enhance compliance. Subsequently, the treatment protocol was  
184 developed after extensive discussions among the authors. The research was approved by the  
185 University of Technology Sydney Human Research Ethics Committee ETH23-8378.

186 Before the treatment protocol was administered, parents received a standard SLP assessment for  
187 early stuttering. That assessment included exploring parent understanding of stuttering causality,

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

188 dealing with any misinformation, and providing a full explanation of current knowledge about the  
189 nature of early stuttering and its childhood onset. Care is taken to ensure that parents know they are  
190 not responsible for the child's stuttering. The methods of the treatment protocol are presented  
191 within the context of the demands and capacities model, with an explanation that two evidence-  
192 based strategies would be used. Because the treatment protocol is simple, short 20-minute video  
193 telehealth consultations are sufficient to administer it. The protocol does not require the child to be  
194 present during clinical consultations, nor does the treatment require any active involvement from  
195 the child. In fact, the child will usually not even be aware the treatment is occurring.

### 196 *Step 1*

197 The clinician uses role play with the parent to demonstrate reducing speech rate by around one  
198 third and pausing for around half a second after each child utterance before replying. Those speech  
199 changes are demonstrated regardless of the parent's usual speech rate and pausing when speaking  
200 with the child. The values of one third speech rate reduction and half a second pausing were  
201 determined during consultation with the advisory group (see above), guided by clinical judgement  
202 about their practical feasibility.

### 203 *Step 2*

204 The parent uses these speaking techniques with the child twice per day for 5–10 minutes each  
205 time, but only in the home environment. The parent records one of these interactions for around 5  
206 minutes using a digital audio recording device such as a mobile phone. During weekly video  
207 telehealth consultations, the clinician reviews a 5-minute recording of the parent implementing the  
208 techniques and discusses any issues the parent may have encountered with using the techniques.

### 209 *Step 3*

210 When the parent is comfortable using the speaking techniques, the parent uses them as much as  
211 is practicable in and outside the home environment. The parent keeps a diary of their use. During

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

212 the weekly consultations, the clinician reviews the diarized information and a 5-minute recording of  
213 the parent implementing the techniques outside the home environment.

214 An SLP used our trial protocol to teach the strategies to parents of three preschoolers who  
215 stuttered, who were younger than 3 years of age (one parent for each child). The three parents  
216 signed their consent after reading a participant information statement. None had consulted a SLP  
217 previously about their child's stuttering, and none had attempted any treatment. The parents  
218 successfully learned to use the simple techniques in a similar manner during a 12-week trial period.  
219 Consistent with the laboratory research reviewed above, and the theoretical soundness of the  
220 procedures, there was a suggestion of possible treatment effects, as shown in Figure 1. The data in  
221 the figure are daily parent-assigned severity ratings using scale where 0 = *no stuttering*, 1 =  
222 *extremely mild stuttering*, 10 = *extremely severe stuttering*. Child 1 was a girl (2 years 7 months)  
223 with moderate–severe stuttering, whose parents reported that she had been stuttering for a couple of  
224 weeks. Her observable stuttering involved mainly repetitions, with some blocks. Child 2 was a girl  
225 (2 years 11 months) with moderate stuttering who had reportedly been stuttering for 6 months. Her  
226 observable stuttering involved mainly repetitions, with some blocks. Child 3 was a girl (3 years 0  
227 months) with mild stuttering, who had reportedly been stuttering for 8–9 months. Her observable  
228 stuttering involved repetitions, prolongations, and blocks. The stuttering severity of Child 1 and  
229 Child 2 improved over the 12 weeks of treatment, with trendlines suggesting continued  
230 improvement toward severity rating scores of 0 (no stuttering) if the treatment continues. Possibly  
231 because of a floor effect from extremely mild stuttering, Child 3 showed only slight reduction of  
232 stuttering severity during the 12 weeks of treatment.

233 These suggestions of a treatment effect are theoretically consistent with a mechanism of action  
234 involving neuroplastic change in response to reduced parent speech rate and increased pausing (see  
235 *Neuroplasticity and Early Stuttering Intervention*, above). An alternative explanation would be that  
236 reduced parent speech rate and increased pausing invoke similar speech changes in children's  
237 speech, which in turn control their stuttering. On the face of it, this seems to be a plausible

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

238 explanation, particularly in the case of reduced speed rate, which has long been known to reduce  
239 stuttering severity (Andrews et al., 1982; Johnson et al., 1937). However, available data cast doubt  
240 on such an explanation, with the bulk of studies showing no clear relationship between the speech  
241 rate of parents and children who stutter (Cardman, & Ryan, 2007; Guitar & Marchinkoski, 2001;  
242 Ratner, 1992; Sawyer et al., 2017; Stephenson-Opsal, & Ratner, 1988; Yaruss & Conture, 1995;  
243 Zebrowski et al., 1996). In fact, two studies showed an inverse relationship (Dehqan et al., 2008;  
244 Meyers & Freeman, 1985), and only one study showed a positive relationship where children's  
245 speech rate decreased in response to parent speech rate decrease (LaSalle, 2015).

246

247 INSERT FIGURE 1 ABOUT HERE

248

249 We propose this protocol as a basis for Phase I clinical trialling, with potential for further  
250 development during Phase II–IV trial evaluation. The latter trials would establish how long a two-  
251 factor treatment would take to establish treatment benefits, and they would evaluate the merits of  
252 the procedure, comparing it to existing treatments and comparing it to no-treatment controls to take  
253 account of variables such as natural recovery and parent speech attention. For that program of  
254 research, outcome measures would include stuttering severity, social anxiety, quality of life, and  
255 attitude to communication.

### 256 **Conclusion: Potential benefits of this two-factor early stuttering intervention**

#### 257 *Potential automation*

258 A stuttering treatment comprising speech rate reduction and pausing between speaking turns has  
259 the advantage of simplicity and that active involvement of children is not required. An additional  
260 advantage to such a treatment is that its simplicity lends itself to website-based automation,  
261 alleviating any demand on under-resourced global SLP healthcare services. A state-of-the-art  
262 automated web version of the treatment could be constructed (Arachchi et al., 2023), with a clinical

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

263 interface of lifelike AI-generated avatar SLPs. Using role play, the AI-generated avatar clinicians  
264 would demonstrate how to slow the speaking rate and how to pause between speaking turns. We  
265 envisage that the avatar clinicians would engage with parents using natural language processing to  
266 provide guidance through the treatment. The web-based treatment would incorporate advanced  
267 speech signal processing techniques to capture key speech features such as power spectrum (Molau  
268 et al., 2001) and provide feedback for speech rate reduction with acoustic measures of syllables  
269 spoken per second and feedback for pausing with acoustic measures of intervocalic intervals in  
270 milliseconds. Fast Fourier transform and advanced filtering techniques would remove unwanted  
271 noise to ensure that analyses focus correctly on the relevant speech signals. Because children who  
272 stutter may have parents who stutter (Darmody et al., 2022), such feedback will need to  
273 accommodate parent stuttering. Although speech signal processing is not yet fully able to recognize  
274 stuttering moments, progress is being made toward that being a realistic prospect (Mujtaba &  
275 Mahapatra, 2028; Shen & Zhang, 2025; Yawatkar et al., 2025).

276 The avatar clinicians would ensure that parents have learned the speaking techniques and  
277 understood the acoustically analyzed outcome. This objective feedback would overcome the  
278 inherent limitation of clinician perception. During Step 3 of the protocol, when parents use the  
279 speaking techniques in everyday conversations with their child, they would log in to the treatment  
280 site once per week and upload a 5-minute audio recording of their conversation with the child. They  
281 would diarize information about their use of the techniques, and log daily severity ratings of their  
282 child's stuttering. The avatar clinicians would provide feedback, recommendations, and  
283 encouragement based on this information.

### 284 *Preventing therapist drift*

285 Therapist drift is when a clinician deviates from delivering a treatment exactly the way it was  
286 designed and is manualized (Waller, 2009). This can occur intentionally or unintentionally, and may  
287 involve altering or omitting core treatment components, applying techniques inconsistently, or  
288 adapting a treatment to a personal style or perceived client needs. Therapist drift may occur for

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

289 many reasons, such as clinical time constraints, fatigue, and burnout. During current best-practice  
290 early stuttering intervention with the Lidcombe Program, therapist drift has been shown to be a  
291 negative influence (Carr Swift et al., 2011). However, automation of our proposed treatment would  
292 make therapist drift impossible. AI-generated avatar SLPs would provide objective and consistent  
293 treatment to parents about their use of the speech techniques. Consequently, parents and children  
294 globally—eventually in many languages—would have constant access to standardized health care  
295 for early stuttering.

### 296 *Cultural transferability*

297 The simplicity of this treatment maximizes the likelihood of it being uniquely culturally  
298 transferable. This would be an advance on existing early interventions that have been developed in  
299 Western cultures. An example is the Lidcombe Program, which relies on parents praising children  
300 for not stuttering. As such, there are reports of this aspect of it not being suitable for other cultures  
301 where the idea of praising children openly can feel unnatural and inappropriate, such as in Malaysia  
302 and Kuwait (Al-Khaledi et al., 2017; Vong et al., 2010). Another issue is that many non-Western  
303 cultures are not open about disability, perceiving it as a source of shame that should remain private.  
304 However, all current early interventions involve direct attention to the child’s speech. For example,  
305 in the Lidcombe Program, parents ask the child to repeat some utterances that are stuttered, and in  
306 the Westmead Program, children are encouraged to use rhythmic speech. With RESTART-DCM  
307 treatment, “now and again, the parent will casually speak using the disfluencies/stutterings the child  
308 makes, with the aim of lessening the child’s sensitivity to this” (Franken et al., 2025, p. 24). These  
309 approaches are not universally culturally appropriate. In principle, the proposed two-factor  
310 treatment of parents reducing speech rate and pausing between speaking turns potentially bypasses  
311 these issues because it does not focus on children in any way, and they may not even know that it is  
312 occurring.

313 **Stepped care intervention immediately after onset**

314 There is a genetic link with stuttering. Based on available genetic modelling, family aggregation  
315 data, and a review of family pedigrees (Darmody et al., 2022), stuttering will occur with 60–70% of  
316 children who have a first- or second-degree relative who stutters. Such children in the community  
317 could be identified by family history and monitored for stuttering onset. As soon as stuttering onset  
318 occurs, immediate two-factor intervention could be administered. This approach could introduce a  
319 stepped-care model to early stuttering intervention. The stepped-care model provides the simplest  
320 health care that is effective. Clients escalate to more resource-intensive and more expensive models  
321 of health care only as needed (Bower & Gilbody, 2005). The two-factor treatment is ideal as a first  
322 intervention in a stepped-care approach, particularly for an automated version. It can be used  
323 immediately after stuttering onset because it requires no active involvement from the child, it  
324 requires no clinician, and it is cost-free. Parents and children would escalate to the more resource-  
325 intensive and costly health care with a real clinician only if stuttering persists. Should Phase I–IV  
326 clinical trials support our proposed two-factor early intervention, adoption of that model by SLP  
327 associations worldwide would be a significant and transformative change to practice and policy of  
328 stuttering treatment. Cost-free early stuttering intervention could be made available to families  
329 globally, regardless of their location or socioeconomic status, alleviating any demand on under-  
330 resourced global SLP healthcare services.

331

332

333

**Acknowledgements**

334 The authors would also like to acknowledge Damien Liu-Brennan for his scientific  
335 copyediting contribution to this publication.

336

337

**Data Availability Statement**

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

338       The datasets generated during and/or analyzed during the current study are available from the  
339       corresponding author on reasonable request.

340

**References**

341

Al-Khaledi, M., Lincoln, M., McCabe, P., & Alshatti, T. (2017) The Lidcombe Program: A series of case studies with Kuwaiti preschool children who stutter. *Speech, Language, and Hearing, 21*, 224–235. <https://doi.org/10.1080/2050571X.2017.1370523>

344

Ambrose, N. G., Yairi, E., Loucks, T. M., Seery, C. H., & Throneburg, R. (2015). Relation of motor, linguistic and temperament factors in epidemiologic subtypes of persistent and recovered stuttering: Initial findings. *Journal of Fluency Disorders, 45*, 12–26. <https://doi.org/10.1016/j.jfludis.2015.05.004>

348

Andrews, G., Howie, P., Dosza, M., & Guitar, B. J. (1982). Stuttering: Speech pattern characteristics under fluency-inducing conditions. *Journal of Speech and Hearing Research, 25*(2), 208–216. <https://doi.org/10.1044/jshr.2502.208>

351

Andrews, C., Trajkovski, N., O'Brian, S., & Onslow, M. (2023). *The Westmead Program treatment guide (Ver. 1.3)* Retrieved from <https://www.uts.edu.au/asrc/resources/westmead-Program>

353

Arachchi, B. B., Sanjeeva J. M. I. P., Wijesooriya H. M. A. H., Chandrasena M. C., Kishara J., Kelegama, T., & Rajapakse V. M. (2023, December). StutterAI: Virtual assistant for stutter detection and analysis through machine learning and natural language processing approaches. In *2023 5th International Conference on Advancements in Computing (ICAC)* (pp. 525-530). IEEE.

357

Berchiatti, M., Badenes-Ribera, L., Ferrer, A., Longobardi, C., & Gastaldi, F. G. M. (2020). School adjustment in children who stutter: The quality of the student-teacher relationship, peer relationships, and children's academic and behavioral competence. *Children and Youth Services Review, 116*(1), Article 105226. <http://dx.doi.org/10.1016/j.childyouth.2020.105226>

361

Bloodstein, O, Bernstein Ratner, N., & Brundage, S. B. (2021). *A handbook on stuttering* (7th ed.). Plural Publishing.

363

Bower, P., & Gilbody, S. (2005). Stepped care in psychological therapies: Access, effectiveness, and efficiency. *British Journal of Psychiatry, 186*, 11–17. <https://doi.org/10.1192/bjp.186.1.11>

364

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

- 365 Boyle, C. A., Decoufle, P., & Yeargin-Allsopp, M. (1994). Prevalence and health impact of  
366 developmental disabilities in US children. *Pediatrics*, *93*(3), 399–403.
- 367 Briley, P. M., O'Brien, K., & Ellis, C. (2019). Behavioral, emotional, and social well-being in children  
368 who stutter: Evidence from the National Health Interview Survey. *Journal of Developmental and*  
369 *Physical Disabilities*, *31*(1), 39–53. <https://doi.org/10.1007/S10882-018-9625-X>
- 370 Caldera, A. V., Wickremasinghe, R., Munasinghe, T. U., Perera, K. M. N., Muttiah, N., Tilakaratne,  
371 D., Peiris, M. K. R. R., Thamilchelvan, E., Sooriyaarachchi, C., Nasma, M. N. and Manamperige, R.  
372 M. (2023). Availability of rehabilitation services for communication disorders in Sri Lanka: a cross-  
373 sectional survey. *BMJ Open*, *13*(4), e071620. <https://doi.org/10.1136/bmjopen-2023-071620>
- 374 Cardman, S., & Ryan, B. P. (2007). Experimental analysis of the relationship between speaking rate and  
375 stuttering during mother–child conversation II. *Journal of Developmental and Physical*  
376 *Disabilities*, *19*(5), 457–469. <https://psycnet.apa.org/doi/10.1007/s10882-007-9063-7>
- 377 Carr Swift, M., O'Brian, S., Hewat, S., Onslow, M., Packman, A., & Menzies, R. (2011). Investigating  
378 parent delivery of the Lidcombe Program. *International Journal of Speech Language Pathology*,  
379 *13*(4), 308–316. <https://doi.org/10.3109/17549507.2011.550692>
- 380 Chang, S. E. (2014). Research updates in neuroimaging studies of children who stutter. *Seminars in*  
381 *Speech and Language*, *35*(2), 67–79. <https://doi.org/10.1055/s-0034-1382151>
- 382 Chang, S. E., Jackson, E. S., Santayana, G., Zavos, G., & Onslow, M. (2025). Contemporary clinical  
383 conversations about stuttering: What does brain imaging research mean to clinicians? *International*  
384 *Journal of Speech-Language Pathology*, *27*(2), 265–271.  
385 <https://doi.org/10.1080/17549507.2024.2327472>
- 386 Craig, A., & Tran, Y. (2014). Trait and social anxiety in adults with chronic stuttering: Conclusions  
387 following meta-analysis. *Journal of Fluency Disorders*, *40*, 35–43.  
388 <https://doi.org/10.1016/j.jfludis.2014.01.001>

389

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

- 390 Darmody, T., O'Brian, S., Rogers, K., Onslow, M., Jacobs, C., McEwen, A., Lowe, R., Packman, A., &  
391 Menzies, R. (2022). Stuttering, family history and counselling: A contemporary database. *Journal of*  
392 *Fluency Disorders*, 73, Article 105925. <https://doi.org/10.1016/j.jfludis.2022.105925>
- 393 Davidow, J. H., Zaroogian, L., & A Garcia-Barrera., M. A. (2016). Strategies for teachers to manage  
394 stuttering in the classroom: A call for research. *Language, Speech, and Hearing Services in Schools*,  
395 47(4), 283–296. [https://doi.org/10.1044/2016\\_lshss-15-0057](https://doi.org/10.1044/2016_lshss-15-0057)
- 396 Dehqan, A., Bakhtiar, M., & Panahi, M. (2008). Relationship between stuttering severity in children and  
397 their mother's speaking rate. *São Paulo Medical Journal*, 126(1), 29–  
398 33. <https://doi.org/10.1590/S1516-31802008000100006>
- 399 De Sonnevile-Koedoot, C., Stolk, E., Rietveld, T., & Franken, M. -C. (2015). Direct versus indirect  
400 treatment for preschool children who stutter: The RESTART randomized trial. *PLoS One*, 10, Article  
401 e0133758. <https://doi.org/10.1371/journal.pone.0133758>
- 402 Erickson, S., Bridgman, K., & Furlong, L. (2022). Australian speech-language pathologists' experiences  
403 and perceptions of working with children who stutter: A qualitative study. *Journal of Fluency*  
404 *Disorders*, 75, Article 105944. <https://doi.org/10.1016/j.jfludis.2022.105944>
- 405 Franken, M. -C., Laroës, E., van Ormondt, J., de Smit, F., & Stipdonk, L. (2025). *RESTART-DCM*  
406 *method* (Rev. ed). <https://restartedcm.nl/wp-content/uploads/2025/02/RestartDCM-Method-2025.pdf>
- 407 Garbarino, J., & Bernstein Ratner, N. (2022). What is the role of questioning in young children's  
408 fluency? *American Journal of Speech-Language Pathology*, 31(5), 2061–2077.  
409 [https://doi.org/10.1044/2022\\_AJSLP-21-00209](https://doi.org/10.1044/2022_AJSLP-21-00209)
- 410 Gerlach, H., Totty, E., Subramanian, A., & Zebrowski, P. (2018). Stuttering and labor market outcomes  
411 in the United States. *Journal of Speech, Language, and Hearing Research*, 61(7), 1649–1663.  
412 [https://doi.org/10.1044/2018\\_JSLHR-S-17-0353](https://doi.org/10.1044/2018_JSLHR-S-17-0353)
- 413 I CAN, & Royal College of Speech and Language Therapists. (2018). *Bercow: Ten Years on*. Retrieved  
414 from <http://www.bercow10yearson.com/>

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

- 415 Iverach, L., Jones, M., McLellan, L. F., Lyneham, H. J., Menzies, R. G., Onslow, M., & Rapee, R. M.  
416 (2016). Prevalence of anxiety disorders among children who stutter. *Journal of Fluency Disorders*,  
417 49, 13–28. <https://doi.org/10.1016/j.jfludis.2016.07.002>
- 418 Iverach, L., O'Brian, S., Jones, M., Block, S., Lincoln, M., Harrison, E., Hewat, S., Menzies, R. G., &  
419 Onslow, M. (2009). Prevalence of anxiety disorders among adults seeking speech therapy for  
420 stuttering. *Journal of Anxiety Disorders*, 23(7), 928–934.  
421 <https://doi.org/10.1016/j.janxdis.2009.06.003>
- 422 Jacobs, M., Gerlach-Houck, H., & Briley, P. (2025). Differential impacts of anticipated success on  
423 employment outcomes among adults who stutter. *American Journal of Speech-Language Pathology*,  
424 34(1), 246–260. [https://doi.org/10.1044/2024\\_AJSLP-24-00202](https://doi.org/10.1044/2024_AJSLP-24-00202)
- 425 Johnson, G., Onslow, M., Carey, B., Jones, M., & Kefalianos, E. (2024). Lidcombe Program telehealth  
426 treatment for children 6-12 years of age: A Phase II trial. *Journal of Fluency Disorders*, 80, Article  
427 106057. <https://doi.org/10.1016/j.jfludis.2024.106057>
- 428 Johnson, W., & Rosen, L. (1937). Studies in the psychology of stuttering: VII: Effect of certain changes  
429 in speech pattern upon frequency of stuttering. *Journal of Speech Disorders*, 2(2), 105–110.  
430 <https://doi.org/10.1044/jshd.0202.105>
- 431 Kelman, E., & Nicholas, A. (2020). *Palin parent-child interaction therapy for early childhood*  
432 *stammering*. Routledge.
- 433 Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E. (2005). Lifetime  
434 prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey  
435 Replication. *Archives of General Psychiatry*, 62(6), 593–602.  
436 <https://doi.org/10.1001/archpsyc.62.6.593>
- 437 LaSalle, L. R. (2015). Slow speech rate effects on stuttering pre- schoolers with disordered phonology.  
438 *Clinical Linguistics & Phonics*, 29(5), 354–377. <https://doi.org/10.3109/02699206.2014.1003970>
- 439

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

- 440 Livingston, L. A., Flowers, Y. E., Hodor, B. A., & Ryan, B. P. (2000). The experimental analysis of  
441 interruption during conversation for three children who stutter. *Journal of Developmental and*  
442 *Physical Disabilities, 12*, 235–266. <https://doi.org/10.1023/A:1009465828258>
- 443 Lowe, R., Jelčić Jakšić, S., Onslow, M., O'Brian, S., Vanryckeghem, M., Millard, S., Kelman, E., Block,  
444 S., Franken, M. -C., Van Eerdenbrugh, S., Menzies, R., Shenker, R., Byrd, C., Bosshardt, H. -G., del  
445 Gado, F., & Lim, V. (2021). Contemporary issues with stuttering: The Fourth Croatia Stuttering  
446 Symposium. *Journal of Fluency Disorders, 70*, Article 105844.  
447 <https://doi.org/10.1016/j.jfludis.2021.105844>
- 448 Guitar, B., & Marchinkoski, L. (2001). Influence of mothers' slower speech on their children's speech  
449 rate. *Journal of Speech, Language, and Hearing Research, 44*(4), 853–861.  
450 [https://doi.org/10.1044/1092-4388\(2001/067\)](https://doi.org/10.1044/1092-4388(2001/067))
- 451 McAllister, J. (2016). Behavioural, emotional and social development of children who stutter. *Journal*  
452 *of Fluency Disorders, 50*, 23–32. <https://doi.org/10.1016/j.jfludis.2016.09.003>
- 453 McAllister, J., Collier, J., & Shepstone, L. (2012). The impact of adolescent stuttering on educational  
454 and employment outcomes: Evidence from a birth cohort study. *Journal of Fluency Disorders, 37*(2),  
455 106–121. <https://doi.org/10.1016/j.jfludis.2012.01.002>
- 456 McGill, N., McLeod, S., Crowe, K., Wang, C., Hopf, S. C. (2021). Waiting lists and prioritization of  
457 children for services: Speech-language pathologists' perspectives. *Journal of Communication*  
458 *Disorders, 91*, Article 106099. <https://doi.org/10.1016/j.jcomdis.2021.106099>
- 459 Meyers, S. C., & Freeman, F. J. (1985). Mother and child speech rates as a variable in stuttering and  
460 disfluency. *Journal of Speech, Language, and Hearing Research, 28*(3), 436–444.  
461 <https://doi.org/10.1044/jshr.2803.444>
- 462 Molau, S., Pitz, M., Schluter, R., & Ney, H. (2001, May). Computing mel-frequency cepstral  
463 coefficients on the power spectrum. In *2001 IEEE international conference on acoustics, speech, and*  
464 *signal processing. Proceedings (cat. No. 01CH37221)* (Vol. 1, pp. 73-76). IEEE.

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

- 465 Mujtaba, D., & Mahapatra, N. (2025). Fine-tuning ASR for stuttered speech: Personalized vs.  
466 generalized approaches. *arXiv:2506.00853v1 [cs.SD]*. <https://doi.org/10.48550/arXiv.2506.00853>
- 467 Neef, N. E., & Chang, S. E. (2024). Knowns and unknowns about the neurobiology of stuttering. *PLoS*  
468 *Biology*, 22(2), Article e3002492. <https://doi.org/10.1371/journal.pbio.3002492>
- 469 Norman, A., Lowe, R., Onslow, M., O'Brian, S., Packman, A., Menzies, R., & Schroeder, L.  
470 (2023). Cost of illness and health-related quality of life for stuttering: Two systematic  
471 reviews. *Journal of Speech, Language, and Hearing Research*, 66(11), 4414–4431.  
472 [https://doi.org/10.1044/2023\\_JSLHR-23-00072](https://doi.org/10.1044/2023_JSLHR-23-00072)
- 473 Onslow, M., & Lowe, R. (2019). After the RESTART trial: Six guidelines for clinical trials of early  
474 stuttering intervention. *International Journal of Language and Communication Disorders*,  
475 54(4), 517–528. <https://doi.org/10.1111/1460-6984.12463>
- 476 Onslow, M., Webber, M., Harrison, E., Arnott, S., Bridgman, K., Carey, B., Sheedy, S., O'Brian, S.,  
477 MacMillan, V., Lloyd, W., Santayana, G., Hodes, G., Smith, K., & Hearne, A. (2025). *The Lidcombe*  
478 *Program Treatment Guide* (Version 1.7c, August 2025). Lidcombe Program Trainers  
479 Consortium. [https://www.uts.edu.au/research/centres/australian-stuttering-research-](https://www.uts.edu.au/research/centres/australian-stuttering-research-centre/resources/lidcombe-program)  
480 [centre/resources/lidcombe-program](https://www.uts.edu.au/research/centres/australian-stuttering-research-centre/resources/lidcombe-program)
- 481 Packman, A., & Onslow, M., & Attanasio, J. (2004). The Demands and Capacities Model: Implications  
482 for evidence-based practice in the treatment of early stuttering. In A. K. Bothe (Ed.), *Evidence-based*  
483 *treatment of stuttering: Empirical issues and clinical implications* (pp. 65–79). Lawrence Erlbaum  
484 Associates.
- 485 Packman, A., Onslow, M., Richard, F., & Van Doorn, J (1996). Syllabic stress and variability: A model  
486 of stuttering. *Clinical Linguistics and Phonetics*, 10(3), 235–263.  
487 <https://doi.org/10.3109/02699209608985174>

488

489

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

- 490 Ratner, N. B. (1992). Measurable outcomes of instructions to modify normal parent-child verbal  
491 interactions: Implications for indirect stuttering therapy. *Journal of Speech, Language, and Hearing*  
492 *Research, 35*(1), 14–20. <https://doi.org/10.1044/jshr.3501.14>
- 493 Reilly, S., Onslow, M., Packman, A., Cini, E., Conway, L., Ukoumunne, O. C., Bavin, E. L., Prior, M.,  
494 Eadie, P., Block, S., & Wake, M. (2013). Natural history of stuttering to 4 years of age: A prospective  
495 community-based study. *Pediatrics, 132*(3), 460–467. <https://doi.org/10.1542/peds.2012-3067>
- 496 Riley, G., & Riley, J. (1985). *Oral motor assessment and treatment: Improving syllable production*.  
497 Pro-Ed.
- 498 Ryan, B. P. (2000). Speaking rate, conversational speech acts, interruption, and linguistic complexity of  
499 20 pre-school stuttering and non-stuttering children and their mothers. *Clinical Linguistics and*  
500 *Phonetics, 14*(1), 25–51. <https://doi.org/10.1080/026992000298931>
- 501 Sawyer, J., Matteson, C., Ou, H., & Nagase, T. (2017). The effects of parent-focused slow relaxed  
502 speech intervention on articulation rate, response time latency, and fluency in preschool children who  
503 stutter. *Journal of Speech, Language, & Hearing Research, 60*(4), 794–809.  
504 [https://doi.org/10.1044/2016\\_JSLHR-S-16-0002](https://doi.org/10.1044/2016_JSLHR-S-16-0002)
- 505 Senate Community Affairs Committee Secretariat (2014, September). Commonwealth of Australia.  
506 [https://www.aph.gov.au/-](https://www.aph.gov.au/-/media/Committees/Senate/committee/clac_ctte/speech_pathology/report/report.pdf)  
507 [/media/Committees/Senate/committee/clac\\_ctte/speech\\_pathology/report/report.pdf](https://www.aph.gov.au/-/media/Committees/Senate/committee/clac_ctte/speech_pathology/report/report.pdf)
- 508 Shen, J., & Zhang, X. (2025). Individual-independent and cross-language detection of speech  
509 disfluencies in stuttering based on multi-adversarial tasks and self-training. *Biomedical Signal*  
510 *Processing and Control, 100*(C), Article 107051. <https://doi.org/10.1016/j.bspc.2024.107051>
- 511 Shenker, R., Rodgers, N., Guitar, B., & Onslow, M. (2023). Contemporary clinical conversations about  
512 stuttering: Neurodiversity and ableism. *Journal of Fluency Disorders, 78*, Article  
513 106014. <https://doi.org/10.1016/j.jfludis.2023.106014>
- 514

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

- 515 Sjøstrand, Å., Kefalianos, E., Hofslundsengen, H., Guttormsen, L. S., Kirmess, M., Lervåg, A., &  
516 Bottegaard Naess, K. -A. (2019). Non-pharmacological interventions for stuttering in children six  
517 years and younger. *Cochrane Database of Systematic Reviews*, 9, Article  
518 CD013489. <https://doi.org/10.1002/14651858.CD013489.pub2>
- 519 Stein, M. B., Baird, A., & Walker, J. R. (1996). Social phobia in adults with stuttering. *American*  
520 *Journal of Psychiatry*, 153(2), 278–280. <https://doi.org/10.1176/ajp.153.2.278>
- 521 Stephenson-Opsal, D., & Ratner, N. B. (1988). Maternal speech rate modification and childhood  
522 stuttering. *Journal of Fluency Disorders*, 13(1), 49–56. [https://doi.org/10.1016/0094-730X\(88\)90027-](https://doi.org/10.1016/0094-730X(88)90027-7)  
523 7
- 524 Tığrak, T. K., Kayıkcı, M. E. K., Kirazlı, M. Ç., & Tığrak, A. (2020). Emotional and behavioural  
525 problems of children and adolescents who stutter: Comparison with typically developing  
526 peers. *Logopedics Phoniatrics Vocology*, 46(4), 186–192.  
527 <https://doi.org/10.1080/14015439.2020.1855472>
- 528 Van Eerdenbrugh, S., Packman, A., Onslow, M., O'Brian, S. (2018). Challenges and strategies for  
529 speech-language pathologists using the Lidcombe Program for early stuttering. *American Journal of*  
530 *Speech-Language Pathology*, 27(3S), 1259–1272. [https://doi.org/10.1044/2018\\_AJSLP-ODC11-17-](https://doi.org/10.1044/2018_AJSLP-ODC11-17-0185)  
531 0185
- 532 Venkatagiri, H. S. (2005). Recent advances in the treatment of stuttering: A theoretical perspective.  
533 *Journal of Communication Disorders*, 38(5), 375–393. <https://doi.org/10.1016/j.jcomdis.2005.03.001>
- 534 Vong, E., Wilson, L., McAllister, L., & Lincoln, M. (2010). Malaysian parents' use of positive  
535 reinforcement: A descriptive study and implications for the Lidcombe Program. *Asia Pacific Journal*  
536 *of Speech, Language and Hearing*, 13(2), 87–108. <https://doi.org/10.1179/136132810805335119>
- 537 Waelkens, V. (2018). *Mini-KIDS—Direct therapy for young children who stutter*. Veerle Waelkens.
- 538 Waller, G. (2009). Evidence-based treatment and therapist drift. *Behaviour Research and Therapy*,  
539 47(2), 119–127. <https://doi.org/10.1016/j.brat.2008.10.018>

## A PROPOSED TWO-FACTOR EARLY INTERVENTION

- 540 Whelan, A. (2019, May 19). 5 tips to share with parents of preschoolers who stutter. *The ASHA*  
541 *LeaderLive*. Retrieved from [https://leader.pubs.asha.org/doi/10.1044/5-tips-to-share-with-parents-of-](https://leader.pubs.asha.org/doi/10.1044/5-tips-to-share-with-parents-of-preschoolers-who-stutter/full/)  
542 [preschoolers-who-stutter/full/](https://leader.pubs.asha.org/doi/10.1044/5-tips-to-share-with-parents-of-preschoolers-who-stutter/full/)
- 543 Whittington Health NHS Trust (2020). *Tips for parents*. Retrieved from  
544 <https://www.whittington.nhs.uk/default.asp?c=29567>
- 545 Williams, D. E., Melrose, B. M., & Woods, C. L. (1969). The relationship between stuttering and  
546 academic achievement in children. *Journal of Communication Disorders*, 2(2), 87–98.  
547 [https://doi.org/10.1016/0021-9924\(69\)90032-X](https://doi.org/10.1016/0021-9924(69)90032-X)
- 548 Winslow, M., & Guitar, B. (1994). The effects of structured turn-taking on disfluencies: A case study.  
549 *Language, Speech, and Hearing Services in Schools*, 25(4), 251–257. [https://doi.org/10.1044/0161-](https://doi.org/10.1044/0161-1461.2504.251)  
550 [1461.2504.251](https://doi.org/10.1044/0161-1461.2504.251)
- 551 Woods, S., Shearsby, J., Onslow, M., & Burnham, D. (2002). Psychological impact of the Lidcombe  
552 Program of early stuttering intervention. *International Journal of Language and Communication*  
553 *Disorders*, 37(1), 31–40. <https://doi.org/10.1080/13682820110096670>
- 554 Yairi, E. (1983). The onset of stuttering in two- and three-year-old children: A preliminary report.  
555 *Journal of Speech and Hearing Disorders*, 48(2), 171–177. <https://doi.org/10.1044/jshd.4802.171>
- 556 Yairi, E., & Ambrose, N. (1992). Onset of stuttering in preschool children: Selected factors. *Journal of*  
557 *Speech and Hearing Research*, 35(4), 782–788. <https://doi.org/10.1044/jshr.3504.782>
- 558 Yaruss, J. S., Coleman, C., & Hammer, D. (2006). Treating preschool children who stutter: Description  
559 and preliminary evaluation of a family-focused treatment approach. *Language, Speech, and Hearing*  
560 *Services in Schools*, 37(2), 118–136. [https://doi.org/10.1044/0161-1461\(2006/014\)](https://doi.org/10.1044/0161-1461(2006/014))
- 561 Yaruss, J. S., & Conture, E. G. (1995). Mother and child speaking rates and utterance lengths in  
562 adjacent fluent utterances: Preliminary observations. *Journal of Fluency Disorders*, 20(3), 257–278.  
563 [https://doi.org/10.1016/0094-730X\(94\)00013-J](https://doi.org/10.1016/0094-730X(94)00013-J)
- 564

## A PROPOSED TWO-FACTOR EARLY INTERVENTION


565 Yawatkar, V., Chow, H. M., & Usler, E. (2025). Automatic temporal analysis of speech: A quick and  
566 objective pipeline for the assessment of overt stuttering. *Behavior Research Methods*, 57(8), Article  
567 228. <https://doi.org/10.3758/s13428-025-02733-z>

568 Zebrowski, P. M., Weiss, A. L., Savelkoul, E. M., & Hammer, C. S. (1996). The effect of maternal rate  
569 reduction on the stuttering, speech rates and linguistic productions of children who stutter: Evidence  
570 from individual dyads. *Clinical Linguistics and Phonetics*, 10(3), 189–206.  
571 <https://doi.org/10.3109/02699209608985171>

A PROPOSED TWO-FACTOR EARLY INTERVENTION

572 Captions

573 Figure 1: Daily parent-assigned severity ratings over 12 weeks for the three children in our trial  
574 protocol. The figure includes a trendline for the severity ratings.



575