

Altered motor control, posture and the Pilates
method of exercise prescription

Dorothy Curnow

Master of Arts (Performance) University of Western Sydney

A thesis submitted as partial requirement
for the degree of
Master of Science by Research

Faculty of Science

University of Technology Sydney

February 2010

CERTIFICATE OF AUTHORSHIP/ORIGINALITY

I certify that the work in this thesis has not previously been submitted for a degree nor has it been submitted as part of requirements for a degree except as fully acknowledged within the text.

I also certify that the thesis has been written by me. Any help that I have received in my research work and the preparation of the thesis itself has been acknowledged. In addition, I certify that all information sources and literature used are indicated in the thesis.

Signature of Student

Production Note:

Signature removed prior to publication.

Abstract

Aims

- To determine whether a basic set of Pilates exercises improves the efficiency of load transfer through the pelvis
- To compare the effects on chronic, mild low back pain (LBP) symptoms of three slightly different Pilates based regimes

Methods

A between subjects equivalent group experimental design was used

- Independent variable: type of exercise training (three groups)
- Dependent variables: efficient load transfer through the pelvis as measured by the stork test in weight bearing; low back pain symptoms

At entry, to establish baseline values, subjects completed an Oswestry Disability Questionnaire and recorded the frequency, intensity and duration of their back pain in an average week. Also, a Stork test was recorded.

Thirty-nine volunteers with mild chronic low back pain (CLBP) were taught four Pilates based exercises before being randomly allocated to one of three groups for the addition of other interventions.

- Groups A received four basic exercises
- Groups B and C received an additional relaxation posture using a specific spinal support
- Group C received an additional postural training exercise

Exercises were performed three times per week for six weeks and recording was done once per week for eight weeks. For each of the eight weeks, subjects recorded frequency, intensity and duration of back pain

At the final assessment, subjects completed another Oswestry Disability Questionnaire, returned their recording sheets, had a stork test recorded and their exercises checked.

Results

Oswestry Disability Questionnaire

Pre and post comparison of answers showed only one statistically significant improvement among subject groups. This was for question one, ‘*Do you have back pain at present?*’ where Group B reported significantly less pain post program compared with pre program (Wilcoxon, $z=-2.496$, $p=0.013$).

Number of days of pain

Group B experienced a statistically significant reduction in the number of days of pain between Week 1 and Weeks 6 to 8 ($F_{7,84}=6.4$, $p=0.0001$). Post hoc analysis using Scheffé showed significant differences between Week 1 and Weeks 6, 7 and 8 and between Week 2 and Week 8 ($p<0.05$). There were statistically significant differences by week within Group C ($F_{7,77}=3.29$, $p=0.0041$), but they only show up with Fisher ($p<0.05$) and they were between Week 1 and Weeks 6, 7 and 8, Week 2 and Weeks 6, 7 and 8, Week 3 and Weeks 6 and 7 and between Week 4 and Week 7.

Some of the improvements were lost once exercising ceased at the end of week 6

Duration of back pain episodes

All groups experienced a reduction in the mean length of the shortest, longest and average pain episodes. At week eight all groups had subjects who were pain free (Group A: 7.7%, Group B: 30.8%, Group C: 25%) and in Group B, no subjects reported pain episodes longer than six hours. However, differences were not statistically significant for the duration of this study.

Some of the improvements were lost once exercising ceased at the end of week 6

Intensity of pain across all lengths of pain episodes

While, all groups experienced a reduction in the intensity of pain across all lengths of pain episodes, the only statistically significant inter group difference involved Group A and Group B on the shortest pain episodes at Week 6, with Group B experiencing a greater reduction in pain intensity than Group A (Kruskal Wallis, $p=0.02$)

And for all groups, intensity of pain tended to rise once exercising ceased

Stork test in weight bearing

These results were inconclusive and this could have been because total subject numbers were small. As the same person carried out all tests, it was unlikely to reflect changes in measurement method or interpretation. However, it did raise questions regarding the stance for commencement of the test and this led to an investigation to assess the importance of the stance to the test results.

Procedure

A physiotherapist, experienced in taking Stork tests, who was not aware of the purpose of the test, carried out the stork tests. Ten subjects stood in three different standing positions to commence the test and neither subjects nor standing positions were tested consecutively.

- Not one subject had the same result recorded from all three positions
- There was no pattern to the results observed
- Two subjects had different results recorded from all three positions

Observations

- For correct execution of the exercises:
 - Subjects required follow up supervision
 - Those who had more individual attention learnt faster than those who had less
 - Understanding the intent of the exercise was more important than other factors such as age
- During periods of nonpractice correct execution was quickly forgotten

- Good acceptance of the program - 95% of the subjects decided to continue
 - However, compliance became an issue – the fewer the LBP symptoms, the less likely the subjects were to comply

Conclusions

Current pain symptoms (Oswestry Disability Questionnaire) showed a reduction in pain and all groups experienced statistically significant reductions in the frequency, duration and intensity of pain across the weeks of exercising. However, effects were not statistically significant between the groups, except for one instance.

Results indicate that other factors such as postural training and relaxation may impact on exercise programs and as the psoas muscle was the intended target for relaxation and training, it may indicate its importance in exercise rehabilitation.

Consistency of practice, supervision and follow up are important for correct exercise execution and once exercising ceases, reductions in LBP symptoms tend to diminish.

Stork test results and subsequent examination suggested that further investigation of the testing process was necessary and a further study was designed.

Aim of the stork test study

- To examine whether changing the stance of the starting position for the stork test on the weight-bearing leg altered the test outcome.

Methods

Seventy-four healthy, mobile adult volunteers took part. The exclusion criteria were stroke, spinal fusion, significant scoliosis, antalgic gait, hip or knee replacement or significant leg length difference. Subjects could suffer from minor ailments, including low back pain.

It was a within subjects experimental design with the independent variables being the four different stances for the commencement of the test and the dependent variable being the stork test which was measured using thumb locations on the posterior superior iliac spine and the second sacral spinous process.

All 296 tests were videoed and subsequently reviewed on a computer screen using a horizontal grid to identify the vertical direction and relative distance of thumb movement.

Results

Consistency between stance pairs was at best fair, with Kappa values ranging from $\kappa = -0.03$ to $\kappa = 0.34$. In conclusion, the starting stance influenced the outcome of the stork test with more than 64% of subjects failing to have consistent results across all four stances.

Discussion

For a comparable stork test measurement pre and post treatment, the same starting stance should be used. In addition the study showed that:

- The thumb on the innominate could rise, lower or remain at the same level
- The thumb on the sacrum could rise, lower or remain at the same level
- These movements could be independent

Thus, it is important to take note of the relative movement between the innominate (os coxa or pelvic bone) and sacral thumb positions since these can move independently during the test.

Acknowledgements

This research was made possible because of the contribution of many individuals and the University of Technology Sydney. I thank the following for their time and generous contribution at various stages of the research:

The subjects who volunteered for this time consuming exercise project

S T Boris Choy for his statistical advice and contribution

Narelle Smith for her statistical advice

Pat McLeod for assisting with data entry

Trish Wisbey-Roth for her encouragement

Jennifer Hoult for assisting with the initial stance and stork tests

Danielle Garrett for being the model for the photographs for the Journal of Bodywork and Movement Therapies

Jane Bramley for the preparation of the photographs

The subjects who volunteered for the stork tests

Pat McLeod, Laurinda Bailey and Nigel Lewis for their help with the setting up and filming of the stance and stork test project

Bruce Weaver for statistical advice and assistance with post hoc analysis of Chi-squared

Narelle Smith for statistical advice and assistance with the stork test data

Michelle Little and Julia Ting for their help, advice and constant support

The University of Technology Sydney and the Department of Medical and Molecular Biosciences for providing space for the project to take place

I have a special thank you for my supervisors Deirdre Cobbin and Jennifer Wyndham. Very individual in their approaches, they have been unfailing in their exceedingly generous contribution of expertise and time. I am very appreciative of their interest and assistance.

Finally, I would like to thank my husband, Nigel Lewis, for his support, encouragement and enthusiasm and my mother Ethel Curnow.

Contents

Abstract	i
Acknowledgements	vi
Contents	vii
Figures	x
Tables	xi
Chapter I: Introduction and aims	1
Background to the study	1
Aims	5
Format of thesis	6
Chapter II: Literature review	7
Introduction	7
Problems with exercise based studies	7
Identification of sub-sets in CLBP	10
Variable control	11
Studies involving direction specific exercise for CLBP	12
Studies involving Pilates for CLBP	14
Chapter III: Methods	17
Study design	17
Stork test	17
Advertising for volunteers	17
Initial screening	18

Exercises	19
Recording	20
Final assessment	20
Initial stork test collection	21
Statistical tests	22
Chapter IV: Results	23
Age and gender	23
Oswestry Disability Questionnaire	23
Number of days of pain	24
Duration of back pain episodes	25
Intensity of pain across all lengths of pain episodes	26
Stork test in weight bearing	28
Chapter V: Discussion	29
Observations	29
Oswestry Disability Questionnaire	30
Completion of program	31
Exercises	31
Measurements	37
Conclusions	38
Problems	39
Future directions	39
Chapter VI: Stork test on the weight bearing leg applied to four different stances for the commencement of the test	40
Introduction	40
Methods	41
Results	45
Discussion	52

Appendices

- Appendix I: Letter to subjects
- Appendix II: Health history
- Appendix III: Initial data
- Appendix IV: Oswestry Disability Questionnaire
- Appendix V: Initial assessment
- Appendix VI: Consent form
- Appendix VII: Exercise instructions
- Appendix VII: Postural exercise instructions
- Appendix IX: Frequently asked questions
- Appendix X: Subjects' recording sheets
- Appendix XI: Final assessment
- Appendix XII: Extra question for Oswestry Disability Questionnaire
- Appendix XIII: Protocols for subjects

Publications to date

Curnow, D. Cobbin D. Wyndham J. Choy STB. Altered motor control, posture and the Pilates method of exercise prescription. Proceedings of the 6th Interdisciplinary World Congress on Low Back & Pelvic Pain Barcelona, Spain 2007

Curnow, D. Cobbin D. Wyndham J. Choy STB. Altered motor control, posture and the Pilates method of exercise prescription. Proceedings of the Australian Pilates Method Association National Conference Melbourne, Australia 2008

Curnow, D. Cobbin D. Wyndham J. Choy STB. Altered motor control, posture and the Pilates method of exercise prescription. Journal of the Bodywork and Movement Therapies 2009; 13(1)

Figures

Figure 3.1: Subject flow through the study	21
Figure 4.1: Oswestry Disability Questionnaire	23
Figure 4.2: Number of days of pain each week	24
Figure 4.3: Duration of pain on the average length of pain episodes	25
Figure 4.4: Intensity of pain on the average length of pain episodes	26
Figure 4.5: Intensity of pain on the short length of pain episodes	27
Figure 5.1: Exercise 1 – the abdominal curl	32
Figure 5.2: Exercise 4 – the oblique abdominal curl	33
Figure 5.3: Exercise 2 – the side lying double leg lift	33
Figure 5.4: Exercise 3 – the spine extension	34
Figure 5.5: Postural training	35
Figure 6.1: Consistent stork test results across the stances	51
Figure 6.2: Positive stork test results across the stances	53

Tables

Table 3.1: Age and gender distribution for subjects who completed the final assessment	20
Table 6.1: Age range and gender breakdown of subjects	42
Table 6.2: Consistency of results across the stances for cephalad movement of the innominate thumb	47
Table 6.3: Consistency of results across the stances for cephalad movement of the innominate thumb without cephalad of the caudal thumb	48
Table 6.4: Consistency of results across the stances for caudal movement of the sacral thumb without caudal movement of the innominate thumb	50