

**THE BIOLOGY, ECOLOGY AND CONSERVATION  
OF WHITE'S SEAHORSE *HIPPOCAMPUS WHITEI***



**by**

**DAVID HARASTI**

B.Sci. (Hons), University of Canberra – 1997

This thesis is submitted for the degree of Doctor of Philosophy  
School of the Environment, University of Technology, Sydney, Australia.

July 2014

## CERTIFICATE OF ORIGINAL AUTHORSHIP

*I certify that the work in this thesis has not previously been submitted for a degree nor has it been submitted as part of requirements for a degree except as fully acknowledged within the text.*

*I also certify that the thesis has been written by me. Any help that I have received in my research work and the preparation of the thesis itself has been acknowledged. In addition, I certify that all information sources and literature used are indicated in the thesis.*

Signature of Student:



Dhovast

---

## ABSTRACT

Seahorses are iconic charismatic species that are threatened in many countries around the world with several species listed on the IUCN Red List as vulnerable or endangered. Populations of seahorses have declined through over-exploitation for traditional medicines, the aquarium trade and for curios and through loss of essential habitats. To conserve seahorse populations in the wild, they are listed on Appendix II of CITES, which controls trade by ensuring exporting countries must be able to certify that export of seahorses is not causing a decline or damage to wild populations. Within Australia, seahorses are protected in several states and also in Commonwealth waters.

The focus of this study was White's seahorse *Hippocampus whitei*, a medium-sized seahorse that is found occurring along the New South Wales (NSW) coast in Australia. The species is listed as 'data deficient' on the IUCN Red List and there is little research information available to assist in the conservation of the species. Research on *H. whitei* was undertaken from 2006-2009 and primarily focused on determining the species' life history parameters, its distribution and relative abundance, habitat preferences and site fidelity, and response to marine protected area (MPA) protection and habitat modification. Research primarily occurred within Sydney Harbour and Port Stephens.

Field surveys found that *H. whitei* is endemic to coastal estuaries along 300 km of NSW coastline. It is a medium-sized seahorse (max  $L_T$  162 mm) that displays rapid

growth (Port Stephens: females  $L_{\infty} = 149.2$  mm and  $K = 2.03$  per year and males  $L_{\infty} = 147.9$  mm and  $K = 2.52$ ; Sydney Harbour: females  $L_{\infty} = 139.8$  mm and  $K = 1.28$  and males  $L_{\infty} = 141.6$  mm and  $K = 1.22$ ), becomes sexually mature at approximately 6 months, and can live for up to 5 years in the wild. The species displays life-long monogamy with several pairs observed breeding over three consecutive breeding seasons, and strong site fidelity with seahorses remaining at the same site for up to 56 months for males and 49 months for females. Adult male and female *H. whitei* exhibited a significant preference for sponge and soft coral *Dendronephthya australis* habitats whilst juveniles had a strong preference for gorgonian *Euplexaura* sp. habitat. *Hippocampus whitei* in Port Stephens were significantly less abundant within the no-take MPA and there was a negative correlation with predator abundance. Long-term monitoring of *H. whitei* in Port Stephens found that populations declined over a period of six months for no apparent reason; however, they recovered within three years. A manipulative experiment undertaken on protective swimming nets in Sydney Harbour found *H. whitei* had a positive association with epibiotic growth and proximity to the sea floor. An experiment on the effects of flash photography found it had no significant effect on movements, behaviour, or site persistence of *H. whitei* and concluded that flash photography by divers is a safe and viable survey technique for this species. The information obtained from this study should contribute towards a reassessment of the species under the IUCN Red List and also provides the necessary data to ensure adequate management of the species within NSW waters.

## PREFACE

The contents within this thesis constitute my own work with several of the chapters having previously been published or are currently under review. As this thesis is based on chapters made up of published papers, there will be some repetitiveness across each of the chapters. References and formatting are based on the *Journal of Fish Biology* format. The development and implementation of each of the research projects within this thesis were of my own doing; however, I have had assistance from co-authors for various components of this study which I have outlined below.

Chapter 3 has been published as “Harasti, D., Martin-Smith, K., and Gladstone, W. (2013) Population dynamics and life history of a geographically restricted seahorse, *Hippocampus whitei*. *Journal of Fish Biology* **81**, 1297-1314.” All the diving work for this study was undertaken by myself and my two co-authors, my PhD supervisors, provided advice on the study design and guidance in the preparation of the manuscript.

Chapter 4 has been submitted to *Journal of Fish Biology* and is currently being considered for publication. Both co-authors provided advice and guidance in writing up this research. Professor Gladstone provided advice on the study design to determine habitat preferences whilst Dr Martin-Smith provided advice on monitoring seahorse movements using a towed GPS system.

Chapter 5 is in process of being submitted to a journal for consideration for publishing. All of the monthly monitoring for this four year study was undertaken by myself with both my co-authors (Martin-Smith and Gladstone) contributing to the study with assistance with the study design and guidance on the writing of the manuscript.

Chapter 6 has been published as “Harasti, D., Glasby, T., and Martin-Smith, K. (2010) Striking a balance between retaining populations of protected seahorses and maintenance of swimming nets. *Journal of Aquatic Conservation: Marine and Freshwater Ecosystems* **20**, 159-166. Dr Glasby provided advice in the development of the study design and statistical analysis for the manipulative experiment. Dr Martin-Smith provided the data that he had collected previously on seahorses at Clifton Gardens that was incorporated into this manuscript whilst I was responsible for the establishment of the experimental study and subsequent diving surveys.

Chapter 7 was a collaborative study with Beth Moore and Dr Hamilton from the California Academy of Sciences and was supported with funds from the Sea Life Conservation Fund (Sydney Aquarium). The project development, tissue sample collection and final write up were undertaken by myself. Beth Moore was responsible for the genetic analysis and provided advice on the written description of the methods and results. Dr Hamilton provided advice on the study design and provided comments on the chapter.

Chapter 7 has been published as “Harasti, D and Gladstone, W. (2013). Does underwater flash photography affect the behaviour, movement and site persistence of seahorses?” *Journal of Fish Biology* **83**, 1344-1353. Professor Gladstone assisted with this project by providing guidance on statistical analysis and preparation of manuscript. Dr Keith Martin-Smith also provided feedback on a draft manuscript that helped improve the publication.

## ACKNOWLEDGEMENTS

*I dedicate this thesis to my father, Dennis Harasti, who passed away unexpectedly in 2005, aged 52, and my grandfather, Taffy Williams, who passed away in 2013, aged 80.*

This PhD project arose from a meeting with Dr Keith Martin-Smith whilst he was working on seahorses at Clifton Gardens in Sydney Harbour. I'm very grateful that Keith took me on board as a student and he has provided excellent guidance, friendship and encouragement since this research commenced.

Professor William (Bill) Gladstone has provided excellent support as my university supervisor, first through the University of Newcastle and now at University of Technology Sydney. Bill has always provided great feedback on the drafts that I have sent and I'm very grateful of his supervision which has helped improve my research and made me a better scientist.

Thanks to Dr Tim Glasby who provided assistance with statistical analysis and filled a great role of 'unofficial supervisor'. Thanks for always responding to my stats questions at such short notice and helping me work through some of the more complicated analyses. You have made me think a lot more about the importance of planning the statistical analysis prior to rushing off and commencing any new study in the future.

I'm grateful to Pam and Chris Norman, former owners of Pro Dive Nelson Bay, who provided great support throughout the duration of the diving surveys and hundreds

of air-fills. Thanks to the Sea Life Conservation Fund (Sydney Aquarium) who provided some funding assistance for the research undertaken on the Manly net and the genetics study. My colleague and friend Chris Gallen provided great assistance with helping develop the GIS maps for the various chapters.

I'm very grateful of the support that my family has provided, especially my Mum who has been forever strong since Dad passed away. Finally, this thesis would not have been possible without the amazing support from my partner Suzanne. Outside of the fact that her excel wizardry has been of great benefit to me, especially since she taught me pivot tables, she has performed wonders in raising our two little boys (William and Ben) over the past 4 years. Thank you for being so understanding, loving and supportive, especially over the past 6 months whilst I was finishing the write up, and ensuring that I had enough 'quiet' time to finally get this thesis completed.

## TABLE OF CONTENTS

CERTIFICATE OF ORIGINAL AUTHORSHIP.....	ii
ABSTRACT .....	iii
PREFACE .....	v
ACKNOWLEDGEMENTS.....	viii
TABLE OF CONTENTS .....	x
LIST of TABLES.....	1
LIST of FIGURES.....	2
CHAPTER 1 .....	6
Justification and outline of research .....	6
CHAPTER 2 .....	9
Background .....	9
2.1 Outline .....	9
2.2 Seahorses .....	9
2.3 Conservation of seahorses.....	16
2.4 <i>Hippocampus whitei</i> – current knowledge.....	22
2.4.1 Description.....	22
2.4.2 Distribution.....	23
2.4.3 Reproductive biology.....	24
2.4.4 Habitat use .....	25
2.4.5 Home range .....	26
2.5 Conclusion.....	27
CHAPTER 3 .....	28
Population dynamics and life history of a geographically restricted seahorse, <i>Hippocampus whitei</i> .....	28
3.1 Abstract.....	29
3.2 Introduction .....	30
3.3 Materials and Methods.....	33
3.4 Results.....	42
3.5 Discussion.....	52
3.6 Acknowledgements .....	59
CHAPTER 4 .....	60

Ontogenic and sex-based differences in habitat preferences and site fidelity of the White's seahorse <i>Hippocampus whitei</i> .....	60
4.1 Abstract.....	61
4.2 Introduction .....	62
4.3 Materials and Methods.....	66
4.4 Results.....	76
4.5 Discussion.....	82
4.6 Acknowledgements .....	87
CHAPTER 5 .....	88
Seahorse abundance is reduced within a no-take Marine Protected Area. ....	88
5.1 Abstract.....	89
5.2 Introduction .....	90
5.3 Materials and Methods.....	93
5.4 Results.....	99
5.5 Discussion.....	105
5.6 Acknowledgements .....	112
CHAPTER 6 .....	113
Striking a balance between retaining populations of protected seahorses and maintenance of swimming nets.....	113
6.1 Abstract.....	114
6.2 Introduction .....	115
6.3 Materials and Methods.....	117
6.4 Results.....	125
6.5 Discussion.....	134
6.6 Acknowledgements .....	139
CHAPTER 7 .....	140
Are populations of the White's seahorse <i>Hippocampus whitei</i> genetically distinct across localities in New South Wales? .....	140
7.1 Abstract.....	141
7.2 Introduction .....	143
7.3 Materials and Methods.....	146

7.4	Results.....	152
7.5	Discussion.....	157
7.6	Acknowledgements .....	160
CHAPTER 8 .....		161
	Does underwater flash photography affect the behaviour, movement and site persistence of seahorses? .....	161
8.1	Abstract.....	162
8.2	Introduction .....	163
8.3	Materials and Methods.....	166
8.4	Results.....	172
8.5	Discussion.....	176
8.6	Acknowledgements .....	178
CHAPTER 9 .....		179
	General Discussion and Conclusion.....	179
9.1	Introduction .....	179
9.2	Study Results .....	180
9.2.1	Life History Parameters.....	180
9.2.2	Habitat usage and preferences.....	181
9.2.3	Population abundance .....	182
9.2.4	Genetics .....	183
9.2.5	Impacts of flash photography .....	183
9.3	Conservation implications .....	184
9.4	Future research .....	186
9.5	Conclusion.....	188
REFERENCES.....		190

## LIST OF TABLES

Table 3.1. GPS coordinates of survey locations in Sydney Harbour and Port Stephens.....	35
Table 3.2. Growth parameters ( $L_{\infty}$ and $K$ from the von Bertalanffy growth model) and natural mortality estimates for <i>Hippocampus whitei</i> from two different regions. $n$ = the total number of growth increments used in model with some individuals having been re-measured on more than one occasion.....	46
Table 4.1. Descriptions of habitat types from habitat preference study and observed habitat depth range.....	72
Table 5.1. Summary of hypotheses tested in the long term monitoring of <i>H. whitei</i> and predator abundance within and outside a marine protected area (MPA) with details of statistical analysis performed and PERMANOVA results. ....	102
Table 6.1. Summary of hypotheses tested in the manipulative cleaning experiment on the Manly net with details of statistical analysis performed and Analysis of Variance (ANOVA) results.....	128
Table 7.1. Locations of tissue sample collection for <i>Hippocampus whitei</i> in New South Wales.....	147
Table 7.2. Specimen information and collection localities for <i>H. whitei</i> specimens used in the pilot study. "X" is used to mark sequences that were used for the pilot study, while "X*" marks sequences that were only used initially to determine the variability at each locus. Variable sites are listed in numbers of nucleotide substitutions for CO1 sequences. Alignment locations are listed as the base numbers where substitutions were found in the CO1 alignment. ....	154
Table 7.3. CO1 pair-wise distance matrices, with distances (uncorrected $p$ ) listed as percentages. A) Distances between only the variable samples of <i>H. whitei</i> found in the pilot study. The highest observed distance is highlighted in yellow. B) Distances of closely related seahorse species, using CO1 sequences obtained from GenBank. The lowest distance between two species is shown in bold with *. ....	156

## LIST OF FIGURES

Figure 2.1. Morphology and measuring of seahorses (modified from Lourie *et al.* 1999b).....24

Figure 3.1. Known distribution of *Hippocampus whitei* in New South Wales – Australia with insert boxes of Port Stephens and Sydney Harbour study sites.....34

Figure 3.2. Mean  $\pm$  S.E. population estimates for male and female *Hippocampus whitei* on Manly net swimming enclosure (May 2007 – February 2008).....43

Figure 3.3. Estimated von Bertalanffy growth curve for males and female *Hippocampus whitei* for Port Stephens and Sydney Harbour. ● = mean size at maturity for male and ■ = mean size at first reproduction in Port Stephens. ● = smallest observed mature male (with pouch) and ■ = small pregnant male in Sydney Harbour. ♂ = oldest male and ♀ = oldest female observed in Port Stephens. ▲ = Size at birth of *H. whitei* from (Lourie *et al.*, 1999). .....45

Figure 3.4. Long term monogamy in *Hippocampus whitei* over three breeding seasons. Data is shown for those pairs ( $n=9$ ) that were resighted paired over consecutive breeding seasons. 1= Sighting of individual. Shading shows when individuals were found together during a survey while hatched shading indicate one of the original paired seahorses from the 2008-2009 breeding season was seen to pair up in the following breeding season with another individual. ....51

Figure 4.1. Location of study sites, Port Stephens, New South Wales – Australia.....71

Figure 4.2. Availability (expressed as % occurrence in all transects in each site) of habitats at the three study sites in January 2009. Seahorse Gardens = grey bars, Pipeline = white bars and Little Beach = black bars. ....78

Figure 4.3a. Habitat preferences scores ( $\pm 95\%$  CIs) of adult *Hippocampus whitei* (males = dark bars, females = light bars) across 3 sites in Port Stephens, New South Wales, Australia based on 1146 individual *H. whitei* observations from Jan 2008 – Dec 2009. Preference scores with a lower confidence interval  $>1$  (indicated by dashed line) indicate a significant habitat preference. ....79

Figure 4.3b. Habitat preferences of juvenile <i>Hippocampus whitei</i> across 3 sites in Port Stephens, New South Wales, Australia based on 100 observations from Jan 2008 – Dec 2009.....	79
Figure 4.4. Habitat availability (%) for the various size classes of the soft coral <i>Dendronephthya australis</i> across all three sites in Port Stephens. Seahorse Gardens = grey bars, Pipeline = white bars and Little Beach = black bars.....	80
Figure 4.5. Standardised habitat preference scores for adult and juvenile <i>Hippocampus whitei</i> combined across all three sites in relation to height classes for <i>Dendronephthya australis</i> .....	80
Figure 5.1. Location of study sites, Port Stephens, New South Wales – Australia.....	96
Figure 5.2. Mean monthly abundance of <i>H. whitei</i> ( $\pm$ S.E.) at four sites within Port Stephens for 2006-2009.....	100
Figure 5.3. Monthly abundance of adult <i>H. whitei</i> recorded in 60 min dive surveys from 2006-2009 at the non-sanctuary site Seahorse Gardens, Port Stephens.....	101
Figure 5.4. Monthly mean abundance ( $\pm$ S.E.) for <i>H. whitei</i> and predators (fish and octopus) for each site from 2007-2009.....	101
Figure 5.5. Monthly mean abundance ( $\pm$ S.E.) in 2007-2009 of seahorse predators at two sites within the Fly Point Sanctuary Zone and at two sites outside the Sanctuary Zone.....	104
Figure 5.6. Relationship between monthly seahorse abundance and predator abundance from 2007 – 2009 at each site.....	104
Figure 6.1. Location of study sites in Sydney Harbour.....	118
Figure 6.2. Total abundance of seahorses on swimming enclosure net at Clifton Gardens, Sydney Harbour, over the period 2003-2008. ■ = <i>Hippocampus whitei</i> , ● = <i>Hippocampus abdominalis</i> .....	126
Figure 6.3. Total abundance of <i>Hippocampus whitei</i> on Manly swimming enclosure net over 12 month period (May 2007 – April 2008). Dashed line indicates when manipulative cleaning was undertaken. Breeding season is shown with double arrow from September to February.....	129

Figure 6.4. Mean percentage ( $\pm$ S.E.) of total abundance of seahorses (average of randomly selected 28 animals per treatment each time) across treatments following net cleaning (average of seven surveys November 07– February 08). Treatment descriptions are Control (A) = 100% epibiotic growth; 100% Clean (B) = all epibiotic growth removed; Top Clean (C) = epibiotic growth removed from surface to 1.2m from seafloor; Bottom Clean (D) = epibiotic growth removed from seafloor to a height of 1.2m from seafloor; Other = Buffer zones and other sections of net not included in experiment (100% epibiotic growth). Treatments not significantly different in post-hoc tests marked with same letter (a or b) above bars.....131

Figures 6.5a–d. Distribution of seahorses among treatments four months after allocation to different net cleaning treatments (initially  $n=28$  seahorses per treatment). (a) seahorses initially in Treatment A, (b) seahorses initially in Treatment B, (c) seahorses initially in Treatment C, (d) seahorses initially in Treatment D. Treatments same as those described in Figure 6.4. ....132

Figure 6.6. Mean ( $\pm$ S.E.) of maximum distance moved by animals from each treatment. Treatments same as those described in Figure 6.4. ....133

Figure 7.1. Geographic distribution of *Hippocampus whitei* in New South Wales and sampling locations. ....148

Figure 8.1. Frequency of occurrence of responses of *Hippocampus whitei* ( $n=10$  surveys combined) to three experimental treatments ( $n=8$  individuals per treatment): flash photography, flash photography + handling, and photography without flash and no handling (control). No response =  (white bars with black diagonal stripe), actively hid = , changed holdfast = , swam away =  (black bars). ....173

Figure 8.2. Mean ( $\pm$ S.E.) number of *H. whitei* across three treatments ( $n=8$  per treatment), responding (with all responses combined) for each weekly survey ( $n=10$ ) [light coloured bars ()]. Mean ( $\pm$ S.E.) distance moved by *H. whitei* across each treatment ( $n=6$ ) for each weekly survey ( $n=9$ ) [dark coloured bars ()]. ....174

Figure 8.3. Cumulative site persistence of *Hippocampus whitei* across all three treatments from January 2009 – December 2009. Flash = — (black line), Flash + handled = — (grey line) and Control (no flash) = — — (dashed grey line).....175