THE DEVELOPMENT OF LIPOSOME ENCAPSULATED CALCIUM PHOSPHATES FOR BONE REGENERATION

A thesis submitted in fulfilment of the requirements for the degree of Doctor of Philosophy

by

KANTHI LEWIS

B.Sc. Applied Chemistry (Hons)

2010



© Kanthi Lewis 2010

CERTIFICATE OF AUTHORSHIP/ORIGINALITY

I, Kanthi Lewis, certify that the work in this thesis has not previously been submitted for a degree nor has it been submitted as part of requirements for a degree except as fully acknowledged within the text.

I also certify that the thesis has been written by me. Any help that I have received in my research work and the preparation of the thesis itself has been acknowledged. In addition, I certify that all information sources and literature used are indicated in the thesis.

> Production Note: Signature removed prior to publication.

Signature

Acknowledgements

First and foremost I would like to thank Dad. You always taught me to see through everything I commit to, no matter how tough it gets. Thanks for all the encouragement, love, support and belief you always had in me. Knowing how proud you have always been, even now that you are not here to remind me, always gives me strength to persevere.

Without you this would not have been possible, and this thesis is dedicated to your memory.

I would like to thank my wonderful supervisors Prof Besim Ben-Nissan, Dr Stella Valenzuela and Dr Louise Evans who have been incredibly supportive, and given me guidance these last 4 years. Always generous with your time and energy, I have learnt more than I ever thought possible, and I could not have asked for better supervisors.

Prof Racquel LeGeros, taking me on and mentoring me through my year at NYU, the experience was invaluable, and it has been an absolute privilege to learn from you.

I would also like to thank all the UTS staff particularly Mr Mark Berhkahn, Mr Ric Wurher, and Dr Ronald Shimmon, for help with the X-ray, Imaging, Mapping, Organic Synthesis and NMR, and Ms Linda Foley, Ms Era Koirala, all your help with ordering and generally make life as a PhD student much easier.

Dr Kristina Warton, Mr Nicholas Archer and Ms Stephanie Dowdell for all the assistance with the PCR, for always being there to help when I got to my wits end.

Dr Murray Killingsworth from Electron Microscopy Laboratory, South Western Area Pathology Service, for processing, sectioning and TEM imaging the liposomes.

Dr Dindo Mijares, I appreciate all your time and effort in the lab and helping me adjust. Dr Yuexun Liu for your help with the PCR data analysis

Dr Catherine Kealley, you always listened and understood. Always a few steps ahead, you were a great source of advice, so often my shoulder to lean on, and a wonderful friend.

My fellow PhD students, Dr Julia Ting, and Dr Dakrong Pissuwan, who made the lab a much brighter place, and always willing to be a helping hand.

I would like to thank all my family and friends for all your love and support

ABSTRACT

Osteoporosis, a degenerative bone disorder, is one of the leading causes of morbidity in the elderly. Proper nutrition plays a role in the prevention and treatment of osteoporosis. Intake of calcium and Vitamin D are some of the most important nutritional factors, and supplementation remains the gold standard and first line of treatment for low bone mineral density and osteoporosis. Supplementation can prevent bone loss and reduce fracture risk.

This work set about to produce, characterise and encapsulate for direct delivery to the bone various micro and nano sized calcium based mineral compounds which may be beneficial to bone health, using the precipitation method and biomimetic processes.

Calcium phosphate mineral was produced and characterised, including hydroxyapatite (Hap), dicalcium phosphate dihydrate (DCPD), as well as multiphase and substituted calcium phosphates using biomimetic process. Standard simulated body Fluid (SBF) solution was modified, creating a high carbonate solution which better mimics the bone environment, and produces precipitates more similar to bone than traditional low carbonate SBF, as confirmed using Fourier Tansform Infrared Spectroscopy (FTIR) and X-Ray Diffraction (XRD).

The use of liposomes as a delivery vesicle for the calcium mineral was evaluated using FTIR, XRD, Electron Dispersive Spectroscopy, Mass Spectrometry Transmission and Scanning Electron Microscopy, and X-Ray Mapping. The calcium mineral from aqueous solutions and prepared HAp and DCPD was incorporated into the liposome.

Functional groups were synthesised based on a published structure used to target the bone marrow macrophage, and incorporation into liposomes was confirmed using Nuclear Magnetic Resonance Spectroscopy and FTIR. Preliminary cell culture studies showed no direct effect on osteoblast like Mg63 or Saos-2 cells or osteoclast resorption, measured by bone collagen release.

Macrophage response was explored using U937 cell line. Expression of TNF- α and IL-1, markers of inflammation, increased with liposome treatments compared to the negative control but decreased compared the positive control. The Mg63 cells given U937 supernatants showed liposomes increased OPG production, but this was regardless of mineralisation.

The calcium based mineral compounds were produced, characterised, successfully encapsulated using liposomes and functionalised to improve uptake at the bone site. This shows the potential to deliver calcium to the bone, however further work to inhibit inflammation and increase the calcium dose to elicit greater cell response is required before this approach can be developed as a treatment option.

Table of Contents

i
ii
iii
vii
X
xvii
xviii

Chapter 1 Introduction

1.1 Significance	2
1.2 Aims	2
1.3 Summary of thesis outline	3
1.4 Outline of the thesis	4

Chapter 2 Literature review

2.1 Bone	7
2.2 Bone Classification	7
Bone Microstructure	8
2.3 Bone Composition	9
Organic Matrix	10
Inorganic Phase	11
Bone Cells	13
Lipids	14
Interstitial fluid	14
2.4 Bone Decomposition	15
2.5 Osteoporosis pathophysiology	15
Current treatments for Osteoporosis	18
Model Systems for studying Osteoporosis pathogenesis	21
2.6 The importance of Lipids in bone health	25
2.7 Liposomes	27
2.4 Bone Cell Activity	32
Markers of bone activity	34
Macrophage bone activity	36
2.8 References	44

Chapter 3 Mineral Production and Characterisation

3.1 Introduction	51
3.2 Mineral Production	54
Production of Simulated body fluid and Precipitation	54
Production of mineral from alternate calcified solutions	60
Production of Calcium phosphate mineral	62

3.3 Mineral Characterisation Methods	65
Fourier transform Infrared Spectroscopy	65
X-Ray Diffraction	66
3.4 Mineral Characterisation	68
3.5 Summary of Chapter findings	77
3.6 References	78

Chapter 4 Liposome and functional group production and Characterisation

4.1 Introduction	80
4.2 Preparation of Liposomes	82
4.3 Methods of Characterisation	84
Transmission Electron Microscopy	84
Nuclear Magnetic Resonance Spectroscopy	86
4.4 Characterisation of Liposomes	88
4.5 Functional Group production method	94
4.6 Functional group characterisation	95
4.7 Production of functionalised liposomes	100
4.8 Characterisation of functionalised liposomes	102
4.9 Summary of chapter findings	103
4.10 References	104

Chapter 5 Liposome Mineral incorporation

5.1 Introduction	106
5.2 Methods of Characterisation	109
Scanning Electron Microspcopy	110
Xray Mapping	111
Inductively coupled plasma- Mass Spectrometry	112
5.3 Characterisation of Mineralised Liposomes	113
5.4 Characterisation of Sample Set 1	118
5.5 Characterisation of Sample Set 2	124
5.6 Characterisation of Sample Set 3	129
5.7 Summary of Chapter findings	139
5.8 References	140

Chapter 6	Osteoblast and osteoclast response	
6.1 Intro	duction	142
6.2 Prima	ary Cell Culture	144
B	one Collection	144
С	ell culture method for osteoblast isolation	145
6.3 Cell I	Lines	147
6.4 Meth	ods for determining Cytotoxicity and Proliferation	150
С	ytotoxicity and liposome dose determination	150
P	roliferation assay method	152
6.5 Meth	ods of cell differentiation determination	153
A	Ikaline Phosphatase Activity	153

154
155
157
162
166
167
169
171
173
176
177

179 180 181

Chapter	7 Macrophage cell culture
7.1	Introduction
7.2	U937 cell culture methods
	Proliferation
	Differentiation

Differentiation	184
Background to PCR	184
PCR method	186
PCR results	194
7.3 Mg63 response to U937 cytokines	196
Proliferation and differentiation	196
Alkaline Phosphatase activity	197
Osteoprotegerin production	198
7.4 Discussion of results	199
7.5 Summary of chapter findings	202
7.6 References	203

Chapter 8	Conclusions and Further Work	
8.1 Conclu	usions and further work	206
8.2 References		217
Appendix A	X-ray Diffraction reference patterns	219
Appendix B	Statistical analysis for cell culture	221

List of Abbreviations

Alkaline phosphatase (ALP) Back scattered electron (BSE) Bone mineral density (BMD) Bovine serum albumin (BSA) Calcium acetate $(Ca(Ac)_2)$ Colony stimulating factor-1 (CSF-1) Dicalcium phosphate dihydrate (DCPD) Environmnental Scanning Electron Microscope (ESEM) Enzyme linked immunosorbent assay (ELISA) Ethyleme diaminetetra acetic acid (EDTA) Extracellular matrix (ECM) Foetal Bovine Serum (FBS) Fourier Transform Infrared Spectroscopy (FTIR) Gaseous secondary electron (GSE) Hormone replacement therapy (HRT) Horseradish peroxidise (HRP). Hydroxyapatite (Hap) Inductively coupled plasma (ICP). Institutional Animal Care and Use Committee (IUCAC) Insulin like growth factor 1 (IGF-I) Interferon- γ (IFN γ), Interleukin (IL), Latent TGF-β binding protein (LTBP)-1 Macrophage colony stimulating factor (MCSF) Macrophage inflammatory protein-1 alpha (MIP-1a)

Mass spectrometry (MS) Monocyte chemoattractant protein-1 (MCP-1) Monoclonal antibodies (Mabs) Propidium Iodide (PI) Raloxifene (RAL) Receptor Activator for Nuclear Factor K B Ligand (RANK-L) Repetitions per minute (RPM) New York University (NYU) Nuclear factor-jB ligand /RANK ligand (RANKL) Nuclear magnetic Resonance Spectroscopy (NMR). Osteoprotegerin (OPG) Ovariectomized (OVX) Parathyroid hormone (PTH), Phosphate Buffered Saline (PBS) Phosphatidycholine (PC) Polymerase Chain Reaction (PCR) p-nitrophenol (pNp) p-nitrophenol phosphate (p-Npp) Reactive oxygen species (ROS) Ribonucleic Acid (RNA) Scanning electron microscope (SEM) Simulated body Fluid (SBF) Small integrin-binding ligand N-linked glycoprotein (SIBLING) family Tartrate-resistant acid phosphatase type 5b (TRAcP-5b) Tetramethylbenzidine (TMB) T helper 1 $(T_H 1)$ Transmission electron microscopy (TEM).

Transforming growth factor (TGF)- β

Tumour-necrosis factor (TNF)

United States Department of Agriculture (USDA).

University of Technology, Sydney (UTS)

Width of Field (WOF)

X-ray Diffraction (XRD)

X-Ray Mapping (XRM)

7-dehydrocholesterol (7- DHC)

1,25-dihydroxyvitaminD₃ (1,25(OH)₂D₃).

1,2 distearoyl-sn-glycero-3-phospho-ethanolamine-N-[monomethoxy poly(ethylene glycol) 5000 (PEG-DSPE)

1,5 Dipalmmitoyl-L-glutamate-N-succinic acid (LGSA)

List of Figures

Chapter 1

No Figures in Chapter 1

- Figure 2.1 Examples of the four bone types shown in the human skeleton.
- Figure 2.1 Microstructure of bone modified
- Figure 2.2 Unit cell structure of HAp
- Figure 2.4 Cell activity in bone, showing osteoblasts in bone formation and osteoclasts resorbing bone
- Figure 2.5 An imbalance between bone formation and resorption leads to trabecular thinning (left) and eventual loss of trabecular connectivity (right)
- Figure 2.6 Cationic lipids activating the cellular activity
- Figure 2.7 Structure of SA
- Figure 2.8 A schematic representation of different liposome types
- Figure 2.9 Liposomes with the four types of functionalisation.
- Figure.2.10 Schematic of active osteoclast and osteoclasts
- Figure 2.11 Bone resorbtion and formation
- Figure 2.12 Osteoclast differentiation pathway
- Figure 2.13 Cytokines affecting the osteoclast.
- Figure 2.14 Monocyte differentiation
- Figure 2.15 Cationic lipid effect on cell activity
- Figure 2.16 Activated T cells induce osteoclast genesis

- Figure 3.1 Outline of mineral production and characterisation.
- Figure 3.2 Synthesis setup for the preparation of hydroxyapatite
- Figure 3.3 Diffraction from crystal planes.
- Figure 3.4 FTIR spectrum of mineral produced from a) low carbonate SBF b) high carbonate SBF c) low carbonate SBF with Flouride
- Figure 3.5 FTIR spectrum of a) Hydroxyapatite b) Dicalcium phosphate dehydrate c) calcium carbonate
- Figure 3.6 X-ray diffraction pattern of mineral obtained from a) low carbonate SBF solution with JCPDS sodium chloride reference pattern b) high carbonate SBF solution with JCPDS sodium chloride reference pattern c) low carbonate SBF solution containing fluoride with JCPDS sodium chloride reference pattern
- Figure 3.7 X-ray diffraction pattern hydroxyapatite with JCPDS hydroxyapatite reference
- Figure 3.8 X-ray diffraction pattern of Hydroxyapatite with JCPDS calcium hydrogen phosphate hydrate reference pattern
- Figure 3.9 X-ray diffraction pattern of dicalcium phosphate dihydrate with JCPDS dicalcium phosphate dihydrate (Brushite) reference pattern
- Figure 3.10 X-ray diffraction pattern of calcium carbonate with JCPDS calcium carbonate reference pattern

- Figure 4.1 Outline of production and characterisation of liposomes
- Figure 4.2 Structure of Phosphatidycholine
- Figure 4.3 Mulitlammelar liposomes with hydrophobic tails grouping together, and polar heads in contact with aqueous solution.
- Figure 4.4 TEM image of unilamellar liposomes with mineralised portions indicated by green arrows.
- Figure 4.5 TEM image of Unilamellar liposomes with some multilamellar samples beginning to form and shown with red arrows.
- Figure 4.6 TEM image of a mix of multilamellar and unilamellar liposomes produced in a calcifying buffer unilamellar liposomes shown by green arrows in these images and multilamellar liposomes in red.
- Figure 4.7 TEM image of a mix of multilamellar and unilamellar liposomes produced in a calcifying buffer
- Figure 4.8 TEM image of a unilamellar liposome produced in a calcyfying buffer
- Figure 4.9 TEM image of multinlamellar liposomes in a calcyfying buffer
- Figure 4.10 TEM image of multilamellar lipsomes in a calcifying buffer.
- Figure 4.11 The structure of compound 1. l-glutamic acid, *N*-(3-carboxy-1-oxopropyl)-, 1,5dihexadecyl ester
- Figure 4.12 The structure of L-glutamine, the four Hydrogen environments that appear in the NMR spectrum are numbered
- Figure 4.13 The structure of hexadecyl alcohol, the four Hydrogen environments that appear in the NMR spectrum are numbered
- Figure 4.14 H-NMR spectrum of starting materials, a) L-glutamine b) hexadecyl alcohol
- Figure 4.15 H-NMR spectrum of compound 1- a) before purification b) after purification
- Figure 4.16 H-NMR spectrum of Compound 2- 1,5 dipalmmitoyl-L-glutamate-N-succinic acid
- Figure 4.17 FTIR spectrum of a) compound 1 after purification b) compound 2-1,5 Dipalmmitoyl-L-glutamate-N-succinic acid
- Figure 4.18 H-NMR spectrum of liposome a) without PEG b) with spontaneously incorporated PEG

- Figure 5.1 Summary of Liposome Characterisation
- Figure 5.2 TEM image of a multilamellar liposomes, produced in SBF
- Figure 5.3 Image of liposomes taken using ESEM in SE mode showing multilamellar liposomes.
- Figure 5.4 EDS spectra of the liposomes on the TEM grid
- Figure 5.5 Backscatter images of liposomes in SBF solution.
- Figure 5.6 EDS of liposomes plus SBF.
- Figure 5.7 Pseudo colour images sample set 1 a) liposomes in PBS Control b) Liposomes in PBS plus HAp c) Liposomes in PBS plus DCPD
- Figure 5.8 EDS sample set 1 a) liposomes in PBS Control b)Liposomes in PBS plus HAp c) Liposomes in PBS plus DCPD.
- Figure 5.9 scatter diagrams sample set 1 a) liposomes in PBS Control b) Liposomes in PBS plus HA c) Liposomes in PBS plus DCPD, showing the correlation between calcium and phosphate
- Figure 5.10 Elemental mapping diagram liposomes in PBS Control
- Figure 5.11 Elemental mapping diagram Liposomes in PBS plus HA
- Figure 5.12 Elemental mapping diagram Liposomes in PBS plus DCPD
- Figure 5.13 Representation of calcium ions binding to the phosphate layer of the liposome
- Figure 5.14 Pseudo colour images of sample set 1 a) liposomes in low CO₃ SBF b)
 Liposomes in low CO₃ SBF +F c) Liposomes in high CO₃ SBF, showing the distribution of calcium, phosphate and the background silicon.
- Figure 5.15 Scatter diagrams produced from X-ray maps for sample set 1 a) liposomes in low CO₃ SBF b) Liposomes in low CO₃ SBF +F c) Liposomes in high CO₃ SBF, showing the correlation between calcium and phosphate in the sample.
- Figure 5.16 Elemental mapping diagram, liposomes in low CO₃ showing the distribution of selected elements over the sample.
- Figure 5.17 Elemental mapping diagrams Liposomes in low CO₃ SBF +F showing the distribution of selected elements over the sample.
- Figure 5.18 Elemental mapping diagrams Liposomes in high CO₃ SBF, showing the distribution of selected elements over the sample.
- Figure 5.19 EDS images produced from the whole sample area for set 1 a) liposomes in low CO₃ SBF b) Liposomes in low CO₃ SBF +F c) Liposomes in high CO₃ SBF

- Figure 5.20 Pseudo colour images of a) liposomes produced with CaCl₂ b) liposomes produced with CaCl₂ plus NaHCO₃ c) liposomes produced with CaCl₂ plus K₂HPO₄ plus MgSO₄ solution.
- Figure 5.21 Scatter diagrams of a) liposomes produced with CaCl₂ b) liposomes produced with CaCl₂ plus NaHCO₃ c) liposomes produced with CaCl₂ plus K₂HPO₄ plus MgSO₄ solution.
- Figure 5.22 Images and EDS over the entire image of a) liposomes produced with CaCl₂ b) liposomes produced with CaCl₂ plus NaHCO₃ c) liposomes produced with CaCl₂ plus K₂HPO₄ plus MgSO₄ solution.
- Figure 5.23 Elemental mapping diagrams of liposomes produced with CaCl₂
- Figure 5.24 Elemental mapping diagrams of liposomes produced with CaCl₂ plus NaHCO₃
- Figure 5.25 Elemental mapping diagrams of liposomes produced with CaCl₂ plus K₂HPO₄ plus MgSO₄ solution.
- Figure 5.26 Mapping image with a scatter diagram showing the correlation between calcium and phosphorus, the region outlined in a black box, as the areas of high calcium and phosphorous are shown in yellow in the corresponding mapping image.
- Figure 5.27 ESEM image of the NaCO₃ plus CaCl₂ sample with EDS spectra obtained for the portion outlined with a red box.
- Figure 5.28 Pseudo colour image of the sample containing CaCl₂ KHPO₄ and MgSO₄ with Sulphur shown in blue, phosphorous in red and calcium in green
- Figure 5.29 ESEM image with EDS spectra obtained for the portion outlined with a red box.
- Figure 5.30 ESEM image, with scatter diagram and EDS spectra of the area selected in red.
- Figure 5.31 X-ray diffraction pattern for the solution with two brushite phase present, as shown by the JCPDS reference patterns [11], shown in purple and green.

- Figure 6.1 Summary of osteoblast and osteoclast cell culture studies
- Figure 6.2 Bone samples in complete bone media for cell culture using explants method.
- Figure 6.3 a) Cell death for Saos-2 cell line with varied liposome concentrations shows a significant increase in cell death for the 2mg/ml after day 3 b) cell death represented as a change from the control sample.
- Figure 6.4 a) Cell death for Mg63 cell line with varied liposome concentrations shows a significant increase in cell death for the 2mg/ml after day 3 b) cell death represented as a change from the control sample.
- Figure 6.5 a) Proliferation for Mg63 cell line with non functionalised liposome samples. Cell numbers are shown after 3,5 and 7 days of growth.
- Figure 6.6 Proliferation for Mg63 cell line with funtionalised liposome samples. Cell numbers are shown after 3,5 and 7 days of growth.
- Figure 6.7 Proliferation for Saos-2 cell line with functionalised liposome samples. Cell numbers are shown after 3,5 and 7 days of growth.
- Figure 6.8 Proliferation for Mg63 cell line with functionalised liposome samples. Cell numbers are shown after 3,5 and 7 days of growth.
- Figure 6.9 ALP activity for Mg63 cell line with non functionalised liposome samples.
- Figure 6.10 ALP activity for Mg63 cell line with functionalised liposome samples.
- Figure 6.11 ALP activity for Saos-2 cell line with non functionalised liposome samples.
- Figure 6.12 ALP activity for Saos-2 cell line with functionalised liposome samples.
- Figure 6.13 Mg63 Osteocalcin production after 10 days, with different liposome samples.
- Figure 6.14 Mg63 Osteoprotegerin production after 10 days, with different liposome samples.
- Figure 6.15 Tibia obtained from a female Sprague Dawley rat.
- Figure 6.16 Trap assay for osteoclast precursors treated with differentitation factors and, liposome samples, shows all samples positive for TRAP, indicating that differentation has occured.
- Figure 6.17 Osteoclasts, collagen release from bone in the presence of liposome samples.

- Figure 7.1 Summary of cell culture work in chapter 7
- Figure 7.2 a) Proliferation for U937 cell line with non functionalised liposome samples. Cell numbers are shown after 2, and 3, days of growth.
- Figure 7.3 a) Proliferation for U937 cell line with functionalised liposome samples. Cell numbers are shown after 2 and 3 days of growth.
- Figure 7.4 A) diagramatic representation of DNA, RNA and protein shown [8] with B) schematic DNA RNA and Protein and involvement in various processes [7].
- Figure 7.5 Phases of a PCR reaction [5]
- Figure 7.6 PCR program used for IL-1, TNF, BTF, and OPG
- Figure 7.7 Melting curve obtained for BTF-3 the house keeping gene, showing the melting point at 88 $^{\circ}\mathrm{C}$
- Figure 7.8 Melting curve obtained for TNF-α aplification, showing the melting point at 83°C
- Figure 7.9 Melting curve obtained for OPG amplification, showing the product melting point at 87 $^{\rm o}{\rm C}$
- Figure 7.10 Melting curve obtained for Il-1 amplification, showing the product melting point at 86 $^{\circ}\mathrm{C}$
- Figure 7.11 Melting curves of negative controls without template DNA. No melting points are present, as no products have formed.
- Figure 7.12 Electrophoresis gels run at 60V for 90mins, showing PCR product sizes corresponding to the expected values shown in table 7.1
- Figure 7.13 Proliferation of Mg63 cell line at 5 and 7 days, with U937 supernatants
- Figure 7.14 ALP activity changes in the Mg63 cell line with the U937 supernatants

List of Equations

Equation 2.1	Conversion of DCPI	Conversion of DCPD to HAp		
$CaHPO_4 + Ca_4(PO_4)_2O \not E Ca_5(PO_4)_3OH$				
Equation 3.1	Bicarbonate bufferin	Bicarbonate buffering system		
$CO_2(g) + H_2O \iff CO_2(aq) + H_2O$		(1) dissolved CO ₂		
$CO_2(aq) + H_2O \Leftrightarrow H_2CO_3$		(2) carbonic acid		
$H_2CO_3 \Leftrightarrow H^+ + HCO_3^-$		(3) bicarbonate	pKa ₁ =6.35 at 25°C	
$HCO_{3} \Leftrightarrow H^{+} + CO_{3}^{2}$		(4) carbonate	pKa ₂ =10.33at 25°C	
Equation 3.2	Frequency of molect	Frequency of molecular Vibrations		
	$U = \frac{1}{2\pi} \sqrt{\frac{k(m_1 + m_2)}{m_1 m_2}}$			
Equation 3.3 Relationship between wavelength, incident radiation and			nt radiation and	
diffraction				
	$n\lambda = 2dsin\theta$			
Equation 4.1 Calculation of Area of PEG dispersion				
	$APEG = \pi R_F^2$			
Equation 4.2	n 4.2 Definition of Florey radius			
	$RF = N^{3/5}a$			
Equation 4.3	Percentage of lipid s	Percentage of lipid surface covered with PEG		
$\mathbf{R} = \mathbf{A}_{PEG} \mathbf{x} \mathbf{M} / \mathbf{A}_{lipid}$				
Equation 5.1	Calculation of ALP	Calculation of ALP activity		
$IU = \mu mol/(L \cdot min)$				
	$= \frac{(ODsample_{\tau} - ODsample_{0}) \cdot 1000 \cdot \text{Re} actionVol}{t \cdot \epsilon \cdot l \cdot samplevol}$			
$= \frac{(ODsample_{1} - ODsample_{0}) \cdot \text{Re} actionVol}{(ODcalibrator - ODH_{2}O) \cdot SampleVol \cdot t}$				
$(ODcalibrator - ODH_2O) \cdot Sample Vol \cdot t$				
Equation 7.1. Change between gene of interest and housekeeping gene (HKG)				

Equation 7.1. Change between gene of interest and housekeeping gene (HKG)

Gene 1 Ct – HKG Ct = Δ Ct

Equation 7.2. Sample change compared to control sample

Sample $\Delta Ct - Control \Delta Ct = X$

Equation 7.3. Find fold increase, as each Cycle represents and exponential increase Fold increase (or decrease) = Power (X,2)

List of Tables

Table 1.2 Biologically Significant Calcium phosphate compounds

Table 2.1 Properties of the Osteoblast Phenotype

Table 3.1 Compounds used for preparation of 1L of Kukubos SBF

Table 3.2 Compounds used for preparation of 500ml of Kanthi's SBF

Table 3.3 Concentrations of Ions in SBF solutions compared to blood plasma

Table 3.4 pH readings of high carbonate SBF with storage at 37 $^{\circ}\mathrm{C}$ and 5% CO2

Table 3.5 Ion concentrations (mmol) in solutions used.

Table 3.6 Different processes available for the production of Hap

Table 3.7 Assignment of IR vibrational bands to functional groups

Table 3.8 Calcium phosphate compounds and major identifying x-ray diffraction peaks (2θ) and expected relative intensities (I) values taken from JCPDS {JCPDS, 1995 #311}

Table 5.1 Concentrations of ions compared to standard solutions

Table 6.1 Collagenase Digest composition

Table 6.2 Bone growth media

Table 6.3 Summary of Cell Culture results

Table 7.1 Primer sequences with resulting product size, and mRNA, NCBI ascension number.

Table 7.2 Change in fold expression of OPG in U937 with liposome treatments

Table 7.3 Change in fold expression of TNF in U937 with liposome treatments

Table 7.4 Change in fold expression of IL-1 in U937 with liposome treatments

Table 7.5 Change in fold expression of OPG in Mg63 with U937 supernatants