

# A genomic island integrated into *recA* of *Vibrio cholerae* contains a divergent *recA* and provides multi-pathway protection from DNA damage

Rita A. Rapa,<sup>1,2</sup> Atiqul Islam,<sup>1</sup> Leigh G. Monahan,<sup>1</sup>  
Ankur Mutreja,<sup>3</sup> Nicholas Thomson,<sup>3</sup> Ian G. Charles,<sup>1</sup>  
Harold W. Stokes<sup>1</sup> and Maurizio Labbate<sup>1,2\*</sup>

<sup>1</sup>*ithree Institute, <sup>2</sup>Department of Medical and Molecular Biosciences, University of Technology, PO Box 123 Broadway, Sydney, NSW 2007, Australia.*

<sup>3</sup>*Wellcome Trust Sanger Institute, Cambridge, UK.*

## Summary

Lateral gene transfer (LGT) has been crucial in the evolution of the cholera pathogen, *Vibrio cholerae*. The two major virulence factors are present on two different mobile genetic elements, a bacteriophage containing the cholera toxin genes and a genomic island (GI) containing the intestinal adhesin genes. Non-toxigenic *V. cholerae* in the aquatic environment are a major source of novel DNA that allows the pathogen to morph via LGT. In this study, we report a novel GI from a non-toxigenic *V. cholerae* strain containing multiple genes involved in DNA repair including the recombination repair gene *recA* that is 23% divergent from the indigenous *recA* and genes involved in the translesion synthesis pathway. This is the first report of a GI containing the critical gene *recA* and the first report of a GI that targets insertion into a specific site within *recA*. We show that possession of the island in *Escherichia coli* is protective against DNA damage induced by UV-irradiation and DNA targeting antibiotics. This study highlights the importance of genetic elements such as GIs in the evolution of *V. cholerae* and emphasizes the importance of environmental strains as a source of novel DNA that can influence the pathogenicity of toxigenic strains.

## Introduction

*Vibrio cholerae* is a common inhabitant of marine and estuarine waters and is the causative agent of the

diarrheal disease cholera. Although there are over 200 O-antigen serogroups among *V. cholerae* strains, only two, O1 and O139, are known to cause pandemics of cholera disease (Kaper *et al.*, 1995). Lateral gene transfer (LGT) has largely contributed to the emergence of new pandemic strains of cholera (Faruque and Mekalanos, 2003; Keymer and Boehm, 2011). The appearance of the O139 serogroup and the so-called hybrid strains in the early 1990s are prime examples (Ramamurthy *et al.*, 2003; Safa *et al.*, 2009). Mobile genetic elements (MGEs) have been pivotal in the evolution of *V. cholerae* including diverse elements, such as the genomic islands (GIs) VPI-1, VPI-2, VSP-1, VSP-2, an integrative conjugative element, and the bacteriophage CTX (Faruque and Mekalanos, 2003; Grim *et al.*, 2010). GIs are defined as large chromosomal regions that have features suggestive of recent LGT (Boyd *et al.*, 2008). They have the capacity to excise and form circular intermediates and often target tRNA loci for their integration. In *V. cholerae*, GIs have been implicated in causing human disease and in environmental survival. For example, the replacement of the O1 classical biotype by the O1 El Tor biotype in the 1960s is suggested to be due to the acquisition of VSP-1 and VSP-2 that have probably enhanced epidemic spread (Faruque and Mekalanos, 2003). VPI-2 is a 57.3 kb island integrated at *tRNA-Ser* and encodes a neuraminidase important for converting higher-order sialogangliosides to GM1 gangliosides, the receptor for cholera toxin (Galen *et al.*, 1992). Moreover, VPI-1 encodes for the toxin-coregulated pilus (TCP), an essential intestinal colonization factor, as well as the accessory colonization factor (ACF), and virulence regulators ToxT and TcpPH (Everiss *et al.*, 1994; Murphy and Boyd, 2008). Non-O1/O139 *V. cholerae* strains are considered to be the major source of laterally acquired DNA for O1/O139 strains (Meibom *et al.*, 2005) thus, a better understanding of the diverse genetic elements present in the *V. cholerae* species is important for predicting and mitigating the emergence of new pandemic strains.

In bacteria, errors in DNA can occur as part of normal DNA replication or can be induced by external stimuli (e.g. UV irradiation) (Janion, 2008). There are several

Received 31 January, 2014; accepted 13 May, 2014. \*For correspondence. E-mail maurizio.labbate@uts.edu.au; Tel. (61-2) 9514 4064; Fax (61-2) 9514 8206.

genetic systems involved in error-free DNA repair including base excision repair (BER), nucleotide excision repair (NER), recombinational DNA repair and mismatch repair (MMR) (Ratray and Strathern, 2003; Janion, 2008; Polosina and Cupples, 2010; Lenhart *et al.*, 2012). However, if DNA damage is extensive the mutagenic phase of the SOS response is triggered (Goodman, 2002). This response is mediated by DNA polymerases that replicate past template lesions in a process called translesion DNA synthesis (TLS) that is inherently error-prone (Goodman, 2002). For example, DNA polymerase V, encoded by the *umuDC* operon (Patel *et al.*, 2010). The SOS induction of error-prone polymerases is considered a final response where although induced mutation(s) may be deleterious to the host cell, this is balanced against the need for rapid DNA repair (Goodman, 2002). An alternative view for the function of error-prone polymerases is that they act to generate genetic diversity that may have a role in environments where the host is maladapted by providing a bank of pre-existing genetic diversity within that population, some of which may confer a positive selective advantage. To support this second view, transcription of error-prone polymerases has been observed in the absence of SOS inducing DNA damage (Yeiser *et al.*, 2002). Furthermore, error-prone polymerase mutants are less competitive than the parent cells during starvation (McKenzie *et al.*, 2000; Yeiser *et al.*, 2002; Tark *et al.*, 2005), and some antibiotics (e.g. quinolones) induce the SOS mutagenic response increasing the frequency of resistant mutants (Pidcock and Wise, 1987; Ysern *et al.*, 1990).

In this study we report a novel GI inserted into *recA* of *V. cholerae* non-O1/O139 strain S24 isolated from an estuarine river in Sydney, Australia. This strain lacks the major virulence factors: cholera toxin and the toxin-coregulated pilus, thus is not capable of causing cholera. The GI carries (i) a *recA* gene phylogenetically distant from the disrupted host *recA*, designated *recA*<sub>RME</sub>; (ii) a *umuDC* operon, designated *umuDC*<sub>RME</sub>, encoding DNA polymerase V; and (iii) genes encoding hypothetical proteins, proteins with DNA processing domains including a MutL domain involved in MMR, and proteins involved in site-specific recombination. The GI can excise as a closed circle and preferentially inserts into a specific site within *recA*. We also show that *recA*<sub>RME</sub> is functional and provides protection from UV irradiation, a common source of DNA damage encountered in the shallow waters of marine and estuarine environments. Furthermore, the GI provides protection from the antibiotics bleomycin and ciprofloxacin. Acquisition of this GI by O1/O139 toxigenic *V. cholerae* would not only enhance survival of this pathogen in the natural environment but may also provide enhanced protection from DNA targeting antibiotics such as ciprofloxacin.


## Results and discussion

### Identification of a novel genomic island in *V. cholerae* S24 containing *recA*

S24 is an environmental, non-O1/O139 *V. cholerae* strain isolated from Georges River in Sydney, Australia, as described in a previous study (Islam *et al.*, 2013). It was noted during multilocus sequence analysis of housekeeping genes *adk*, *gyrB*, *mdh* and *recA* using primers designed to amplify the *V. cholerae* S24 *recA*, (designated here *recA*<sub>S24</sub>), that a product of ~ 1.5 kb was identified instead of the expected ~ 850 bp. When the *V. cholerae* S24 draft genome sequence (to be released at a later date) was interrogated, it was noted that *recA*<sub>S24</sub> was present on two separate contigs. PCR, using primers designed to sequence within these contigs, was used to close this region of the genome (described in *Experimental procedures*) resulting in a final contig of 262,869 bp. Within this contig and disrupting *recA*<sub>S24</sub> at 494 bp into the 1065 bp gene was a GI of 32,787 bp we have designated *recA* mobile element (RME). Consistent with RME being a mobile genetic element, the GC content is 41.3% compared with the genome average of 47.2%, it encodes mobility functions (see below) and is bordered by 9 bp inverted repeats, designated IR<sub>R</sub> (for *recA* end) and IR<sub>I</sub> (for integrase end) (Fig. 1). Bioinformatic analysis of the GI identified 23 coding sequences (CDSs) (Fig. 1) including a complete copy of *recA*, designated *recA*<sub>RME</sub> at the IR<sub>R</sub> end and a phage integrase at the IR<sub>I</sub> end. To our knowledge, this is the first mobile genetic element associated with the lateral movement of the critical gene, *recA*.

A number of other genes similar to those known to be involved in DNA processing are also found in RME including *umuDC* encoding the error-prone DNA polymerase V and a gene encoding a protein with a partial domain found in MutL (COG0323; Fig. 1), a component of the MMR pathway (Polosina and Cupples, 2010). A number other genes on RME, homologous to those involved in DNA processes include those encoding a ParB-like nuclease (91% identity to *Vibrio alginolyticus* 12G01; WP\_005381205.1), a redox sensitive transcriptional activator with a SoxR-domain (96% identity to *Vibrio* sp. 712i1; WP\_017634100.1), a type II restriction enzyme containing a methylase subunit (77% identity to *Vibrio splendidus*; WP\_017082665.1) and a helicase (90% identity to *Vibrio brasiliensis* LMG 20546; WP\_006880978.1).

Two insertion sequence (ISVvu4) elements were identified at positions 12,877 – 14,083 and 15,897 – 17,103 (striped boxes in Fig. 1) of RME. In both instances, 7 bp direct repeats (DR) were evident bordering the ISVvu4 elements indicating insertion by transposition. The DR for each ISVvu4 element is different, indicating independent




Locus tag	start	stop	orientation	Putative protein identification
RME001	204	1271	+	Recombinase protein RecA
RME002	1373	2593	-	Par-B like nuclease
RME003	2783	3916	+	Hypothetical
RME004	6225	4204	-	MutL mismatch repair domain-containing protein
RME005	6547	9219	+	Type II restriction enzyme, methylase subunit
RME006	9247	11268	+	Hypothetical; DEAD/DEAH box helicase
RME007	11284	12531	+	Hypothetical; YeeC-like nuclease domain-containing protein
RME008	13036	14025	+	Transposase of ISVvu4
RME009	14731	14114	-	Hypothetical
RME010	15192	14734	-	Hypothetical
RME011	15609	15199	-	Hypothetical; DnaJ domain-containing protein
RME012	16056	17045	+	Transposase of ISVvu4
RME013	17645	17211	-	Redox-sensitive transcriptional activator, SoxR
RME014	17963	18844	+	Permease of the drug/metabolite transporter (DMT) superfamily
RME015	21105	19225	-	Hypothetical; chromosome segregation ATPase-containing protein
RME016	22080	21598	-	Hypothetical
RME017	23939	22683	-	Error-prone, lesion bypass DNA polymerase V (UmuC)
RME018	24199	23939	-	Error-prone, lesion bypass DNA polymerase V (UmuD)
RME019	24984	25739	+	Multi domain XerS site-specific tyrosine recombinase XerS
RME020	27461	26001	-	Hypothetical
RME021	28181	27465	-	Hypothetical
RME022	30198	28165	-	Phage integrase family domain protein
RME023	32131	30185	-	Hypothetical

**Fig. 1.** Genetic structure and gene content of the *recA* genomic island. The RME contains 9 bp inverted repeats at each end ( $IR_R$  and  $IR_i$ ) and 23 ORFs inclusive of the transposase genes from the ISVvu4 elements (striped boxes). The ISVvu4 elements are abutted by 7 bp direct repeats (DR) indicating insertion by transposition. RME contains multiple genes in DNA repair including a full copy of *recA* (RME001), the *umuDC* operon (RME017 and RME018) encoding the two subunits of DNA polymerase V and a gene encoding a protein with a MutL mismatch repair domain (RME004).

insertion events. *In silico* removal of the ISVvu4 elements from the sequence did not restore any CDSs indicating that their insertion had not led to gene disruption. As expected, the promoter regions of both *recA<sub>RME</sub>* and the *umuDC<sub>RME</sub>* operon have the characteristic LexA binding sequence of CTGT-(AT)<sub>4</sub>-ACAG indicating control by the SOS response (Wertman and Mount, 1985; Sanchez-Alberola *et al.*, 2012). Present on RME are also genes putatively involved in mobilization/integration such as a phage integrase (RME022) and a site-specific recombinase XerS (RME019) (Fig. 1).

Phylogenetic analysis of *recA* sequences from the *Vibrionaceae* determined that *recA<sub>S24</sub>* is characteristic

of *recA* genes found within the *V. cholerae* clade, whereas *recA<sub>RME</sub>* is not. It does, however, group with other more distantly related *recA* genes found in other members of the *Vibrio* genus (Fig. 2). This is consistent with *recA<sub>RME</sub>* having been acquired by LGT. *recA* is an excellent phylogenetic marker for resolving relationships within the *Vibrionaceae* family (Stine *et al.*, 2000; Thompson *et al.*, 2004). Although the acquisition of a divergent *recA* in *V. cholerae* S24 is easily evident, this data reminds us that LGT of critical housekeeping genes like *recA* can and does occur. Less evident would be LGT of *recA* between closely related strains within the *V. cholerae* species confounding phylogenetic trees



**Fig. 2.** Phylogenetic analysis of *recA*<sub>S24</sub> and *recA*<sub>RME</sub> (RME highlighted in red) *recA*<sub>S24</sub> (also highlighted in red) groups with *V. cholerae* strains whereas, *recA*<sub>RME</sub> groups with *recA* from other *Vibrio* species indicating that *recA*<sub>RME</sub> was mobilized from another member of the *Vibrio* genus.


using a single marker (Bapteste *et al.*, 2004; Creevey *et al.*, 2004).

#### *The recA mobile element excises as a closed circle and targets a specific site in recA*

Many GIs are known to excise from their location in the chromosome (Boyd *et al.*, 2008). Analysis of RME suggested that it integrated into *recA*<sub>S24</sub> using site-specific recombination. In site-specific recombination, a DNA recombinase recognizes specific sequences (usually inverted sequences) allowing for DNA breakage and joining reactions that result in integration or excision of the element (Hallet and Sherratt, 1997). Exact excision of RME at IR<sub>R</sub> and IR<sub>I</sub> was predicted to leave behind a 4 bp scar introducing a frame shift in *recA*<sub>S24</sub> (Fig. 3A). To determine whether RME excision would leave behind an excision scar, an inverse PCR was conducted using primers reading out from the IR<sub>R</sub> and IR<sub>I</sub> ends (primers RME-F/RME-R in Table 1). A product of ~ 560 bp was amplified (see gel image in Fig. 3B) and sequenced. Analysis of the sequence showed that excision occurred in one of two possible ways (Fig. 3B): (1) Excision occurred at 2 bp on either side of the IR<sub>R</sub> and IR<sub>I</sub> ends (black arrows in Fig. 3A) and/or (2) precise excision occurred at the end of IR<sub>I</sub> and at 4 bp before the end of

IR<sub>R</sub> (grey arrows in Fig. 3A). Either way, excision was predicted to restore an uninterrupted and therefore functional copy of *recA*<sub>S24</sub> in the chromosome, consistent with site-specific recombination. This was confirmed by amplification of an intact 'empty' insertion site using primers S24-cinA-F/S24-recX-R and excising the predicted ~ 1.6 kb fragment (marked with a diamond in Fig. 3C). A nested PCR was then performed on the purified excised fragment using primers EcoRI-*recA*-F/EcoRI-*recA*-R (see gel image in Fig. 3C) and the product sequenced (Fig. 3C).

To determine whether the RME was capable of translocating from the genome of *V. cholerae* S24 into a new location, a vector (pOriVn<sub>700</sub>-*recA*<sub>S22</sub>; see Fig. 4A) containing the *recA* gene from a closely related strain of *V. cholerae* S24, strain S22 (Islam *et al.*, 2013), was introduced into *V. cholerae* S24 by conjugation. Here, the RME is expected to excise from the genome of *V. cholerae* S24 and insert into *recA*<sub>S22</sub> present on pOriVn<sub>700</sub>-*recA*<sub>S22</sub>. A control vector substituting *recA*<sub>S22</sub> with *gfp* (pOriVn<sub>700</sub>-P<sub>lac</sub>*gfp*; see Fig. 4A) was also introduced into *V. cholerae* S24 by conjugation as a control. Primers (ori6k-R and ori700-F; see Fig. 4A) targeting the vector backbone and the ends of the RME were used in a PCR reaction to determine whether the RME had mobilized into either pOriVn<sub>700</sub>-*recA*<sub>S22</sub> or pOriVn<sub>700</sub>-P<sub>lac</sub>*gfp*. In



**Fig. 3.** A. Sequence abutting insertion of the *recA* genomic island in *V. cholerae* S24. The black and grey arrows demarcate the possible excision points for the RME. B. Sequence and gel image of the product derived from inverse PCR of the excised RME. The sequence shows that excision does not precisely occur at  $IR_R$  and  $IR_I$  and either occurs by two possible methods shown in Fig. 3A (see text for more details). C. Sequence of the 'empty' *recA*<sub>S24</sub> insertion site and translated peptide sequence shows excision restores an uninterrupted *recA*<sub>S24</sub>. The asterisk marks the point of RME insertion. Amplification of the 'empty' *recA*<sub>S24</sub> site gave a faint product (marked by diamond in left gel image). This was excised, purified and a nested PCR (right gel image) was conducted to generate sufficient product for sequencing.

four independent experiments where pOriVn<sub>700</sub>-P<sub>lac</sub>*gfp* was successfully introduced into *V. cholerae* S24 by conjugation, a product was never detected in the transconjugates (see representative gel in Fig. 4B). However, when pOriVn<sub>700</sub>-*recA*<sub>S22</sub> was successfully introduced into *V. cholerae* S24, products were amplified (see representative gel in Fig. 4B) demonstrating insertion of the RME in the equivalent DNA site of *recA*<sub>S22</sub> and in both orientations, with respect to *recA*<sub>S22</sub>. Sequence of the PCR products are shown in Fig. 4C, demonstrating successful insertion of RME into the equivalent *recA*<sub>S24</sub> insertion site into *recA*<sub>S22</sub> in pOriVn<sub>700</sub>-*recA*<sub>S22</sub>.

It should be noted that homologous recombination between *recA*<sub>S22</sub> in pOriVn<sub>700</sub>-*recA*<sub>S22</sub> and *recA*<sub>S24</sub> in the *V. cholerae* S24 genome could result in merodiploids that generate the same amplicons as those for RME inserted in the orientation (relative to *recA*<sub>S24</sub>) found in *V. cholerae* S24 (see Supporting Information Fig. S1 on expected merodiploids). However, in the immediate 2 bp of the  $IR_R$  end for three of the transconjugates (Fig. 4C; ia, iia and iv), there is a G to T substitution and at the immediate 3 bp of the  $IR_I$  end, two transconjugants (Fig. 4C; iia and iv) showed a T to G substitution. Since the *recA*<sub>S22</sub> sequence is identical to *recA*<sub>S24</sub> around the insertion point, homologous recombination should result in identical sequences immediately surrounding the RME. Further-

more, the RME was also found in both orientations, with respect to *recA*<sub>S22</sub> (Fig. 4C). Consequently, homologous recombination is unable to explain these results.

These data show that RME is capable of mobilization and preferentially targets a specific site within *recA*. By carrying its own functional copy of *recA*, the GI does not affect any of the vital cell pathways associated with disruption of this gene during integration. Furthermore, specific targeting of *recA* may be necessary to ensure successful maintenance and dissemination of the GI. Since RecA does not function as a monomer but polymerizes to form a filament structure (Yu *et al.*, 2004), disruption of the indigenous *recA* prevents a situation where two divergent RecA proteins might negatively interact resulting in reduced cell fitness.

#### *The recA mobile element provides E. coli protection from UV irradiation*

The presence of multiple genes involved in DNA repair prompted us to look at whether the GI could protect against a common DNA-damaging process faced by *V. cholerae* – UV irradiation. To investigate if *recA*<sub>RME</sub> has a role in protecting the cell from DNA damage, the RME was cloned into a fosmid and used to transform *recA*<sup>-</sup> *E. coli* strain EPI300. The resultant transformant was

**Table 1.** Primers used in this study.

Primer	Sequence (5'-3')	Target	Source
RME-R	GACGAGTCCAGCTCATGACA	integrase end of <i>recA</i> genomic island	This study
RME-F	GCTGCTAACGCTTTCTGCTT	<i>recA</i> end of <i>recA</i> genomic island	This study
S24-ctg675-F	CGGTTAGGAGGGGCTTTTAG	3' end of contig 675	This study
S24-ctg708-R	TATCGGCTGTGGTTGTTTGA	5' end of contig 675	This study
S24-ctg367-F	TAGCTAGAGCATTGTCATAAGAAAAAGTAAG	3' end of contig 675	This study
S24-ctg367-R	ACTGGCAGCAGAAGAAGCAT	5' end contig 708	This study
S24-cinA-F	CAAGGTTGGCTCAAAGTG	<i>cinA</i> in <i>V. cholerae</i> S24	This study
S24-recX-R	GGCATCACTCAAATACCCTA	<i>recX</i> in <i>V. cholerae</i> S24	This study
S24-recA-F	CTGGAATTTGTGATGCATT	<i>recA</i> in <i>V. cholerae</i> S24	This study
EcoRI-recA-F <sup>a</sup>	TTTT <b><u>GAATTC</u></b> TGGACGAGAATAAACAGAAGG	<i>recA</i> in <i>V. cholerae</i> S22 & S24	This study
EcoRI-recA-R <sup>a</sup>	TTTT <b><u>GAATTC</u></b> AAACTCTTCTGGCACC	<i>recA</i> in <i>V. cholerae</i> S22 & S24	This study
EcoRI-Ori700-R <sup>a</sup>	TTTT <b><u>GAATTC</u></b> CGCGCTATCGCTTGTCG	<i>ori<sub>pB1067</sub></i> of pOriVn700	This study
EcoRI-OriR6K-F <sup>a</sup>	TTTT <b><u>GAATTC</u></b> GTGTTCTGTGTCCTCAAAATTG	<i>ori6k</i>	This study
Ori700-F	CCCTATTCCCTCTTTAGTCCCTGC	<i>ori<sub>pB1067</sub></i> of pOriVn700	This study
Ori6K-R	TAACGCACTGAGAAGCCC	<i>ori6k</i>	This study
S24-phage-Int-F	GCCAAGATATGGCAGGAAAA	Integrase in <i>recA</i> genomic island	This study
S24-phage-Int-R	GGACGCTACCCAGTGAATGT	Integrase in <i>recA</i> genomic island	This study
<i>recA</i> -F	TGGACGAGAATAAACAGAAGGC	<i>recA</i>	(Boucher <i>et al.</i> , 2011)
<i>recA</i> -R	CCGTTATAGCTGTACCAAGCGCCC	<i>recA</i>	(Boucher <i>et al.</i> , 2011)
pCC2FOS-FP	GTACAACGACACCTAGAC	pCC2FOS sequencing primers (F)	Epicentre Biotechnologies
pCC2FOS-RP	CAGGAAACAGCCTAGGAA	pCC2FOS sequencing primers (R)	Epicentre Biotechnologies
<i>recA</i> -Tn5-F	CGCTCATAAGTCAGTAATGCTTCA	<i>recA</i> on genomic island. Used to screen for Tn5 insertion.	This study
<i>umuC</i> -Tn5-F	GATGTATGGCTGAATCGACCA	<i>umuC</i> on genomic island. Used to screen for Tn5 insertion.	This study
KAN-2 FP-1	ACCTACAACAAAGCTCTCATCAACC	Forward primer inside Tn5 used to screen for Tn5 insertion.	Epicentre Biotechnologies
KAN-2 RP-1	GCAATGTAACATCAGAGATTTTGAG	Reverse primer inside Tn5 used to screen for Tn5 insertion.	Epicentre Biotechnologies

a. Bold and underlined sequence shows the *EcoRI* restriction site.

subjected to UV-C irradiation. Fig. 5A shows that the presence of the RME element conferred enhanced bacterial cell survival when exposed to 0.8 mJ cm<sup>-2</sup> of UV-C. From Fig. 5A and B it can be seen that EPI300 and EPI300 transformed by vector only controls are completely killed by exposure to 20 s 0.8 mJ cm<sup>-2</sup> of UV-C. However, EPI300 transformants containing the RME survive for up to 60 s of UV-C exposure and show a 100-fold increase in survival at 10 s and up to 10,000,000-fold higher survival after 20 s UV-C exposure. Fig. 5A shows that when *recA*<sub>RME</sub> is insertionally inactivated, the level of cell survivability decreases to a level comparable with the vector-only control (Fig. 5B). This demonstrates that *recA*<sub>RME</sub> is functional and is the gene mainly responsible for the protection provided by the presence of the RME. An interesting future question would be whether *recA*<sub>RME</sub> is more efficient in DNA repair

than the host *recA* (i.e. *recA*<sub>S24</sub>). There is precedent for such an idea, in a strain of *Clostridium difficile*, a 4.2 kb insert disrupts a gene encoding a thymidylate synthetase (involved in DNA synthesis and repair) but contains a more functionally active version of the disrupted gene (Knetsch *et al.*, 2011).

#### *The recA mobile element provides E. coli with increased protection against antibiotics*


Since RME has multiple genes involved in DNA repair, we tested whether RME provided enhanced protection against three DNA-targeting antibiotics: nalidixic acid, ciprofloxacin and bleomycin. Minimum inhibitory concentrations (MICs) were determined using nalidixic acid, ciprofloxacin and bleomycin (Table 2). Interestingly, the MIC of the first-generation quinolone, nalidixic acid, did

**Fig. 4.** Translocation of the *recA* genomic island from the genome of *V. cholerae* S24 into a replicating vector containing *recA*<sub>S22</sub>.

A. Genetic structure of the replicating vectors pOriVn700-*recA*<sub>S22</sub> and pOriVn700-*P<sub>lac</sub>Gfp* and the placement of primers *ori6k*-R and *ori700*-F used for amplifying the boundaries of the translocated genomic island are shown.

B. Representative gel showing amplification using vector specific and RME specific primers from colonies derived from conjugation of pOriVn700-*recA*<sub>S22</sub> (lanes 1 and 2) and pOriVn700-*P<sub>lac</sub>Gfp* (lanes 3 and 4) into *V. cholerae* S24. Lane 5 shows negative dH<sub>2</sub>O control.

C. Sequence of products derived using vector-specific and RME-specific primers from PCR of *V. cholerae* S24 transconjugates from four independent conjugations. Each transconjugant is denoted by i, ii, iii and iv. In most instances (iib, iiib and ivb), the same transconjugate showed insertion of RME in both orientations relative to *recA*<sub>S22</sub>. The sequences indicate specific insertion of RME into the same site of *recA*<sub>S22</sub> (the equivalent insertion site in *recA*<sub>S24</sub>).



**Table 2.** Minimal inhibitory concentration (MICs<sup>a</sup>).


Strain	Ciprofloxacin	Bleomycin
EPI300	0.015625	8
EPI300/pCC2FOS	0.015625	8
EPI300/pCC2FOS::RME	0.0625	16
EPI300/pCC2FOS::RME $\Delta$ umuC <sub>RME</sub>	0.0625	8
EPI300/pCC2FOS::RME $\Delta$ recA <sub>RME</sub>	0.015625	8

a. MIC given as  $\mu\text{g ml}^{-1}$ .

not vary between any of the tested strains, including the RME (data not shown). However, when the strains given in Table 2 were tested using ciprofloxacin, a second-generation quinolone, a fourfold increase in MIC was observed for strains containing the RME. In the case of ciprofloxacin, it is apparent that *recA*<sub>RME</sub> is responsible for the increased resistance. When *recA*<sub>RME</sub> is insertionally inactivated from the genomic island, the MIC drops to a level equivalent to that seen for the *E. coli* strain EPI300. The importance of *recA* in protection against ciprofloxacin and other antibiotics has previously been documented [e.g. in *Acinetobacter baumannii* (Aranda et al., 2011)]. RME also provided protection from bleomycin in *E. coli* EPI300. However, in contrast to ciprofloxacin, when *recA*<sub>RME</sub> and *umuC* from the RME (designated *umuC*<sub>RME</sub>)

are both inactivated independently, the MIC against bleomycin is the same as *E. coli* EPI300, indicating that *RecA*<sub>RME</sub> activation of the DNA polymerase V subunit UmuD to UmuD' encoded on the GI is responsible for protections against bleomycin (Patel et al., 2010). Since *umuDC*<sub>RME</sub> was able to provide protection from bleomycin in a genetic background that already contains *umuDC*, it is hypothesized that in the *V. cholerae* species, where *umuDC* is only sporadically found, this element may provide increased protection from DNA damage compared with what was observed here in *E. coli* EPI300.

Apart from DNA repair, *recA* and DNA polymerase V (encoded by *umuDC*) are known to increase spontaneous mutation frequencies resulting in the emergence of antibiotic resistance mutants (Thi et al., 2011). Spontaneous mutation in *V. cholerae* is well documented to cause resistance to a variety of antibiotics (Goss et al., 1965; Gellert et al., 1977; Sugino et al., 1977; Allen et al., 1979; Kitaoka et al., 2011). Here we chose to examine the mutation frequency of two antibiotics, rifampicin which acts on protein synthesis and nalidixic acid which targets DNA replication by inhibiting the A subunit of DNA gyrase. Mutation frequencies after 24 and 48 h showed no differences between *E. coli* containing RME and the controls on 100  $\mu\text{g ml}^{-1}$  rifampicin (data not shown). This may be because of rifampicin acting on protein synthesis and therefore not inducing the SOS response which induces transcription of *umuDC*. However, when the experiments were repeated with 50  $\mu\text{g ml}^{-1}$  nalidixic acid, *E. coli* EPI300 and the vector-only control consistently did not produce any spontaneous mutants, while *E. coli* EPI300/pCC2FOS::RME showed an increased mutation frequency after both 24 and 48 h incubation in the presence of nalidixic acid (Table 3). *E. coli* EPI300/pCC2FOS::RME $\Delta$ umuC<sub>RME</sub> generally had the same mutation frequency as the complete RME. However, experiment 3 (Table 3) showed that this strain produced no mutants after 24 and 48 h. It can be concluded from these experiments that *E. coli* EPI300 with the complete RME provides an adaptive advantage by increasing the mutation rate resulting in subsequent resistance to nalidixic acid, but this effect could not be wholly attributed to the activity of *umuC*<sub>RME</sub>. One possible explanation is the activation of the indigenous *E. coli* UmuD by *RecA* provided by the GI. However, it cannot be excluded that other genes on the RME are elevating the spontaneous mutation rate. Nevertheless, *umuDC*-like operons are commonly associated with mobile genetic elements (Permina et al., 2002; Tark et al., 2005; Hare et al., 2012) and do provide a general adaptive advantage to hosts that house them (Yeiser et al., 2002; Tark et al., 2005). Although we failed in our attempts to transfer RME into seven non-O1/O139 *V. cholerae* strains from Sydney using chitin



**Fig. 5.** Survival of *E. coli* carrying the *recA* genomic island and control strains when exposed to UV-C stress. Time points are given at 0, 10, 20, 40 and 60 s. UV-C exposure was set to 0.8  $\text{mJ cm}^{-2}$ .



transformation, this element is likely to do the same in a *V. cholerae* genetic background.

To conclude, this study reports a novel GI in *V. cholerae* that contains genes involved in multiple DNA repair pathways, including the critical housekeeping gene *recA* and genes encoding DNA polymerase V which in this study, we show to be functional. The presence of other DNA processing genes may provide *V. cholerae* with alternative DNA repair pathways. Since this element can excise from its chromosomal location, it has the potential to mobilize into other strains, such as cholera toxin-producing O1/O139 pandemic strains. Such mobilization could have implications for increased environmental survival or resistance to certain antibiotics.

## Experimental procedures

### Bacterial strains, plasmids and growth conditions

All strains and plasmids used are shown in Table 4. *V. cholerae* strain S24 was collected from Georges River in the greater Sydney (Australia) urban area as previously described (Islam *et al.*, 2013). All *E. coli* and *V. cholerae* strains were routinely grown on Luria–Bertani (LB) broth at 37°C under aerobic conditions. For *E. coli* WM3064, diamminopimelic acid (DAP) was added to a final concentration of 0.3 mM. Spectinomycin was used for *E. coli* and *V. cholerae* at 50 µg ml<sup>-1</sup> and 125 µg ml<sup>-1</sup> respectively. Chloramphenicol was used at 12.5 µg ml<sup>-1</sup>.

### Whole genome sequencing, PCR, DNA extraction and sequencing methods

DNA was extracted using the Wizard genomic DNA purification kit (Promega). Plasmid and PCR/gel extractions were done using PureYield Plasmid Miniprep and Wizard SV Gel and PCR clean-up systems respectively (Promega). Purified DNA from *V. cholerae* S24 was sequenced at the Wellcome Trust Sanger Institute using Illumina-based technology.

All primers used in this study are shown in Table 1. Standard PCR was performed using the PCR master mix (Promega) containing 25 units ml<sup>-1</sup> of *Taq* DNA polymerase, 800 µM dNTPs and 1.5 mM MgCl<sub>2</sub>. Primers were used at a final concentration of 0.5 µM each. All PCRs were performed with 30 cycles of denaturation at 94°C for 30 s, the appropriate annealing temperature for 30 s and an extension of 72°C (1 min kb<sup>-1</sup>) and sequencing performed at Macrogen. From whole genome sequencing (Wellcome Trust Sanger Institute) it became evident that the host *recA* had been disrupted and was present on two separate contigs. The two contigs (contigs 675 and 708) were pieced together by PCR and joined to an intervening third contig to produce a contig of 262,869 bp (contig 367) using primers described in Table 1. The accession number for RME is KJ123688.

### Cloning of RME and transposon mutagenesis of recA genomic island

To clone the *recA* genomic island (RME) from *V. cholerae* strain S24, genomic DNA was digested with *Nae*I and a

**Table 3.** Nalidixic acid<sup>a</sup> mutation frequencies.

Experiment 1		
Strain	Mutation frequency 24 h	Mutation frequency 48 h
EPI300	< 1.1 × 10 <sup>-11</sup>	< 1.1 × 10 <sup>-11</sup>
EPI300/pCC2FOS	< 1.7 × 10 <sup>-11</sup>	< 1.7 × 10 <sup>-11</sup>
EPI300/pCC2FOS::RME	1.4 × 10 <sup>-8</sup> (4)	1.5 × 10 <sup>-8</sup> (4)
EPI300/pCC2FOS::RMEΔ <i>umu</i> C <sub>RME</sub>	1.6 × 10 <sup>-9</sup> (3)	2.1 × 10 <sup>-9</sup> (4)
Experiment 2		
Strain	Mutation frequency 24 h	Mutation frequency 48 h
EPI300	< 1.1 × 10 <sup>-11</sup>	< 1.1 × 10 <sup>-11</sup>
EPI300/pCC2FOS	< 1.7 × 10 <sup>-11</sup>	< 1.7 × 10 <sup>-11</sup>
EPI300/pCC2FOS::RME	8.3 × 10 <sup>-10</sup> (1)	1.4 × 10 <sup>-9</sup> (3)
EPI300/pCC2FOS::RMEΔ <i>umu</i> C <sub>RME</sub>	2.8 × 10 <sup>-9</sup> (6)	4.4 × 10 <sup>-9</sup> (7)
Experiment 3		
Strain	Mutation frequency 24 h	Mutation frequency 48 h
EPI300	< 1.1 × 10 <sup>-11</sup>	< 1.1 × 10 <sup>-11</sup>
EPI300/pCC2FOS	< 1.3 × 10 <sup>-11</sup>	< 1.3 × 10 <sup>-11</sup>
EPI300/pCC2FOS::RME	4.8 × 10 <sup>-10</sup> (1)	1.7 × 10 <sup>-9</sup> (4)
EPI300/pCC2FOS::RMEΔ <i>umu</i> C <sub>RME</sub>	< 9.1 × 10 <sup>-12</sup>	< 9.1 × 10 <sup>-12</sup>

a. Concentration of nalidixic acid = 50 µg ml<sup>-1</sup>.

'<' indicates that zero colonies appeared in all 10 replicates (see *Experimental procedures*).

Numbers in brackets indicates the number of replicates in which one or more colonies appeared.

**Table 4.** List of strains and plasmids.

Strain or plasmid	Relevant genotype <sup>a</sup>	Reference or source
<i>V. cholerae</i> S24	Wild-type (non-O1/O139)	This study
S22	Wild-type (non-O1/O139)	(Islam <i>et al.</i> , 2013)
<i>E. coli</i> DH5 $\alpha$ pir	<i>endA1 glnV44 thi-1 recA1 relA1 gyrA96 deoR nupG</i> $\Phi$ 80d <i>lacZ</i> $\Delta$ M15 $\Delta$ ( <i>lacZYA-argF</i> )U169 <i>hsdR17</i> $\lambda$ pir	(Demarre <i>et al.</i> , 2005)
WM3064	Donor strain for conjugation: <i>thrB1004 pro thi rpsL hsdS lacZ</i> $\Delta$ M15 RP4-1360 $\Delta$ ( <i>araBAD</i> )567 $\Delta$ <i>dapA1341::[erm pir]</i> , Sm <sup>R</sup>	(Saltikov and Newman, 2003)
EPI300 <sup>TM</sup> -T1 <sup>R</sup>	[F <sup>-</sup> <i>mcrA</i> $\Delta$ ( <i>mrr-hsdRMS-mcrBC</i> ) $\Phi$ 80d <i>lacZ</i> $\Delta$ M15 $\Delta$ <i>lacX74 recA1 endA1 araD139</i> $\Delta$ ( <i>ara, leu</i> )7697 <i>galK</i> $\lambda$ : <i>rpsL nupG trfA tonA dhfr</i> ], Sm <sup>R</sup> , Tp <sup>R</sup>	Epicentre Biotechnologies
Plasmids/fosmids		
pCC2FOS	Cloning vector, Cm <sup>R</sup>	
pCC2FOS-RME	pCC2FOS vector containing 32 kb insert from <i>V. cholerae</i> S24. The insert contains the <i>recA</i> GI and surrounding sequence, Cm <sup>R</sup>	This study
pCC2FOS::RME $\Delta$ <i>recA</i> <sub>RME</sub>	pCC2FOS vector containing 32 kb insert from <i>V. cholerae</i> S24. The insert contains the <i>recA</i> GI and surrounding sequence and has <i>recA</i> on the GI insertionally inactivated by Tn5, Km <sup>R</sup> , Cm <sup>R</sup>	This study
pCC2FOS::RME $\Delta$ <i>umuC</i> <sub>RME</sub>	pCC2FOS vector containing 32 kb insert from <i>V. cholerae</i> S24. The insert contains the <i>recA</i> GI and surrounding sequence and has <i>umuC</i> present on the GI insertionally inactivated by Tn5, Km <sup>R</sup> , Cm <sup>R</sup>	This study
pOriVn <sub>700</sub>	Low copy mobilizable vector containing <i>ori</i> <sub>pB1067</sub> (vibrio specific) and <i>ori6K</i> , Sp <sup>R</sup>	(Le Roux <i>et al.</i> , 2011)
pOriVn <sub>700</sub> - <i>recA</i> <sub>S22</sub>	pOriVn <sub>700</sub> with <i>recA</i> from <i>V. cholerae</i> S22 in between <i>ori</i> <sub>pB1067</sub> and <i>ori6K</i> . The <i>recA</i> gene is reading toward <i>ori</i> <sub>pB1067</sub> , Sp <sup>R</sup>	This study
pOriVn <sub>700</sub> -P <sub>lac</sub> <i>gfp</i>	pOriVn <sub>700</sub> with P <sub>lac</sub> <i>gfp</i> cloned in between <i>ori</i> <sub>pB1067</sub> and <i>ori6K</i> , Sp <sup>R</sup>	(Le Roux <i>et al.</i> , 2011)

a. Tc<sup>R</sup>, tetracycline resistance; Sm<sup>R</sup>, streptomycin resistance; Sp<sup>R</sup>, spectinomycin resistance; Cm<sup>R</sup>, chloramphenicol resistance, Km<sup>R</sup>, kanamycin resistance.

library constructed using the CopyControl Fosmid Library Production Kit (Epicentre). *NaeI* digestion of *V. cholerae* strain S24 genomic DNA creates a fragment of 38, 913 bp containing the entire 32, 787 bp RME. The library was screened for a fosmid clone containing the 38, 913-bp *NaeI* fragment using primers targeting the phage integrase in the RME (Table 1). A positive clone designated pCC2FOS-RME was confirmed by sequencing the ends of the cloned insert using the pCC2FOS vector primers FP and RP (Table 1). To create the pCC2FOS no insert control, linearized and dephosphorylated pCC2FOS (Epicentre) was treated with T4 polynucleotide kinase and circularized by ligation. A mutant library of pCC2FOS-RME was constructed using the EZ-Tn5 Kan-2 Insertion Kit (Epicentre Biotechnologies) according to manufacturer instructions. Mutants containing knockouts of individual genes present on the genomic island were screened by PCR using primers reading out from EZ-Tn5 Kan-2 and a primer targeting the gene of interest (Table 1).

#### Phylogenetic analysis

Phylogenetic analysis of *recA*<sub>S24</sub> and *recA*<sub>RME</sub> was done using bioinformatics program Geneious version 6.1.6 and FigTree version 1.4.0. Phylogenetic tree parameters were taken from (Thompson *et al.*, 2004). Distance estimations were obtained using the Jukes and Cantor model and tree built using the neighbour-joining method. Bootstrap percentages were calculated after 100 simulations. The *Campylobacter jejuni* subsp. *jejuni* NCTC 11168 *recA* sequence was used as an outgroup.

#### *recA* targeting experiments

Vector pOriVn<sub>700</sub> and *recA* from a strain of *V. cholerae* S22 that is closely related to *V. cholerae* S24 were amplified using primer pairs EcoRI-Ori700-R/EcoRI-Ori6K-F and EcoRI-*recA*-F/EcoRI-*recA*-R respectively (Table 1). Since the primers contained engineered *EcoRI* sites, the resulting amplicons of *recA*<sub>S22</sub> and pOriVn<sub>700</sub> were purified, digested with *EcoRI* and then ligated together using T4 DNA ligase (Fermentas). The ligation mix was then transformed into *E. coli* DH5 $\alpha$ pir to produce pOriVn<sub>700</sub>-*recA*<sub>S22</sub>. The construct was then extracted and transformed into the conjugation donor strain *E. coli* WM3064.

Conjugations using pOriVn<sub>700</sub>-*recA*<sub>S22</sub> and pOriVn<sub>700</sub>-P<sub>lac</sub>*gfp* were performed by combining equal volumes of overnight cultures in LB from both donor and recipient strains. These were then centrifuged at 3000  $\times$  *g* and re-suspended in 50  $\mu$ L of LB and spotted onto a 0.2  $\mu$ M filter (Millipore) that had been placed on an LB agar plate containing 0.3 mM DAP. Donor and recipient cells were left to incubate for 4 h at 37°C and cells were then removed from the filter by vortexing. The re-suspended cells were then plated on LB + 125  $\mu$ g ml<sup>-1</sup> spectinomycin and incubated at 37°C overnight. One colony per mating was picked and appropriate junction PCR was conducted using primers in plasmid backbone (Table 1; Ori700-F/Ori6K-R) and primers reading out from RME (Table 1; RME-F/RME-R).

#### UV stress experiments

UV stress experiments were adapted from Lin and Wang (2001). Strains were grown for 16–20 h at 37°C with shaking

at 230 r.p.m. in 5 ml LB broth supplemented with appropriate antibiotic. Cells were centrifuged at  $4000 \times g$ , corrected for differences in optical density (OD) at 600 nm and re-suspended in equal volumes of M9 salts (Sambrook *et al.*, 1989) supplemented with  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  to a final concentration of 0.002 M. The entire cell suspension was placed in a clear bottom 10 cm plastic petri dish and subjected to  $0.8 \text{ mJ cm}^{-2}$  UV-C for 0, 10, 20, 40 and 60 s using an Amersham Life Science Ultraviolet Crosslinker. After each time interval, 150  $\mu\text{l}$  aliquot was removed and placed in a 1.5 ml Eppendorf tube in the dark. The remaining liquid culture was thoroughly re-suspended using a pipette to avoid clumping of cells. After the final UV-C exposure time point, cells were diluted in M9 salts+  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  to  $10^{-6}$  and enumerated by the drop plate method on LB agar. Plates were incubated in the dark to prevent photoreactivation at  $37^\circ\text{C}$  overnight and colony-forming units (CFUs) were calculated the following day.

#### Minimum inhibitory concentration experiments and antibiotic mutation frequency experiments

MICs of nalidixic acid, ciprofloxacin and bleomycin were determined by broth microdilution using standard methods (Clinical and Laboratory Standards Institute, 2003) except that LB broth was used as the growth medium instead of Mueller Hinton. Each MIC was performed in triplicate. The mutation frequency experiment was designed using the guidelines described in (Pope *et al.*, 2008). Specifically, mutation frequencies were determined using LB supplemented with  $50 \mu\text{g ml}^{-1}$  nalidixic acid and  $100 \mu\text{g ml}^{-1}$  rifampicin. Ten replicate overnight cultures for each strain were grown in 5 ml LB (chloramphenicol was added for those strains carrying pCC2FOS and derivatives). Each overnight culture was then diluted to  $\sim 10^4$  CFU  $\text{ml}^{-1}$  with fresh LB5 (no chloramphenicol added) and 5 ml for each replicate was transferred into a 15 ml tube and incubated for 16–20 h at  $37^\circ\text{C}$  with shaking at 230 r.p.m. The following day, 200  $\mu\text{l}$  from each tube was spread plated onto LB5 agar supplemented with the appropriate antibiotic (rifampicin or nalidixic acid) and incubated for 24 h and then 48 h at  $37^\circ\text{C}$  when colonies were counted. This was repeated in triplicate. In order to calculate total colony counts, cells were enumerated on LB5 agar with no antibiotic. Note that these experiments were performed in a Class II Biosafety Hood to avoid any contamination. Mutation frequencies were calculated as number of antibiotic-resistant CFUs/total number of CFUs after 24 and 48 h.

#### Acknowledgements

We would like to acknowledge Associate Professor Aaron Darling for his help with the phylogenetic analysis. This work was supported by the ithree institute, UTS. Rita Rapa is a recipient of a UTS postgraduate scholarship. The sequencing of *V. cholerae* S24 was funded under Wellcome Trust grant 098051.

#### References

Allen, J.G., Atherton, F.R., Hall, M.J., Hassall, C.H., Holmes, S.W., Lambert, R.W., *et al.* (1979) Phosphonopeptides

as antibacterial agents: alaphosphin and related phosphonopeptides. *Antimicrob Agents Chemother* **15**: 684–695.

Aranda, J., Bardina, C., Beceiro, A., Rumbo, S., Cabral, M.P., Barbe, J., and Bou, G. (2011) *Acinetobacter baumannii* RecA protein in repair of DNA damage, antimicrobial resistance, general stress response, and virulence. *J Bacteriol* **193**: 3740–3747.

Baptiste, E., Boucher, Y., Leigh, J., and Doolittle, W.F. (2004) Phylogenetic reconstruction and lateral gene transfer. *Trends Microbiol* **12**: 406–411.

Boucher, Y., Cordero, O.X., Takemura, A., Hunt, D.E., Schliep, K., Baptiste, E., *et al.* (2011) Local mobile gene pools rapidly cross species boundaries to create endemicity within global *Vibrio cholerae* populations. *MBio* **2**: e00335-10.

Boyd, E.F., Almagro-Moreno, S., and Parent, M.A. (2008) Genomic islands are dynamic, ancient integrative elements in bacterial evolution. *Trends Microbiol* **17**: 47–53.

Clinical and Laboratory Standards Institute. (2003) *Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically*.

Creevey, C.J., Fitzpatrick, D.A., Philip, G.K., Kinsella, R.J., O'Connell, M.J., Pentony, M.M., *et al.* (2004) Does a tree-like phylogeny only exist at the tips in the prokaryotes? *Proc Biol Sci* **271**: 2551–2558.

Demarre, G., Guérout, A., Matsumoto-Mashimo, C., Rowe-Magnus, D.A., Marlière, P., and Mazel, D. (2005) A new family of mobilizable suicide plasmids based on broad host range R388 plasmid (IncW) and RP4 plasmid (IncP $\alpha$ ) conjugative machineries and their cognate *Escherichia coli* host strains. *Res Microbiol* **156**: 245–255.

Everiss, K.D., Hughes, K.J., and Peterson, K.M. (1994) The accessory colonization factor and toxin-coregulated pilus gene clusters are physically linked on the *Vibrio cholerae* 0395 chromosome. *DNA Seq* **5**: 51–55.

Faruque, S.M., and Mekalanos, J.J. (2003) Pathogenicity islands and phages in *Vibrio cholerae* evolution. *Trends Microbiol* **11**: 505–510.

Galen, J.E., Ketley, J.M., Fasano, A., Richardson, S.H., Wasserman, S.S., and Kaper, J.B. (1992) Role of *Vibrio cholerae* neuraminidase in the function of cholera toxin. *Infect Immun* **60**: 406–415.

Gellert, M., Mizuuchi, K., O'Dea, M.H., Itoh, T., and Tomizawa, J.I. (1977) Nalidixic acid resistance: a second genetic character involved in DNA gyrase activity. *Proc Natl Acad Sci USA* **74**: 4772–4776.

Goodman, M.F. (2002) Error-prone repair DNA polymerases in prokaryotes and eukaryotes. *Annu Rev Biochem* **71**: 17–50.

Goss, W.A., Deitz, W.H., and Cook, T.M. (1965) Mechanism of action of nalidixic acid on *Escherichia coli*. Inhibition of deoxyribonucleic acid synthesis. *J Bacteriol* **89**: 1068–1074.

Grim, C.J., Hasan, N.A., Taviani, E., Haley, B., Chun, J., Brettin, T.S., *et al.* (2010) Genome sequence of hybrid *Vibrio cholerae* O1 MJ-1236, B-33, and CIRS101 and comparative genomics with *V. cholerae*. *J Bacteriol* **192**: 3524–3533.

Hallet, B., and Sherratt, D.J. (1997) Transposition and site-specific recombination: adapting DNA cut-and-paste

- mechanisms to a variety of genetic rearrangements. *FEMS Microbiol Rev* **21**: 157–178.
- Hare, J.M., Bradley, J.A., Lin, C., and Elam, T.J. (2012) Diverse responses to UV light exposure in *Acinetobacter* include the capacity for DNA damage-induced mutagenesis in the opportunistic pathogens *Acinetobacter baumannii* and *Acinetobacter ursingii*. *Microbiology* **158**: 601–611.
- Islam, A., Labbate, M., Djordjevic, S.P., Alam, M., Darling, A., Melvold, J., et al. (2013) Indigenous *Vibrio cholerae* strains from a non-endemic region are pathogenic. *Open Biol* **3**: 120181.
- Janion, C. (2008) Inducible SOS response system of DNA repair and mutagenesis in *Escherichia coli*. *Int J Biol Sci* **4**: 338–344.
- Kaper, J., Morris, J., Jr, and Levine, M. (1995) Cholera. *Clin Microbiol Rev* **8**: 48–86.
- Keymer, D.P., and Boehm, A.B. (2011) Recombination shapes the structure of an environmental *Vibrio cholerae* population. *Appl Environ Microbiol* **77**: 537–544.
- Kitaoka, M., Miyata, S.T., Unterweger, D., and Pukatzki, S. (2011) Antibiotic resistance mechanisms of *Vibrio cholerae*. *J Med Microbiol* **60**: 397–407.
- Knetsch, C.W., Hensgens, M.P., Harmanus, C., van der Bijl, M.W., Savelkoul, P.H., Kuijper, E.J., et al. (2011) Genetic markers for *Clostridium difficile* lineages linked to hypervirulence. *Microbiology* **157**: 3113–3123.
- Le Roux, F., Davis, B.M., and Waldor, M.K. (2011) Conserved small RNAs govern replication and incompatibility of a diverse new plasmid family from marine bacteria. *Nucleic Acids Res* **39**: 1004–1013.
- Lenhart, J.S., Schroeder, J.W., Walsh, B.W., and Simmons, L.A. (2012) DNA repair and genome maintenance in *Bacillus subtilis*. *Microbiol Mol Biol Rev* **76**: 530–564.
- Lin, K., and Wang, A. (2001) UV mutagenesis in *Escherichia coli* K-12: cell survival and mutation frequency of the chromosomal genes *lacZ*, *rpoB*, *ompF*, and *ampA*. *J Exp Microbiol Immunol* **1**: 32–46.
- McKenzie, G.J., Harris, R.S., Lee, P.L., and Rosenberg, S.M. (2000) The SOS response regulates adaptive mutation. *Proc Natl Acad Sci USA* **97**: 6646–6651.
- Meibom, K.L., Blokesch, M., Dolganov, N.A., Wu, C.-Y., and Schoolnik, G.K. (2005) Chitin induces natural competence in *Vibrio cholerae*. *Science* **310**: 1824–1827.
- Murphy, R.A., and Boyd, E.F. (2008) Three pathogenicity islands of *Vibrio cholerae* can excise from the chromosome and form circular intermediates. *J Bacteriol* **190**: 636–647.
- Patel, M., Jiang, Q., Woodgate, R., Cox, M.M., and Goodman, M.F. (2010) A new model for SOS-induced mutagenesis: how RecA protein activates DNA polymerase V. *Crit Rev Biochem Mol Biol* **45**: 171–184.
- Permina, E.A., Mironov, A.A., and Gelfand, M.S. (2002) Damage-repair error-prone polymerases of eubacteria: association with mobile genome elements. *Gene* **293**: 133–140.
- Pidcock, L.J., and Wise, R. (1987) Induction of the SOS response in *Escherichia coli* by 4-quinolone antimicrobial agents. *FEMS Microbiol Lett* **41**: 289–294.
- Polosina, Y.Y., and Cupples, C.G. (2010) Wot the 'L-Does MutL do? *Mutat Res* **705**: 228–238.
- Pope, C.F., O'Sullivan, D.M., McHugh, T.D., and Gillespie, S.H. (2008) A practical guide to measuring mutation rates in antibiotic resistance. *Antimicrob Agents Chemother* **52**: 1209–1214.
- Ramamurthy, T., Yamasaki, S., Takeda, Y., and Nair, G.B. (2003) *Vibrio cholerae* O139 Bengal: odyssey of a fortuitous variant. *Microbes Infect* **5**: 329–344.
- Ratray, A.J., and Strathern, J.N. (2003) Error-prone DNA polymerases: when making a mistake is the only way to get ahead. *Annu Rev Genet* **37**: 31–66.
- Safa, A., Bhuiya, A.N., Murphy, D., Bates, J., Nusrin, S., Kong, R.Y.C., et al. (2009) Multilocus genetic analysis reveals that the Australian strains of *Vibrio cholerae* O1 are similar to the pre-seventh pandemic strains of the El Tor biotype. *J Med Microbiol* **58**: 105–111.
- Saltikov, C.W., and Newman, D.K. (2003) Genetic identification of a respiratory arsenate reductase. *Proc Natl Acad Sci USA* **100**: 10983–10988.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular cloning: A laboratory manual*.
- Sanchez-Alberola, N., Campoy, S., Barbé, J., and Erill, I. (2012) Analysis of the SOS response of *Vibrio* and other bacteria with multiple chromosomes. *BMC Genomics* **13**: 58.
- Stine, O.C., Sozhamannan, S., Gou, Q., Zheng, S., Morris, J.G., Jr, and Johnson, J.A. (2000) Phylogeny of *Vibrio cholerae* based on *recA* sequence. *Infect Immun* **68**: 7180–7185.
- Sugino, A., Peebles, C.L., Kreuzer, K.N., and Cozzarelli, N.R. (1977) Mechanism of action of nalidixic acid: purification of *Escherichia coli* *nalA* gene product and its relationship to DNA gyrase and a novel nicking-closing enzyme. *Proc Natl Acad Sci USA* **74**: 4767–4771.
- Tark, M., Tover, A., Tarassova, K., Tegova, R., Kivi, G., Hórák, R., and Kivisaar, M. (2005) A DNA polymerase V homologue encoded by TOL plasmid pWW0 confers evolutionary fitness on *Pseudomonas putida* under conditions of environmental stress. *J Bacteriol* **187**: 5203–5213.
- Thi, T.D., Lopez, E., Rodriguez-Rojas, A., Rodriguez-Beltran, J., Couce, A., Guelfo, J.R., et al. (2011) Effect of *recA* inactivation on mutagenesis of *Escherichia coli* exposed to sublethal concentrations of antimicrobials. *J Antimicrob Chemother* **66**: 531–538.
- Thompson, C.C., Thompson, F.L., Vandemeulebroecke, K., Hoste, B., Dawyndt, P., and Swings, J. (2004) Use of *recA* as an alternative phylogenetic marker in the family *Vibrionaceae*. *Int J Syst Evol Microbiol* **54**: 919–924.
- Wertman, K.F., and Mount, D.W. (1985) Nucleotide sequence binding specificity of the LexA repressor of *Escherichia coli* K-12. *J Bacteriol* **163**: 376–384.
- Yeiser, B., Pepper, E.D., Goodman, M.F., and Finkel, S.E. (2002) SOS-induced DNA polymerases enhance long-term survival and evolutionary fitness. *Proc Natl Acad Sci USA* **99**: 8737–8741.
- Ysern, P., Clerch, B., Castaño, M., Gilbert, I., Barbé, J., and Llagostera, M. (1990) Induction of SOS genes in *Escherichia coli* and mutagenesis in *Salmonella typhimurium* by fluoroquinolones. *Mutagenesis* **5**: 63–66.

Yu, X., VanLoock, M.S., Yang, S., Reese, J.T., and Egelman, E.H. (2004) What is the structure of the RecA-DNA filament? *Curr Protein Pept Sci* **5**: 73–79.

### Supporting information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

**Fig. S1.** Possible insertion and homologous recombination events of the *recA* genomic island with pOriVn<sub>700</sub>-*recA*<sub>S22</sub>. Production of merodiploids because of homologous

recombination between *recA*<sub>S22</sub> and *recA*<sub>S24</sub> could result in two genetic structures (crossover 1 and crossover 2). In both instances, PCR products could be generated using the vector-specific and RME-specific primers used to detect insertion of the RME into *recA*<sub>S22</sub> (broken lines) in the orientation found in the *V. cholerae* S24 genome (insertion 1). Insertion of the RME in the inverse orientation (insertion 2) would generate products using inverse primer pairs. The inverse insertion cannot be explained by homologous recombination and indicates an integration event.

**Appendix S1.** Genbank file of the *recA* mobile element (RME); Accession Number KJ123688.