

Assessment of the endocrine disruption
potential of an advanced tertiary treated sewage
effluent using multiple lines of evidence

Lisa Anne Hamilton

A thesis submitted in fulfilment of the requirements for the degree of Doctor of
Philosophy in Science

University of Technology, Sydney

2010

CERTIFICATE OF AUTHORSHIP/ORIGINALITY

I certify that the work in this thesis has not previously been submitted for a degree nor has it been submitted as part of requirements for a degree except as fully acknowledged within the text.

I also certify that the thesis has been written by me. Any help that I have received in my research work and the preparation of the thesis itself has been acknowledged. In addition, I certify that all information sources and literature used are indicated in the thesis.

Signature of Student

Production Note:
Signature removed prior to publication.

Acknowledgements

First and foremost, I must acknowledge the Australian Research Council and the industry partner organisations Veolia Water Australia and Landcare Research, NZ for funding this project under ARC Linkage Grant LP0560600.

I would like to thank my supervisors and industry representatives Assoc. Prof. Richard Lim, Dr Grant Northcott, Dr Louis Tremblay and Mr Michael Boake for their support and guidance. A special thank you to Richard and Grant for providing mentorship and for both doing a stint assisting in intensive onsite sampling during the fish testing experiments.

I would like to thank Dr Fujio Shirarishi for training me in the two-hybrid yeast assay and for gifting the medER and hER two-hybrid yeasts to my laboratory group at UTS. Additionally I would like to acknowledge the contribution of Dr Fujio Shiraishi and Dr Daisuke Nakajima, in performing the two-hybrid yeast assay and *umu* test on some of the samples. Thanks go to Dr Toshihiro Horiguchi for facilitating the research exchange between NIES and UTS. I would also like to acknowledge and am very grateful to Dr Marianne Woods at CSIRO in Adelaide for analysing the rainbowfish for vitellogenin upregulation and to Dr Anu Kumar for facilitating this contribution.

A big thank you to Jeff Kidd and his team at Gerringong-Gerroa STP; Tiny, Rob, Steve, Pete and Tim for their support, assistance and patience during my onsite sampling and experiments. Thank you for installing the piping for the fish tests and letting me take over your lab.

Another big thanks goes to former fellow EDC students, Dr Chris Rawson, Dr Jenni Gadd and Heather Brown for advice, assistance, guidance and empathy and thanks to Chris Rawson, Alec Davie, Stephen Summer-Hayes and Alison Packham for field assistance collecting mosquitofish and Andrew 'batman' Smith for his week long stint as my lab assistant.

I am very grateful to those who provided me advice, guidance and training through the array of bioassays used in this project. In particular I would like to thank Dr Anne

Collville for her assistance in all things UTS and ecotox, Katherine Trought for training and assistance in the E-screen and ERBA, Dr Katherine James and Amanda Hudson for advice and guidance on qPCR.

Finally, I would like to thank my friends and family for their support, especially my parents who let me lock myself away in their mountain retreat during the thesis write-up process.

Table of Contents

List of Figures	xi
List of Tables	xv
Abstract	xvii
Abbreviations	xix
Chapter 1 Endocrine Disruptors in Domestic Wastewater: A review of the occurrence, methods of assessment and efficacy of current treatment technologies	1
1.1 General Introduction	2
1.1.1 Endocrine disrupting chemicals (EDCs)	3
1.1.2 Endocrine disruption from wastewater.....	4
1.1.3 Causative agents in wastewater	5
1.1.4 Complicating issues in assessing endocrine disruption	6
1.2 Modes of action of endocrine disrupting compounds	6
1.3 Chemical concentrations in wastewater	8
1.3.1 Estrogenic EDCs	8
1.3.1.1 Steroidal estrogens	10
1.3.1.2 Alkylphenol polyethoxylates and Bisphenol A	10
1.3.1.3 Phytoestrogens, phthalates and other xenoestrogens	13
1.4 Treatment technology and EDC removal	14
1.4.1 Partitioning behaviour of EDCs.....	14
1.4.2 Activated sludge treatment of sewage.....	15
1.4.3 Advanced tertiary treatment.....	17
1.5 Bioassay assessment of estrogenic endocrine disrupting chemicals	19

1.5.1	In vitro bioassays	20
1.5.2	In vivo bioassays	23
1.6	Research aim and objectives	24
1.7	Thesis structure	25
 Chapter 2 Description of the Gerringong-Gerroa sewage treatment plant, and chemical and bioassay analysis methodology		26
2.1	Site description: Gerringong-Gerroa sewage treatment plant	27
2.1.1	Location and history	27
2.1.2	Treatment design and technology.....	28
2.2	Sampling regime.....	29
2.2.1	Field collection of wastewater samples	29
2.3	Chemical analysis using GC-MS.....	30
2.3.1	Materials.....	30
2.3.2	Sample filtration and extraction.....	31
2.3.3	GPC clean-up	33
2.3.4	Derivatisation procedures	34
2.3.5	GC-MS detection/analysis	35
2.3.6	Quantification method, compound recovery and detection limits.....	38
2.4	In vitro bioassays.....	41
2.4.1	Two-hybrid yeast assay	41
2.4.1.1	Chemicals and reagents	41
2.4.1.2	Assay protocol.....	42
2.4.1.3	Calibration and EEq calculation	43
2.4.2	Estrogen receptor binding assay (ERBA)	43
2.4.2.1	ERBA calibration and EEq calculations	44
2.4.3	E-screen	45
2.4.3.1	E-screen calibration and EEq calculations.....	46
2.4.4	In vitro bioassay LOD/LOQ.....	46
2.5	In vivo Bioassay – vitellogenin analysis	46
2.5.1	Sourcing fish and animal care and ethics	46

2.5.2	Vitellogenin analysis in moquitofish	48
2.5.2.1	Excising and preserving the livers	48
2.5.2.2	RNA extraction	48
2.5.2.3	RNA quantification	49
2.5.2.4	Quantitative reverse transcriptase PCR	49
2.5.3	Vitellogenin analysis in rainbowfish	51
2.5.3.1	Excising and preserving the livers	51
2.5.3.2	RNA extraction, quantification and qPCR	51
 Chapter 3 The impact of variations in influent load on the efficacy of EDC removal at the Gerringong-Gerroa STP		53
3.1	Abstract.....	54
3.2	Introduction	55
3.3	Methodology.....	57
3.3.1	Study site	57
3.3.2	Sample collection periods	57
3.3.3	Extraction	58
3.3.4	Chemical analysis.....	58
3.3.5	In vitro bioassay.....	59
3.3.6	Data analysis.....	60
3.4	Results.....	60
3.4.1	Operational differences between normal and peak inflow events	60
3.4.1.1	Influent load.....	60
3.4.2	Presence and removal of EDCs	62
3.4.2.1	Influent.....	62
3.4.3	Removal of EDCs through the STP	64
3.5	In vitro bioassay responses to EDCs.....	68
3.6	Discussion	73
3.7	Conclusions	78

Chapter 4 Optimization of the two-hybrid yeast assay for the assessment of EDC potential and the ability of the assay to describe variations in EDC concentrations in complex STP matrices..... 79

4.1	Abstract.....	80
4.2	Introduction	81
4.3	Methodology.....	84
4.3.1	Assay protocol for model compounds.....	84
4.3.1.1	Standard curve, and determining the LOD and LOQ.....	85
4.3.2	Optimising sample processing for maximum response.....	86
4.3.2.1	Extraction method and fractionation of samples	86
4.3.2.2	Florisil clean-up test	87
4.3.3	Synthetic mixture testing	88
4.4	Results.....	89
4.4.1	Estradiol standard dose-response relationship and LOD/LOQs	89
4.4.2	Model compounds.....	91
4.4.3	Optimising sample processing for maximum response.....	94
4.4.3.1	Extraction method and fractionating of samples.....	94
4.4.3.2	Florisil clean-up test	95
4.4.4	Synthetic mixture tests.....	97
4.5	Discussion	100
4.6	Conclusions	103

Chapter 5 Bioassay directed assessment of ER, RAR, AhR disruption potential and genotoxicity of sewage effluent at GGSTP 104

5.1	Abstract.....	105
5.2	Introduction	106
5.3	Methodology.....	109
5.3.1	Sample collection and preparation	109
5.3.2	Sample fractionation process	110

5.3.3	Yeast assay protocol	110
5.3.4	Luminescent umu genotoxicity bioassay	111
5.4	Results.....	112
5.4.1	hER and medER bioassays	112
5.4.2	AhR bioassay.....	114
5.4.3	RAR bioassay.....	115
5.4.4	Umu test	116
5.4.5	Activity of particulate material	117
5.5	Discussion	119
5.6	Conclusions	125
Chapter 6	Partitioning of estrogenic EDC chemicals throughout the treatment process at the Gerringong-Gerroa STP	127
6.1	Abstract.....	128
6.2	Introduction	129
6.3	Methodology.....	132
6.3.1	Sample collection	132
6.3.1.1	Particulate material extraction	133
6.3.2	Data analysis.....	134
6.4	Results.....	135
6.4.1	Recovery of ¹³ C labelled analogues	135
6.4.2	Limits of detection.....	137
6.4.3	Concentration of EDCs in effluent	138
6.4.4	Partitioning between fractions.....	143
6.4.5	Percentage removal.....	144
6.5	Discussion	147
6.6	Conclusions	151

Chapter 7	Characterisation of the efficacy of an advanced tertiary sewage treatment plant to remove estrogenic activity using on-site real-time fish exposure studies.....	153
7.1	Abstract.....	154
7.2	Introduction	156
7.3	Methodology.....	158
7.3.1	Flow-through exposure system	158
7.3.2	Preparation of estradiol stock solution	159
7.3.3	Selection of test fish species.....	159
7.3.4	Fish stocking and maintenance	160
7.3.5	24 hour composite water sample collection	160
7.3.6	Chemical analysis.....	161
7.3.7	Two-hybrid yeast bioassay	161
7.3.8	Vitellogenin analysis by quantitative PCR.....	161
7.3.9	Data analysis.....	162
7.4	Results.....	163
7.4.1	Experimental conditions.....	163
7.4.2	Concentrations of EDCs in the water of the fish exposure tanks	163
7.4.3	Two-hybrid yeast response in the 24 hour composite samples	168
7.4.4	Vitellogenin up-regulation in <i>Gambusia holbrooki</i> (Girhard 1859) and <i>Melanotaenia fluviatilis</i> (Castelnau 1878).....	170
7.5	Discussion	173
7.6	Conclusions	178
Chapter 8	General discussion and conclusions.....	179
8.1	General discussion	180
8.2	Conclusions	185
8.3	Recommendations for future research.....	186

References..... 189

Appendix 1 208

List of Figures

- Figure 1.1: Representation of a cell showing the process of gene activation. Ligands (e.g. hormones such as estradiol) enter the cell and bind to the nuclear receptor (NR) coactivator forming a homodimer that then binds to the receptor responsive element triggering transcription and activation of the gene (from Janosek et al 2006).7
- Figure 2.1: Schematic of Gerringong-Gerroa STP with sampling locations. 1 – influent, 2 – post denitrifying tank, 3 –clarifier, 4 –sand filters, 5 –ozone, 6 - biologically activated carbon filters, 7 –microfiltration, 8 –UV disinfection, 9 –storage dam (Source: Boake 2006).29
- Figure 3.1: a) Influent concentration (ng/L) and b) total daily mass loading (mg/day) of compounds that were consistently above the detection limits in the influent. Black dots represent concentrations/loads measured in samples collected during normal flow conditions, white dots represent concentrations/loads measured in samples collected during peak flow conditions.64
- Figure 3.2: Two-hybrid yeast assay estradiol equivalence (EEq) for the ECx10 response of samples taken during the peak season sampling. Black bars represent the hER yeast response, while white bars are the medER yeast response. Error bars are ± 1 standard deviation (n=3). BDL= below detection limits. The detection limits were <0.6 ng/L for hER yeast assay and <1.4 ng/L for medER assay.69
- Figure 3.3: ERBA (black bars) and E-screen (white bars) EEq for the EC₅₀ (ng/L) response of samples taken on three out of the six sampling occasions during the study. Error bars represent ± 1 standard deviation (n=3). BDL= below detection limits. The detection limits were <2.9 ng/L for the ERBA assay and <0.2 ng/L for the E-screen assay.70
- Figure 3.4: Ability of chemical analysis results to predict the bioassay response when scaled for potency in the ERBA (n=10), E-screen (n=13), hER two-hybrid yeast (n=12) and medER two-hybrid yeast (n=11) bioassays. The solid line is the regression equation while the dashed line is the isometric line. The closer the regression line is to the isometric line the better the relationship between the chemistry results and the bioassay response.72
- Figure 4.1: Estradiol standard dose response curves for hER and medER two-hybrid yeast with 95% confidence interval. The solid line represents the average total/blank (T/B) chemiluminescent response for standard curve for 25 separate assays conducted over a 12 month period. The 95% confidence interval is represented by the dashed lines.90
- Figure 4.2: Regression equations for the linear portion of the average dose response relationship of estradiol in the hER two hybrid yeast assay (solid line) and the medER yeast assay (dashed line). The quantification point of T/B=10 is shown with a dotted line. Low concentration data points of the dose response relationship including those not part of the linear portion, for the hER (diamonds) and medER (squares) are also shown, however y-axis is cropped for clarity and regression also includes data points above T/B 100.91
- Figure 4.3: Model compound response for hER two-hybrid yeast with 95% confidence intervals (n=3). Solid lines are regression models using Equation 4-1.93

Figure 4.4: Model compound response for medER yeast with 95% confidence intervals (n=3). Solid lines are regression models using Equation 4-1.	93
Figure 4.5: Sample dilution curves for the spiked sandfilter samples tested with Oasis and Empore extraction protocols, with and without florisil fractionation. Only the curve for the semi-polar fraction is shown as the non-polar and polar fractions had no response in the assay. Error bars represent the 95% confidence intervals.	95
Figure 4.6: Difference in EEq (ng/L) measured for three different influent samples with and without florisil clean-up. The black bars represent direct assays without florisil clean-up and the white bars are those that have been put through florisil	96
Figure 4.7: Reduction in sample toxicity in the two-hybrid yeast assay after florisil clean-up. Diamonds represent the untreated samples and squares represent the response of the same sample with florisil clean-up prior to testing. Graphs 1a-3a are the hER response curves while graphs 1b-3b are the corresponding medER response curves. T/B = total/background chemiluminescence. EV = equivalent volume.	97
Figure 4.8: Relationship between the hER and medER two-hybrid yeast assays response to effluent samples. The solid line is the regression line ($y=1.0535x + 0.7213$, $R^2=0.832$) and the dotted line is the 1:1 isometric line.	98
Figure 4.9: Relationship for a) hER and b) medER responses with either the chemical predicted EEq of the effluent samples or to the bioassay response of synthetic simulated effluent samples. Diamonds are EEq predicted by the chemical data vs the actual bioassay response, solid line is the actual chemical data regression, triangles are synthetic bioassay responses vs the bioassay response to the real effluent samples, dashed line is synthetic sample data regression, and the dotted line is the isometric line	99
Figure 5.1: hER activity measured by the hER two-hybrid yeast assay. Results are the EC_{x10} of the sample fractions expressed as the equivalent concentration of estradiol (ng/L). Black bars represent the response in the non-polar fraction, grey bars represent the response in the semi-polar fraction and white bars represent the response of the polar fraction.	113
Figure 5.2: medER activity measured by the medER two-hybrid yeast assay. Results are the EC_{x10} of the sample fractions expressed as the equivalent concentration of estradiol (ng/L). Black bars represent the response in the non-polar fraction, grey bars represent the response in the semi-polar fraction and white bars represent the response of the polar fraction.	113
Figure 5.3: AhR activity measured by the AhR two-hybrid yeast assay. Results are the EC_{x10} of the sample fractions expressed as the concentration equivalent to β -naphthoflavone. Black bars represent the response in the non-polar fraction, grey bars represent the response in the semi-polar fraction and white bars represent the response in the polar fraction.	115
Figure 5.4: RAR activity measured by the RAR two-hybrid yeast assay. Results are the EC_{x10} of the sample fractions expressed as the equivalent concentration of all-trans retinoic acid (ng/L). Black bars represent the response in the non-polar fraction, grey bars represent the response in the semi-polar fraction and white bars represent the response in the polar fraction.	116
Figure 5.5: Genotoxicity as measured by expression of the umu gene in <i>S. typhimurium</i> TL210. Sample responses are expressed as the equivalent concentration in either 4-nitroquinoline-1-oxide (4NQO) in ng/L for the samples not incubated with	

metabolic activator (-S9) and as benzo(a)pyrene (BaP) in ng/L for samples incubated with metabolic activator (+S9). Analysis was conducted on the September 2007 samples.117

Figure 5.6: a) AhR and b) RAR activity from particulate fraction of sewage samples taken from different stages of treatment. Particulate samples were for the raw, post-DN and dam (dam1) were collected on the 7 September 2007 and samples from the sandfilter, O3/BAC, microfiltration and dam (dam2) were collected on the 25 January 2007. Black bars represent the response in the non-polar fraction, grey bars represent the response in the semi-polar fraction and white bars represent the response of the polar fraction.118

Figure 6.1: Average concentration of steroid estrogens a) E₂, b) E₁, c) E₃ and 4) the anti-microbial triclosan in the aqueous and particulate phases of effluent samples from different stages of wastewater treatment at GGSTP. Black bars represent the aqueous concentration while the grey bars are the particulate concentrations. Error bars are the standard deviation. n=6 for each grouping, though the BAC, microfiltration and UV particulate fractions were analysed as two composite samples containing 3 individual samples each.141

Figure 6.2: Average concentrations of a) BPA, b) technical NP, c) 4tOP and 4) 4tAP in the aqueous and particulate phases of effluent samples from different stages of wastewater treatment at GGSTP. Black bars represent the aqueous concentration while the grey bars are the particulate concentrations. Error bars are the standard deviation. n=6 for each grouping, though the BAC, microfiltration and UV particulate fractions were analysed as two composite samples containing 3 individual samples each.142

Figure 6.3: Partitioning of EDCs between the aqueous and particulate phases a) in the post-DN tank, n=6 for BPA, triclosan and TNP, n=3 for E₁ and E₃, n=1 for E₂ and b) from the clarifier/sandfilter/ozone treatment stages. n= 2 for E₂, n=8 for triclosan, n=1 for 4nNP, n=10 for 4tOP, n=13 for TNP, n=9 for BPA. Black bars represent aqueous fraction and grey bars represent particulate fraction. Values on bars represent the standard deviation.143

Figure 6.4: Average percentage of EDCs remaining in the aqueous and particulate fractions after each treatment stage and the implied removal based on the concentration reduction for a) E₁ b) E₃, c) triclosan d) BPA, e) TNP and f) 4tOP. Black bars represent the concentration remaining in the aqueous phase, grey bars the particulate phase, white bars the apparent removal and cross-hatched bars represent the maximum additional removal if concentrations below detection limits were actually complete removal to nil.146

Figure 7.1: Average ± standard deviation (n=6) concentrations of EDCs measured in 24 hour composite samples taken from a representative tank for each treatment of the fish flow-through exposure experiments a) May 2008, b) November 2008 and c) February 2009. Error bars represent the standard deviation. N.D. – not detectable167

Figure 7.2: Average ± standard deviation (n=6) of estrogenic response in the hER (black bars) and medER (white bars) two-hybrid yeast assay for the 24-hour composite samples taken in a) the May 2008, b) the November 2008 and c) the February 2009 fish flow-through exposure experiment. Columns with the same letter above them are not significantly different (p>0.05). Lower case letters are for hER and upper case for medER responses.169

Figure 7.3: Vitellogenin upregulation relative to the average vitellogenin upregulation in fish from the control tanks for the a) May 2008 exposure experiment and b) the November 2008 exposure experiment for *G. holbrooki* and c) the February 2009 exposure experiment for *M. fluviatilis*. Error bars represent the standard error of the mean (SEM). * indicates <0.05 and *** indicates <0.001 for the significant difference to the control by Dunnett test.172

List of Tables

Table 1.1: Emerging contaminants in wastewater that could cause ED (from Chang et al. 2009).....	9
Table 1.2: Hormone concentrations in raw sewage influent or primary treated effluent. All concentrations reported in ng/L	11
Table 1.3: Alkylphenol and Bisphenol concentrations in raw sewage influent or primary treated effluent. All concentrations reported in ng/L.....	12
Table 1.4: Examples of in vitro bioassays for the screening of estrogenic ED potential.	22
Table 1.5: Summary of reported concentrations of xenoestrogens exposure resulting in the induction of vitellogenin production in male fish (sourced from Brown et al. 2001).....	24
Table 2.1: NSW DECCW effluent emission license requirements	28
Table 2.2: TFAA derivatised compounds retention times and acquisition ions.....	36
Table 2.3: MSTFA derivatised compounds retention time and acquisition ions	37
Table 2.4: Limits of detection for TFAA derivatised samples (concentration within wastewater sample based on 1.0 L for influent, post-DN, field and MSTFA blanks and 2.0 L of sample for ozone and a derivitisation volume of 250 μ L).....	39
Table 2.5: Limits of detection for MSTFA derivatised samples (concentration within wastewater sample based on 1.0 L for influent, post-DN, field and MSTFA blanks and 2.0 L of sample for ozone and a derivitisation volume of 200 μ L).....	40
Table 2.6: Primer sequences and melting temperatures (T_m) of the vitellogenin target and 18S housekeeping genes. Lower case lettering indicates base pairs involved in the hairpin structure quenching the LUX fluorophore.	50
Table 3.1: Average daily influent load parameters based on online monitors at the GGSTP. Hydraulic retention time (HRT) estimates were calculated using a standard formula for the STP design.	61
Table 3.2: General influent/effluent quality parameters. Results are based on the average of two samples collected and analysed during the sampling periods as part of the STP's routine monitoring by Sydney Water Corporation. COD - chemical oxygen demand, TSS – total suspended solids, total N – total nitrogen, total P – total phosphorus, CBOD – chemical biological oxygen demand.....	62
Table 3.3: Steroidal estrogen concentrations. Concentration (ng/L) range of steroid estrogens under normal and peak flow conditions (n=3 samples for each sampling period).	66
Table 3.4: Synthetic phenolic compounds. Concentration range (ng/L) of phenols and triclosan during normal and peak flow conditions.	67
Table 3.5: Limits of detection (LOD) and limits of quantification (LOQ) ranges encountered when conducting the different bioassays in ng/L. The LOD and LOQ varied each time the assay was run due to slight differences in background signal (affecting the LOD), and standard EC_{50} response (affecting the LOQ).	68
Table 3.6: Potencies of the some of the expected EDCs in wastewater in the ERBA, E-screen and hER and medER two-hybrid yeast assays. Potencies are relative to 17 β -	

estradiol and are calculated by dividing the EC ₅₀ of 17β-estradiol by the EC ₅₀ of the test compound (or EC _{x10} for the two-hybrid yeast assay). NA – not assessed.	71
Table 4.1: Two-hybrid yeast assay EC _{x10} ± standard deviation (n=18) and the equivalent LOQ for a 1.0 L sample concentrated to 100 μL before assaying. Sample LOD based on blank + 3 standard deviations.....	90
Table 4.2: Potency of estrogenic EDCs relative to estradiol in the two-hybrid yeast test (E ₂ EC _{x10} / compound EC _{x10}) based on molar response ratios.....	92
Table 4.3: EC _{x10} averages ± standard deviation (n=3) for replicate samples extracted using either Oasis column or Empore disk, with and without florisil fractionation. The samples are 1.0 L sandfilter process water spiked with 28 ng of estradiol. Responses are reported in EEq (ng/L). “RSD” is the relative standard deviation.	94
Table 4.4: Average percentage (%) of the real bioassay sample response measured in the synthetic sample simulations tested with the two hybrid yeast assay and predicted EEq calculated from the chemical data.	99
Table 5.1: Relative percentages of the total AhR or RAR yeast response between the particulate and aqueous phases for the total response of the three solvent fractions. n.a. = not applicable as both phases were below the detection limits.....	119
Table 6.1: Ranges of Log K _{OC} ^a (Campbell et al. 2006) and Log K _{OW} ^b (Yoon et al. 2007), ^c (Ahel & Giger 1993) found for some estrogenic EDCs.	131
Table 6.2: Percentage recovery ± standard deviation of ¹³ C labelled concentrations from aqueous samples. n=6, difference to the recovery measured in the blank was tested using Dunnett’s test with unequal variance correction, significance is represented by *p<0.05, **p<0.01, ***p<0.001	136
Table 6.3: Percentage recovery ± standard deviation of ¹³ C labelled analogues from particulate samples. n=6 for post-DN, clarifier, sandfilter and dam samples. Ozone, microfiltration, final and blank are average recoveries of two composite samples. BAC is only one composite sample. Difference to the recovery measured in the blank was tested using Dunnett’s test with unequal variance correction, significance is represented by ***p<0.001	137
Table 6.4: Detection limits for the EDCs analysed for in different sample matrices. Concentrations are in ng/L	138
Table 7.1: Water quality conditions measured during on-site flow-through fish exposure experiments. Averages are from daily measurements of all tanks during the experimental period.....	165
Table 7.2: Average ± standard deviation of recovery (%) for different sample matrices and different isotopic labelled EDC analogues. The ¹³ C labelled compounds were used as the surrogate standard in the November 2008 and February 2009 samples, while the deuterated (d4 or d16) compounds were used for the May 2008 samples. ...	166
Table 7.3: Average cycle thresholds (Ct) for of vitellogenin and the housekeeping gene 18S and the average normalised vtg up-regulation (ΔCt) and change in Ct relative to the normalised vtg up-regulation in control tank fish (-ΔΔCt). Error (±) is the standard deviation and brackets contain number of quantifications for treatments that did not have quantifiable vtg for all fish (n).....	171

Abstract

In Australia, due to increased uncertainties over security of water supply because of unpredictable drought and flood cycles, alternative water sources are being investigated for commercial, agricultural, industrial and domestic supply, including the option of reusing treated sewage effluents. However, sewage effluent is a known source of estrogenic endocrine disrupting chemicals (EDCs) in the environment. Exposure to sewage effluents containing steroid estrogens and xenoestrogens can cause developmental and behavioural reproductive abnormalities in fish and other aquatic animals. As such, risk of endocrine disruption is one of the water quality issues that needs to be evaluated when assessing the appropriate level of treatment required for reuse applications. The Gerringong-Gerroa sewage treatment plant (GGSTP), currently employs advanced tertiary treatment technology to treat domestic sewage from two small coastal towns (Gerringong and Gerroa), which receive large seasonal influxes of holiday makers. In this study, the efficacy of the treatment at the GGSTP in removing estrogenically active chemicals was assessed using a multi-tiered assessment approach, incorporating chemical analysis, *in vitro* bioassays and *in vivo* fish exposure studies.

The raw sewage influent was found to contain steroidal estrogens; 17β -estradiol (E_2), estrone (E_1) and estriol (E_3) as well as synthetic phenolic xenoestrogens; 4-tert-octylphenol, Bisphenol A and technical nonylphenol at concentrations commonly found in sewage influents. The influent also displayed high levels of activity in the two-hybrid yeast *in vitro* bioassay. However, the final effluent had no detectable concentrations of steroidal estrogens, no estrogenic activity in the two-hybrid yeast assay and only infrequent occurrence of low concentrations of synthetic phenols. Biodegradation by activated sludge treatment provided significant, but incomplete removal of measured EDCs and estrogenic activity, with the in-line combination of ozone oxidation and biologically activated carbon filtration reducing the remaining estrogenic activity to undetectable levels. EDCs in both the dissolved and particulate phases of the effluent were removed by the treatment process and the efficacy of treatment was not compromised by increases in influent flow during the peak holiday seasons. Treatment of the effluent at the GGSTP was also successful at reducing retinoic acid receptor

(RAR) activity and genotoxicity to below detection limits and greatly reducing arylhydrocarbon receptor (AhR) activity.

On-site real-time exposure tests using the mosquitofish (*Gambusia holbrooki*) and rainbowfish (*Melanotaenia fluviatilis*) demonstrated that the final effluent did not elicit up-regulation of vitellogenin, a well known biomarker of exposure to estrogenic EDCs. Despite the presence of residual concentrations of E₁ and the *in vitro* activity in effluent after being processed through activated sludge treatment, clarification and sandfiltration, vitellogenin up-regulation was not detected in fish exposed to this partially treated effluent. Overall, the results provide evidence that the application of advanced tertiary treatment technology to domestic sewage can produce a final effluent that is unlikely to pose an endocrine disruption risk to the aquatic biota.

Abbreviations

4nNP	4- <i>n</i> -nonylphenol
4nOP	4- <i>n</i> -octylphenol
4NQO	4-nitroquinoline-N-oxide
4tAP	4- <i>tert</i> -amylphenol
4tOP	4- <i>tert</i> -octylphenol
AhR	Arylhydrocarbon receptor
AP	Alkylphenol
APEO/APE	Alkylphenol polyethoxylates
AS	Activated sludge
atRA	<i>all-trans</i> retinoic acid
BAC	Biological activated carbon
BaP	Benzo[a]pyrene
BDL	Below detection limits
BPA	Bisphenol A
COA	Chemical advanced oxidants
DAFF	Dissolved air floatation filtration
DCM	Dichloromethane
DE	Diethylether
DECCW	Department of Environment Climate Change and Water
DMSO	Dimethylsulfoxide
DN	Denitrification
DNA	Dioxyribose nucleic acid
E ₁	Estrone
E ₂	17β-estradiol
E ₃	Estriol
ED	Endocrine disruption
EDC	endocrine disrupting chemical/compound
EE ₂	17α-ethynylestradiol
EEq	Estradiol equivalent (concentration)
EPA	Environmental Protection Agency
ER	Estrogen receptor
ERBA	Estrogen receptor binding assay
ERE	Estrogen response element
EROD	Ethoxyresorufin-O-deethylase
ESI	Electrospray ionisation
GAC	Granular activated carbon
GC-MS	Gas chromatography mass spectrometry
GC-MS-MS	Gas chromatography tandem mass spectrometry
GGSTP	Gerringong-Gerroa Sewage Treatment Plant
GPC	Gel permeation chromatography

H ₂ SO ₄	Sulphuric acid
HAH	Halogenated aromatic hydrocarbon
hER	Human estrogen receptor
HRT	Hydraulic retention time
ISTD	Internal standard
Kow	Octanol-water partitioning coefficient
LC-MS-MS	Liquid chromatography tandem mass spectrometry
LOD	Limits of detection
LOQ	Limits of quantification
MCF-7	Michigan Cancer Foundation -7
medER	Medaka (<i>Oryzias latipes</i>) estrogen receptor
MeOH	Methanol
MLSS	Mixed liquor suspended solids
mRNA	messenger ribonucleic acid
MSTFA	n-methyl-N-(trimethyl-silyl) trifluoroacetamide
NP	Nonylphenol
NPE	Nonylphenol polyethoxylates
NSW	New South Wales
O ₃	Ozone
OP	Octylphenol
OPE	Octylphenol polyethoxylates
PAC	Particulate activated carbon
PAH	Polycyclic aromatic hydrocarbon
PCB	Polychlorinated biphenyl
PCR	Polymerase chain reaction
PNEC	Proposed no effect concentration
PPCPs	Pharmaceuticals and personal care products
QLD	Queensland
qPCR	Quantitative polymerase chain reaction
RAR	Retinoic acid receptor
RNA	Ribonucleic acid
RO	Reverse osmosis
RT-PCR	Reverse-transcriptase polymerase chain reaction
SPE	Solid phase extraction
SRT	Solids retention time
STP	Sewage treatment plant
TCDD	2,3,7,8-tetrachlorodibenzo-p-dioxin
TFAA	Trifluoroacetic anhydride
TIE	Toxicity Identification Evaluation
T _m	Melting temperature
TNP	Technical nonylphenol
UV	Ultraviolet light
Vtg	Vitellogenin

YES
 αE_2
 β -NP

Yeast estrogen screen
17 α -estradiol
 β -naphthoflavone