Tai Chi and Stress: A Randomised Controlled Trial and Chinese Medicine Pattern Diagnostics

Shuai Zheng

A thesis submitted as partial requirement for the degree of Doctor of Philosophy (Science)

Faculty of Science

University of Technology, Sydney

December 2014

Certificate of Authorship/Originality

I certify that the work in this thesis has not previously been submitted for a degree nor has it been submitted as part of requirements for a degree except as fully acknowledged within the text.

I also certify that the thesis has been written by me. Any help that I have received in my research work and the preparation of the thesis itself has been acknowledged. In addition, I certify that all information sources and literature used are indicated in the thesis.

Production Note: Signature removed prior to publication.

Shuai Zheng

Abstract

Stress is a growing problem in modern society and in severe cases could potentially lead to hospitalisation. The ancient Chinese mind body exercise of *Tai Chi* (TC) is practiced worldwide by millions of people daily and is often accredited with a plethora of health benefits, including reduction of stress. There has been a growing interest in the scientific community to evaluate the efficacy of TC practice for a range of diseases and conditions, in particular the effects of TC in the improvement of psychological and mental health. However the term "stress" itself is a common diagnosis used by both patients and practitioners alike to describe a "condition" generally regarded as subjective in nature and as such each individual will likely present with varying somatic or cognitive signs and symptoms. Currently there are no definitive diagnosis or signs and symptoms for "stress" for both biomedicine and Chinese Medicine (CM). This thesis evaluates the efficacy of TC as an intervention for stress through a randomised controlled trial (RCT) and the use of a questionnaire to identify the CM diagnostic patterns associated with stress

The RCT involved fifty participants who were randomly allocated into one of three groups; TC (n=17), exercise (n=16) or a wait list control group (WL) (n=16). Both TC and exercise groups were required to complete 5 hours per week of either TC or exercise for 12 weeks (total of 3600 minutes), whilst WL maintained their normal lifestyle. State Trait Anxiety Inventory (STAI) which assesses both state anxiety and trait anxiety was used as the primary outcome measure. Secondary outcome measures were the Perceived Stress Scale 14 (PSS14), blood pressure, heart rate variability, visual analogue scale and short form 36. Data were collected at baseline, midway at week 6 and at the completion of the trial at week 12. A two-way ANOVA with repeated measures followed by Bonferroni's post-hoc test was used for statistical analysis. The stress questionnaire was constructed using the signs and symptoms for General Anxiety Disorder and commonly

reported signs and symptoms for stress were cross referenced against TCM textbooks to relate each sign and symptom to possible patterns (*zheng*) and then cross referenced again against the signs and symptoms list to form two gender specific questionnaires. Pattern identification measurement was based on percentage of signs and symptoms present against possible number of signs and symptom per pattern for each gender.

Results for the RCT showed that there were significant improvements from baseline for both TC and exercise groups for the outcomes of STAI, PSS14, VAS as well as mental health and vitality domains of the SF36. Furthermore there were significant differences between groups for TC and the wait list control group for both state and trait anxiety as well as the mental health domain. The stress questionnaire results indicate that the top three pattern for both genders were Heart Qi deficiency, Heart Blood deficiency and Liver Blood deficiency. This result is different to the commonly accepted idea that stress is associated with Liver Qi stasis, however despite the reliability of the instrument a larger cohort size will be needed to ascertain the validity of these findings.

It is hoped that the results from these two studies will be incorporated into future research in both TC and stress diagnosis. The findings from the RCT revealed that TC effectively reduces stress levels in healthy individuals as early as week 6 and may provide a safer and less strenuous therapeutic alternative to exercise. Whilst the results from the stress questionnaire will hopefully help CM clinicians with their understanding of stress related symptoms the choosing the correct treatment principles for their patients.

List of referred papers, conference presentations and posters arising from the research

Publications

ZHENG, S., Lal, S.,Meier, P., Sibbritt, D., Zaslawski, C. 2014. "Protocol: The Effect of 12 Weeks of Tai Chi Practice on Anxiety in Healthy but Stressed People Compared to Exercise and Wait-list Comparison Groups: A Randomized Controlled Trial." *Journal of Acupuncture and Meridian Studies*, 7 (3), 159-165

ZHENG, S., Meier, P., Zaslawski, C. 2014. "Development of a novel questionnaire for the Traditional Chinese Medicine pattern diagnosis of stress", Poster Abstract, *Journal of Integrative Medicine*, 12 (3), 297.

ZHENG, S., Lal, S.,Meier, P., Sibbritt, D., Zaslawski, C. 2014. "The effects of twelve weeks of Tai Chi practice on anxiety in stressed but healthy people compared to exercise and wait-list groups: a randomised controlled trial", Poster Abstract, *Journal of Integrative Medicine*, 12 (3), 297-298.

ZHENG, S., Meier, P., Zaslawski, P. 2013. "WFAS 2013: Selected Conference Abstracts - Development of a novel questionnaire for the Traditional Chinese Medicine pattern diagnosis of stress." *Australian Journal of Acupuncture and Chinese Medicine*, 8 (2), 25-32.

Oral Presentations

Speaker at an international conference

ZHENG, S. 2013. "Development of a novel questionnaire for the Traditional Chinese Medicine pattern diagnosis of stress." *World Conference on Acupuncture World Federation of Acupuncture and Moxibustion Societies*, 2-4th November 2013, Sydney Australia...

Invited speaker at an international conference

ZHENG, S. 2012. "Tai Chi and Stress: A randomised controlled pilot study." *International Scientific Acupuncture and Meridian Symposium 2012*, Sydney, 5-7th

October 2012.

Poster Presentations

ZHENG, S., Meier, P., Zaslawski, C. 2014. "Development of a novel questionnaire for the Traditional Chinese Medicine pattern diagnosis of stress". Society for Acupuncture Research and the Chinese Association of Acupuncture and Moxibustion Conference, Beijing, China. 30th May to 1st June 2014.

ZHENG, S., Lal, S., Meier, P., Sibbritt, D., Zaslawski, C. 2014. "The effects of twelve weeks of Tai Chi practice on anxiety in stressed but healthy people compared to exercise and wait-list groups: a randomised controlled trial". Society for Acupuncture Research and the Chinese Association of Acupuncture and Moxibustion Conference, Beijing, China. 30th May to 1st June 2014.

Zheng, S., Lal, S., Meier, P., Zaslawski, C. 2013. "The effects of six weeks of Tai chi practice on anxiety in healthy but stressed individuals compared to an exercise only comparison a randomised controlled trial: Pilot Study. Society for Acupuncture Research Conference, Ann Arbor, Michigan, USA. 18th to 21st April 2012.

Zheng, S., Meier, P., Zaslawski, C. 2011. "The effects of six weeks of Tai chi practice on anxiety in healthy but stressed individuals compared to an exercise only comparison a randomised controlled trial: Pilot Study. International Forum & Exhibition on Integrative Medicine, Sydney, Australia. 21st to 23rd October 2011.

Acknowledgements

This thesis could not have been completed without the contributions and support of so many and my sincerest gratitude to all those who have helped me on this journey. I would like to thank some of them in this acknowledgement.

I would like to sincerely thank all the participants who volunteered for the study, both those who were involved in the randomised controlled trial as well as those who completed the stress questionnaire. None of this would be possible without their generosity and immeasurable contributions to broaden our knowledge and understanding.

My thanks go especially to my principal supervisor, Associate Professor Chris Zaslawski, whose endless patience and commitment to my study gave it form and structure. His wisdom and guidance has helped me every step of the way, from protocol design and recruitment to the final edits that has polished this thesis. Chris was not only my PhD supervisor but also a great mentor and I am proud to say that I was his student.

My co-supervisor Associate Professor Peter Meier, I thank you for your support during my PhD journey. Associate Professor Sara Lal for her expertise in neuroscience and heart electrophysiology, as well as allowing me to share her laboratory to collect the physiological data for the clinical trial component of my PhD.

My sincere thanks to Professor David Sibbritt from the Faculty of Health whose advice and expertise in statistics have helped me immensely in this undertaking. My thanks to my colleague Christine Berle, whose humour and wisdom have kept me sane through my PhD journey. I would also like to thank the Chinese Lions Club of Sydney for their generous scholarship which helped support me during my years as a PhD student.

I would also like to thank all of my family for their patience and continuous support during these four years and especially to my beautiful fiancée Dr Christine Kim, who kept me going with her advice, expertise, prayers and immeasurable love.

Finally, I would like to thank God, to whom deserves all the glory!

Contents

Abstract	iii
List of referred papers, conference presentations and posters research	_
Contents	X
List of Figures	xiv
List of Tables	XV
Chapter I: Introduction and aims	17
1.1 Background to the studies	17
1.1.1 Concept of Stress	18
1.2 Study Aims.	19
1.2.1 Tai Chi Randomised Controlled Trial (RCT)	19
1.2.2 Stress questionnaire	20
1.3 Format of thesis.	20
Chapter II: Literature review	23
2.1. Introduction	23
2.2 Theories of Stress	23
2.3 Stress hormones	26
2.4 Current Treatments for Stress	27
2.5 History and development of Tai Chi	29
2.6 Tai Chi Research	30
2.7 Tai Chi Research in Psychological Health	32
Chapter III: Methods – Tai Chi RCT	41
3.1 Trial design and protocol	41

3.2 Participant Recruitment	41
3.3 Inclusion and Exclusion Criteria.	42
3.4 Data Collection Location	43
3.5 Ethical Considerations	44
3.6 Intervention	44
3.6.1 Tai Chi Rationale	44
3.6.2 Tai Chi Instruction Method	45
3.6.3 Tai Chi Dosage	46
3.6.4 Active control	46
3.6.5 Non active control	47
3.7 Adverse events and discontinuation from the study	47
3.8 Outcome Measures	48
3.8.1 State Trait Anxiety Inventory	48
3.8.2 Perceived Stress Scale 14	49
3.8.3 Heart Rate Variability	49
3.8.4 Blood Pressure	50
3.8.5 Short Form 36 Health Survey	51
3.8.6 Visual Analogue Scale	51
3.9 Statistical Methods	51
3.9.1 Sample Size and Randomisation	51
3.9.2 Data Management and Statistical Analysis	52
3.10 Method Discussion	52
Chapter IV: Results – Tai Chi RCT	55
4.1 Primary Outcome Measures – State Trait Anxiety Inventory (STAI)	
4 1 1 State Anxiety (Y-1)	57

4.1.2 Trait Anxiety (Y-2)			
4.1.3 State and Trait Anxiety	59		
4.2 Secondary Outcome Measures	60		
4.2.1 Perceived Stress Scale 14	60		
4.2.2 Visual Analogue Scale (VAS)	61		
4.2.3 Blood Pressure	63		
4.2.4 Short Form 36 Health Survey (SF36)	66		
4.2.5 Heart Rate Variability	77		
4.3 Summary	81		
Chapter V: Discussion – Tai Chi RCT	83		
5.1 State Anxiety	83		
5.2 Trait Anxiety	86		
5.3 Perceived Stress Scale 14	87		
5.4 Visual Analogue Scale	88		
5.5 Blood pressure	88		
5.6 Short Form 36	89		
5.7 Heart Rate variability	92		
5.8 Limitations and Future Directions	94		
Chapter VI: Methods – Stress Questionnaire	96		
6.1 Instrument Design	96		
6.1.1 Initial Biomedical Sign and Symptom Selection	96		
6.1.2 Chinese Medicine Diagnostic Pattern Differentiation and Sign and Sympton	n		
Integration	97		
6.2 Ethical Considerations	100		
6.3 Intra-rater Reliability	100		

6.4 Data Collection	100
Chapter VII: Results – Stress Questionnaire	101
7.1 Intra Rater Reliability	101
7.2 Internal consistency	101
7.3 Pattern Differentiation	101
7.4 Symptoms	105
VIII: Discussion – Stress Questionnaire	107
8.1 Reliability	107
8.2 Pattern and Symptoms	107
8.3 Limitations	110
Chapter IX Overview and Recommendations	113
References	115
Appendices	123
Appendix 1	123
Appendix 2	128
Appendix 3	137
Appendix 4	139
Appendix 5	154
Appendix 6	157

List of Figures

Figure 1: Flow chart of recruitment and completion of the outcome measures	43
Figure 2: State anxiety score in Tai Chi, exercise and wait list	57
Figure 3: Trait anxiety score in Tai Chi, exercise and wait list	59
Figure 4: Perceived stress scale 14 score in Tai Chi, exercise and wait list	61
Figure 5: Visual analogue scale 14 score in Tai Chi, exercise and wait list	62
Figure 6: Mean arterial pressure in Tai Chi, exercise and wait list	64
Figure 7: Systolic blood pressure in Tai Chi, exercise and wait list	65
Figure 8: Diastolic blood pressure in Tai Chi, exercise and wait list	66
Figure 9: Physical functioning scores in Tai Chi, exercise and wait list	67
Figure 10: Role – physical scores in Tai Chi, exercise and wait list	68
Figure 11: Bodily pain scores in Tai Chi, exercise and wait list	69
Figure 12: General health scores in Tai Chi, exercise and wait list	70
Figure 13: Vitality scores in Tai Chi, exercise and wait list	71
Figure 14: Social functioning scores in Tai Chi, exercise and wait list	73
Figure 15: Role – emotional scores in Tai Chi, exercise and wait list	74
Figure 16: Mental health scores in Tai Chi, exercise and wait list	76
Figure 17: Low frequency scores in Tai Chi, exercise and wait list	78
Figure 18: High frequency scores in Tai Chi, exercise and wait list	79
Figure 19: Low frequency: high frequency ratio in Tai Chi, exercise and wait list	80
Figure 20: Total power in Tai Chi, exercise and wait list	81
Figure 21: Flow chart of stress questionnaire instrument design	99
Figure 22: Mean percentage of symptoms per pattern in males and females	103

List of Tables

Table 1: Mean Baseline Parameters For Participants in the Tai Chi RCT	56
Table 2: Symptoms Commonly Associated With Stress	97
Table 3: Mean Percentage Of Symptoms Per Pattern in Males And Females	102
Table 4: Frequency Of Most Common Patterns	105