

**Multivariate and
multiway analysis of
hyperspectral and
fluorescence
landscape data**

by

Helen Rutlidge

A thesis submitted for the

Degree of Doctor of Philosophy (Science)

University of Technology, Sydney

March, 2011

Dedicated to Michael Coleman and Peter Rutledge

"For some life lasts a short while, but the memories it holds lasts forever."

Laura Swenson

Certificate of authorship and originality

I certify that the work in this thesis has not previously been submitted for a degree nor has it been submitted as part of the requirements for a degree except as fully acknowledged within the text.

I also certify that the thesis has been written by me. Any help that I have received in my research work and the preparation of the thesis itself has been acknowledged. In addition, I certify that all the information sources and literature used are indicated in the thesis.

Helen Rutledge

Production Note:
Signature removed prior to publication.

31/03/2011

Acknowledgements

Whilst this has been my project for the last five years there have a number of people who have contributed to the success of it. Firstly to Dr Brian Reedy, my supervisor, who first started me on this journey. Your guidance and encouragement have enhanced this process for me and made the project what it is.

I would like to thank Professor Peter Ralph for his suggestion and assistance with the algal cultures in this project. Thank you to Vinod Kumar and Penelope Ajani for their technical assistance with the algal cultures.

I thank Dr Dominic Hare and Christine Austin for all of their assistance with laser ablation inductively coupled plasma. Both of you were always willing to answer any questions I had and provide assistance when required.

I would also like to Fiona Burger, Jonathan Mak and Anne Rutledge for collecting soil samples for me.

Thanks also go to everyone in Office 4.60 and other fellow students (past and present) who I have shared this journey with. You have always been willing to celebrate the good times and find the silver lining in the bad. All of you have made being at uni enjoyable.

I would like to thank my family and friends for their support and understanding whilst I have been at university. Thank you also for making sure I had a life outside my PhD.

Table of contents

CHAPTER 1: INTRODUCTION	2
1.1 HYPERSPECTRAL IMAGING.....	2
1.1.1 <i>UV-visible imaging</i>	4
1.1.1.1 Fluorescence imaging.....	6
1.1.2 <i>Infrared imaging techniques</i>	6
1.1.2.1 Near infrared imaging	7
1.1.2.2 Mid-infrared mapping and imaging.....	9
1.1.3 <i>Raman mapping and imaging</i>	12
1.1.4 <i>Molecular mass spectrometry imaging</i>	14
1.1.5 <i>Elemental mapping</i>	16
1.1.5.1 X-ray mapping techniques	16
1.1.5.2 Laser ablation inductively coupled mass spectroscopy.....	17
1.2 ANALYSIS OF HYPERSPECTRAL IMAGES	18
1.2.1 <i>Preprocessing methods</i>	19
1.2.1.1 Savitzky-Golay derivatives	20
1.2.1.2 Mean-centring	20
1.2.1.3 Variance scaling	20
1.2.1.4 Autoscaling.....	20
1.2.1.5 Vector normalisation.....	21
1.2.2 <i>Clustering techniques</i>	21
1.2.2.1 Distance measures.....	21
1.2.2.2 Hierarchical cluster analysis.....	22
1.2.2.3 Fuzzy C-means clustering.....	24
1.2.2.4 Self-organising feature map.....	25
1.2.3 <i>“Factor analysis” techniques</i>	26

1.2.3.1	Principal component analysis	27
1.2.3.2	Partial least squares analysis	28
1.2.4	<i>End member techniques</i>	30
1.2.4.1	Spectral angle mapper	30
1.2.4.2	Spectral feature fitting	31
1.2.4.3	Mixture tuned matched filter	32
1.2.5	<i>Multiway techniques</i>	33
1.2.5.1	PARAFAC.....	33
1.2.5.2	Tucker models.....	34
1.2.5.3	Application of multiway techniques on hyperspectral images	34
1.2.6	<i>Dimension reduction strategies</i>	35
1.3	PROJECT AIMS.....	35
CHAPTER 2:	CLASSIFICATION OF HETEROGENEOUS SOLIDS USING INFRARED HYPERSPECTRAL	
IMAGING	37	
2.1	INTRODUCTION.....	37
2.1.1	<i>Hyperspectral imaging</i>	37
2.1.2	<i>Dimension reduction strategies</i>	38
2.1.2.1	PCA on PC1.....	38
2.1.2.2	ASMULIM	39
2.1.2.3	MOLMAP	39
2.1.3	<i>Classification techniques</i>	39
2.1.3.1	Discriminant analysis.....	40
2.1.3.2	Soft independent modelling of class analogy.....	40
2.1.4	<i>Aims</i>	41
2.2	EXPERIMENTAL AND DATA ANALYSIS METHODS	42
2.2.1	<i>Sample preparation</i>	42
2.2.2	<i>Instrumental</i>	43
2.2.3	<i>Data preprocessing</i>	44

2.2.4	<i>Data analysis</i>	44
2.2.4.1	ASD and MIQ methods	44
2.2.4.2	Average and median methods	46
2.2.4.3	Clustering methods	46
2.2.4.4	PCA on PC1	47
2.2.4.5	ASMULIM	48
2.2.4.6	Endmember techniques	48
2.3	RESULTS AND DISCUSSION.....	49
2.3.1	<i>Choice of compounds</i>	49
2.3.2	<i>Sampling considerations</i>	49
2.3.3	<i>Particle size vs pixel size</i>	50
2.3.4	<i>Comparison of classification approaches</i>	51
2.3.5	<i>Endmember classification results</i>	56
2.4	CONCLUSIONS	57
CHAPTER 3: CLASSIFICATION OF SUBURBAN SOIL SAMPLES USING HYPERSPECTRAL IMAGING ...		59
3.1	INTRODUCTION.....	59
3.1.1	<i>Soil</i>	59
3.1.1.1	Forensic analysis of soil.....	59
3.1.1.2	Mid-infrared analysis of soil.....	60
3.1.1.3	Elemental analysis of soil	60
3.1.2	<i>LA-ICP-MS</i>	62
3.1.2.1	LA-ICP-MS calibration.....	62
3.1.3	<i>Overall aims</i>	63
3.2	EXPERIMENTAL AND DATA ANALYSIS METHODS	64
3.2.1	<i>Preliminary study</i>	64
3.2.1.1	Sample preparation	64
3.2.1.2	Infrared imaging	65

3.2.1.3	LA-ICP-MS elemental imaging.....	65
3.2.1.4	Data analysis	67
3.2.2	<i>Final study</i>	67
3.2.2.1	Sample preparation	67
3.2.2.2	Element selection	68
3.2.2.3	LA-ICP-MS elemental imaging.....	69
3.2.2.4	Data analysis	70
3.3	RESULTS AND DISCUSSION.....	71
3.3.1	<i>LA-ICP-MS normalisation</i>	71
3.3.2	<i>Preliminary soil sample set</i>	72
3.3.3	<i>Final soil sample set</i>	76
3.3.3.1	Element selection	76
3.3.3.2	Classification results	78
3.3.3.3	Comparison of different imaging approaches.....	81
3.4	CONCLUSIONS	82
CHAPTER 4: PIGMENT ANALYSIS IN ALGAE USING FLUORESCENT LANDSCAPES AND PARAFAC ...		85
4.1	INTRODUCTION.....	85
4.1.1	<i>Algae</i>	85
4.1.1.1	Pigment analysis	85
4.1.1.2	Algal fluorescence.....	86
4.1.1.3	Adaptation of algae to light	87
4.1.2	<i>Fluorescent landscapes</i>	88
4.1.3	<i>PARAFAC</i>	89
4.1.4	<i>Aims</i>	89
4.2	EXPERIMENTAL AND DATA ANALYSIS METHODS	90
4.2.1	<i>Samples</i>	90
4.2.2	<i>Light table</i>	90

4.2.3	<i>Preliminary study</i>	91
4.2.4	<i>Final study</i>	91
4.2.5	<i>Measurements</i>	92
4.2.5.1	Fluorescence landscapes.....	92
4.2.6	<i>Linearity Studies</i>	93
4.2.7	<i>Data analysis</i>	93
4.3	RESULTS AND DISCUSSION.....	94
4.3.1	<i>Preliminary investigations</i>	94
4.3.1.1	Algal landscapes.....	94
4.3.1.2	Linearity.....	95
4.3.2	<i>Preliminary study</i>	97
4.3.2.1	Chlorophyll a concentration	97
4.3.2.2	PARAFAC results	98
4.3.3	<i>Final study</i>	102
4.3.3.1	Chlorophyll a concentration	102
4.3.3.2	Yield measurements	103
4.3.3.3	PARAFAC results	103
4.4	CONCLUSIONS	107
CHAPTER 5: APPLICATION OF PARAFAC TO LATENT FINGERPRINT VISUALISATION		109
5.1	INTRODUCTION.....	109
5.1.1	<i>Fingerprints on difficult surfaces</i>	110
5.1.1.1	Hyperspectral imaging	110
5.1.1.2	Fluorescence imaging.....	112
5.1.1.3	Nanoparticles	114
5.1.1.4	Other techniques	115
5.1.2	<i>Background correction</i>	116
5.1.3	<i>Aims</i>	116

5.2	EXPERIMENTAL AND DATA ANALYSIS METHODS	117
5.2.1	<i>Samples</i>	117
5.2.1.1	Drink can	117
5.2.1.2	Polymer banknotes.....	117
5.2.1.3	Glossy printed card.....	118
5.2.2	<i>Instrumental</i>	118
5.2.2.1	Polilight / Poliview system	118
5.2.2.2	HSI Examiner™ 100 QD.....	119
5.2.3	<i>Data preprocessing</i>	120
5.2.3.1	Polilight / Poliview system	120
5.2.3.2	HSI Examiner™ 100 QD.....	120
5.2.4	<i>Data analysis</i>	120
5.2.4.1	PARAFAC.....	120
5.2.4.2	PCA.....	121
5.2.4.3	MIA approach	121
5.3	RESULTS AND DISCUSSION.....	121
5.3.1	<i>Aluminium drink can</i>	121
5.3.1.1	PARAFAC.....	123
5.3.1.2	PCA.....	126
5.3.1.3	MIA approach	127
5.3.2	<i>Banknote (Polilight)</i>	128
5.3.2.1	PARAFAC.....	129
5.3.2.2	PCA.....	130
5.3.2.3	MIA approach	132
5.3.3	<i>Banknote (HSI Examiner™)</i>	133
5.3.3.1	PARAFAC.....	135
5.3.3.2	PCA.....	136
5.3.4	<i>Glossy printed card (Polilight)</i>	137

5.3.4.1	PARAFAC.....	137
5.3.4.2	PCA.....	140
5.3.5	<i>Glossy Card (HSI Examiner™)</i>	140
5.3.5.1	PARAFAC.....	141
5.3.5.2	PCA.....	141
5.3.5.3	MIA approach	142
5.3.6	<i>General Discussion</i>	143
5.3.6.1	Comparison of PARAFAC and PCA.....	143
5.3.6.2	Background interference	143
5.3.6.3	MIA approach	146
5.4	CONCLUSIONS	147
CHAPTER 6:	CONCLUSIONS AND FURTHER WORK	149

List of figures

Figure 1-1 Diagrammatical representation of an hyperspectral data cube (image from Tahtouh et al. ²).	2
Figure 1-2 A portion of the electromagnetic spectrum, showing representative molecular processes that occur when light in each region is absorbed (image adapted from Harris ⁴).	3
Figure 1-3 Approximate stretching vibrations in the MIR region. ⁸	9
Figure 1-4 Classification of methods for hyperspectral image analysis (adapted from Gendrin et al. ⁶⁶).	19
Figure 1-5 An example of a dendrogram from HCA.....	24
Figure 1-6 Schematic of the SOFM training process. ²⁴⁵	26
Figure 1-7 The MIA approach (A) The scatter score plot (PC1 vs PC2). (B) The defined classes in the score plot of (A). (C) The classes in the original image. ²⁷⁰	28
Figure 1-8 Diagrammatical representation of PLS, where X is the spectral data, c is the reference values, T is the common scores matrix, P and q are loading matrices, and E and f are error matrices. ²⁴⁷	29
Figure 1-9 An illustration of the difference in the way that SAM (left panel) and SFF (right panel) compare a known endmember spectrum to an unknown pixel spectrum. ²⁷⁷	32
Figure 1-10 Diagrammatical representation of PARAFAC.....	33
Figure 2-1 Schematic of the general approach to dimension reduction and classification of the resulting feature vectors.	42
Figure 2-2 Schematic of data treatment for the MIQ method (data is treated analogously in the ASD method); "M1 S1" is Mixture 1-Sample1, etc.....	45
Figure 2-3 Construction of median-interquartile range "super-spectrum" used in the MIQ method.....	46

Figure 2-4 Conversion of a cluster image (such as from HCA) to a feature vector for an image; this vector is simply a histogram of the number of pixels in the image that belong to each cluster.....	47
Figure 2-5 Representative spectra from a reflectance FTIR spectral image of Mixture 1 (25% w/w of each anthracene, anthraquinone, aspirin, and citric acid).....	50
Figure 2-6 Two perspectives of the discriminant scores plot obtained by classifying scores from the PCA of the single-point spectra.	53
Figure 2-7 Two perspectives of the discriminant scores plot obtained by classifying scores from the PCA of the MIQ super-spectra from the images.	54
Figure 3-1 PCA scores plot for LA-ICP-MS data using silicon as IS.	71
Figure 3-2 PCA scores plot for LA-ICP-MS elemental ratios.	72
Figure 3-3 Average MIR spectra for the soil locations.	74
Figure 3-4 LDA scores plot for MIQ of MIR image data (filled circles represent the training samples and open circles test samples).	75
Figure 3-5 LDA scores plot for MIQ of LA-ICP-MS elemental maps (filled circles represent the training samples and open circles test samples).	75
Figure 3-6 LDA scores plot for MIQ of LA-ICP-MS elemental maps from final sample set (filled circles represent the training samples and open circles test samples).....	80
Figure 3-7 PCA scores plot for MIQ of LA-ICP-MS elemental maps from final sample set (filled circles represent the training samples and open circles test samples).....	81
Figure 4-1 A example of a fluorescent landscape.....	88
Figure 4-2 Fluorescent landscape for <i>Chaetoceros</i> sp.....	95
Figure 4-3 Fluorescent landscape for <i>Dunaliella tertiolecta</i>	95
Figure 4-4 Fluorescence intensity vs. chlorophyll a concentration at chlorophyll a peak for <i>Chaetoceros</i> sp..	96

Figure 4-5 Fluorescence intensity vs. chlorophyll a concentration at chlorophyll a peak for <i>Dunaliella tertiolecta</i>	97
Figure 4-6 Chlorophyll a concentration measured using Trilogy Fluorometer in the preliminary study.....	98
Figure 4-7 PARAFAC components in preliminary study.	99
Figure 4-8 PARAFAC scores for component 1 in preliminary study.....	100
Figure 4-9 PARAFAC scores for component 2 in preliminary study.....	100
Figure 4-10 PARAFAC scores for component 3 in preliminary study.....	101
Figure 4-11 PARAFAC scores for component 4 in preliminary study.....	101
Figure 4-12 Chlorophyll a concentration measured using Trilogy Fluorometer in the final study.....	102
Figure 4-13 PARAFAC components in final study.	103
Figure 4-14 PARAFAC scores for component 1 in final study.	105
Figure 4-15 PARAFAC scores for component 2 in final study.	105
Figure 4-16 PARAFAC scores for component 3 in final study.	106
Figure 4-17 PARAFAC scores for component 4 in final study.	106
Figure 5-1 Fluorescent image of ethyl cyanoacrylate-fumed fingerprint on aluminium drink can stained with Rhodamine 6G (λ_{ex} 490 nm, λ_{em} 555 nm).....	122
Figure 5-2 Average fluorescent landscapes from different regions of aluminium drink can..	122
Figure 5-3 PARAFAC score images from ethyl cyanoacrylate-fumed fingerprint on aluminium drink stained with Rhodamine 6G.	124
Figure 5-4 PARAFAC score images after mean centring of fluorescent landscapes from ethyl cyanoacrylate-fumed fingerprint on aluminium drink stained with Rhodamine 6G.....	125

Figure 5-5 PCA score images from ethyl cyanoacrylate-fumed fingerprint on aluminium drink stained with Rhodamine 6G.	126
Figure 5-6 PARAFAC scores scatter plot with the selected pixels green.	127
Figure 5-7 MIA image from PARAFAC of aluminium drink can.....	128
Figure 5-8 MIA image from PCA of aluminium drink can.....	128
Figure 5-9 Fluorescent image from Polilight of fresh fingerprint on banknote powered with Brilliant Yellow ($\lambda_{ex} = 450 \text{ nm}$, $\lambda_{em} = 530 \text{ nm}$).	129
Figure 5-10 PARAFAC score images after mean centring of fluorescent landscapes on banknote powdered with Brilliant Yellow (Polilight).	131
Figure 5-11 PCA score images after mean centring of fluorescent landscapes from fingerprint on banknote powdered with Brilliant Yellow (Polilight).	131
Figure 5-12 MIA image from PARAFAC of banknote.....	132
Figure 5-13 MIA image from PARAFAC of banknote showing ridge detail over black intaglio printing.	132
Figure 5-14 Fluorescent image from HSI Examiner™ of fresh fingerprint on banknote powered with Brilliant Yellow ($\lambda_{ex} = 315 - 400 \text{ nm}$, $\lambda_{em} = 510 \text{ nm}$).	133
Figure 5-15 PARAFAC score images after mean centring of fluorescent landscapes from fingerprint on banknote powdered with Brilliant Yellow (HSI Examiner™).	134
Figure 5-16 Emission loadings from PARAFAC of banknote (HSI Examiner™).....	135
Figure 5-17 PCA score images after mean centring of fluorescent landscapes from fingerprint on banknote powdered with Brilliant Yellow (HSI Examiner™).....	136
Figure 5-18 Fluorescent image from Polilight of ethyl cyanoacrylate-fumed fingerprint on glossy printed card powered with Brilliant Orange ($\lambda_{ex} = 490 \text{ nm}$ & $\lambda_{em} = 610 \text{ nm}$).	137
Figure 5-19 PARAFAC score images after mean centring of fluorescent landscapes from fingerprint on glossy card powdered with Brilliant Orange (Polilight).	138

Figure 5-20 PCA score images after mean centring of fluorescent landscapes from fingerprint on glossy card powdered with Brilliant Orange (Polilight).....	139
Figure 5-21 Fluorescent image from HSI Examiner™ of ethyl cyanoacrylate fumed fingerprint on glossy printed card powdered with Brilliant Orange ($\lambda_{ex} = 450-500$ nm, $\lambda_{em} = 590$ nm).....	140
Figure 5-22 PARAFAC score images after mean centring of fluorescent landscapes from fingerprint on glossy card powdered with Brilliant Orange (HSI Examiner™).....	141
Figure 5-23 PCA score images after mean centring of fluorescent landscapes from fingerprint on glossy card powdered with Brilliant Orange (HSI Examiner™).	142
Figure 5-24 MIA image from PARAFAC of glossy card (HSI Examiner™).....	142
Figure 5-25 Comparison of fluorescent powder emission spectra and PARAFAC loadings for HSI Examiner™ datasets.	144
Figure 5-26 Comparison of fluorescent powder emission spectra and PARAFAC loadings for Polilight datasets.....	145
Figure 5-27 Average emission spectra from 400-450 nm excitation of banknote using HSI Examiner™.....	146

List of tables

Table 2-1 Compositions of the four powder mixtures.	43
Table 2-2 Summary of results for image classification methods.	52
Table 3-1 Suburban soils collected for preliminary study.	64
Table 3-2 Experimental parameters for LA-ICP-MS analysis.	66
Table 3-3 Groups for calculation of elemental ratios in preliminary study.	66
Table 3-4 Suburban soils collected for final study.	68
Table 3-5 Experimental parameters for LA-ICP-MS element selection.	69
Table 3-6 Groups for calculation of elemental ratios in final study.	70
Table 3-7 LDA correct classification percentages for preliminary soil sample set.	73
Table 3-8 ANOVA and Kruskal-Wallis test results for element selection study (N.B. sig. < 0.002 used due to the number of variables).	77
Table 3-9 LDA correct classification percentages for final soil sample set.	79
Table 4-1 Kodak Wratten filters used in light table for algal studies.	91
Table 4-2 Experimental parameters for Cary Eclipse Spectrophotometer in preliminary study.	92
Table 4-3 Experimental parameters for Cary Eclipse Spectrophotometer in final study.	93
Table 4-4 Dilution series for linearity studies.	93
Table 5-1 Main constituents of latent fingerprint deposits. ⁴²⁹	109
Table 5-2 Wavelengths collected for each sample using the Polilight / Poliview system.	119

Table 5-3 Experimental parameters for collection of fluorescent images using HSI Examiner™.
.....119

Abbreviations

ASD	average-standard deviation
ASMULIM	analysis of a set of multivariate images
ATR	attenuated total reflection
chl	chlorophyll
DA	discriminant analysis
DFO	1,8-diazafluorene-9-one
FPA	focal plane array
HCA	hierarchical cluster analysis
HPLC	high pressure liquid chromatography
IS	internal standard
LA-ICP-MS	laser ablation inductively coupled plasma mass spectrometry
LOO	leave-one-out
MIA	multivariate image analysis
MIQ	median-interquartile range
MIR	mid-infrared
NIR	near-infrared
PARAFAC	parallel factor analysis

PCA	principal component analysis
SAM	spectral angle mapper
SIMCA	soft independent modelling of class analogy
SOFM	self organising feature map
UV	ultraviolet

Abstract

The majority of the methods that have been used for the analysis of hyperspectral images have focussed on classification of spectra within an image, and there have been few examples of using whole hyperspectral images (one image per sample) for classification purposes. In this project, feature vectors were created to capture the heterogeneity present in each hyperspectral image for subsequent classification. All of the feature vectors methods were compared to single-point spectra and literature methods. The MIQ (the median-interquartile superspectrum) achieved the best classification of a set of heterogeneous powder mixtures, classifying 100% of the external test set.

Mid-infrared (MIR) hyperspectral images and LA-ICP-MS elemental maps were collected of 'real-world' samples of soil collected from various locations around Sydney. For both the MIR and LA-ICP-MS data, the MIQ feature vector achieved superior classification compared to the averages alone (used to represent single-point spectra). For the Sydney soil samples, the classification performance of the LA-ICP-MS elemental maps was superior to MIR reflection hyperspectral images. For the LA-ICP-MS data, using elemental ratios for calibration proved to be a more suitable strategy than using silicon as an internal standard.

A major application of PARAFAC has been the analysis of fluorescent landscapes. In the other main area of this project, novel applications of PARAFAC were investigated. The first application explored was the use of PARAFAC and fluorescent landscapes for the characterisation of phytoplankton. Algal species were placed under different wavelengths of light and monitored over time to determine their response to these environments. PARAFAC was able to isolate components that were identified as β -carotene, chlorophyll *a* and *b*, and chlorophyll *a* degradation products. The advantage of using PARAFAC for the characterisation of phytoplankton is that it can simultaneously determine all pigment concentrations without the need for prior separation or extraction.

Background interference due to highly coloured / patterned and fluorescence remains an issue with many techniques of fingerprint visualisation. The second application of PARAFAC in this project was to analyse fluorescent landscapes constructed by collecting series of images of fluorescent fingerprints on these traditionally difficult backgrounds. Whilst PARAFAC successfully isolated the fluorescence due to the fingerprint, there was typically

only a small reduction in background interference. However, it is expected that PARAFAC would be more successful for samples where the background is also fluorescent, such as DFO-developed prints on fluorescent yellow paper.