THE INFLUENCE OF DROUGHT, AND OTHER ABIOTIC FACTORS ON TREE WATER USE IN A TEMPERATE REMNANT FOREST

Melanie Zeppel

October 2006

A thesis submitted in fulfilment of the requirements for the degree of Doctor of Philosophy in The Department of Environmental Sciences at The University of Technology, Sydney

Declaration

I certify that the work in this thesis has not previously been submitted for a degree nor has it been submitted as part of requirements for a degree except as fully acknowledged within the text.

I also certify that the thesis has been written by me. Any help that I have received in my research work and the preparation of the thesis itself has been acknowledged. In addition, I certify that all information sources and literature used are indicated in the thesis.

Production Note:
Signature removed prior to publication.

Work from this thesis has directly contributed to the following papers:

Zeppel M.J.B., Yunusa I.A.M. and Eamus D., Daily, seasonal, and annual patterns of transpiration from a stand of remnant vegetation dominated by a coniferous *Callitris* species and a broad-leaved *Eucalyptus* species. (accepted: *Physiologia Plantarum*).

Zeppel M.J.B., and Eamus D., The influence of species, tree size and season on tree water use, Huber value, canopy conductance and canopy decoupling in a native remnant evergreen woodland (submitted: *Tree Physiology*).

Eamus D., Macinnis-Ng C. M., Hose G.C., Zeppel M.J.B., Taylor D.T. and Murray B. M. (2005) Turner Review No. 9: Ecosystem services: an ecophysiological examination. *Australian Journal of Botany* **53**, 1-19.

Zeppel M.J.B. and Eamus D. (2005) Tree water use under conditions of drought. *Agricultural Science* **17**, 8-11.

Zeppel M.J.B., Murray B.R., Barton C. and Eamus D. (2004) Seasonal responses of xylem sap velocity to VPD and solar radiation during drought in a stand of native trees in temperate Australia. *Functional Plant Biology* **31**, 461-470.

Zeppel M.J.B., Murray B.R. and Eamus D. (2003) The potential impact of dryland salinity on the threatened flora and fauna of New South Wales. *Ecological Management and Restoration* **4** (Supplement), 53-59.

Murray B.R., Zeppel M.J.B., Hose G.C. and Eamus D. (2003) Groundwater-dependent ecosystems in Australia: It's more than just water for rivers. *Ecological Management and Restoration* **4**, 110-113.

Acknowledgements

Many people have been generous with their time, equipment, and support during my candidature.

I have been lucky to have three supportive supervisors. Professor Derek Eamus has always been there to answer questions, provide interesting scientific questions and make sure that the 'bar' is high enough. Dr Brad Murray has encouraged me, improved my writing style dramatically, and taught me much about biology, and statistics. Dr Craig Barton has patiently helped with calibrating equipment, and proof read many drafts.

I especially would like to thank my lab, for most importantly for keeping me well fed, and listening to me. Cate has been a wonderful guide, cook, friend, and listened to all my crazy ideas, as well as trying to keep me obeying the rules which are in place for my own benefit (no texting while driving!). Thanks to Grant for always being positive, and supportive of the life/work balance, as well as the amazing wig collection. Isa has shared his extensive knowledge of soil-plant interactions. Dan has been quietly reminding me of the excitement and keenness a new PhD candidate has.

Many DIPNR staff have provided equipment, data and useful suggestions. It has been a privilege to work on the Recharge Validation Project. Rick Young, from Tamworth provided climate and Bowen ratio data, while Mark Littleboy showed me that my data were useful for a state-wide salinity project, as well as providing insights into the insanity of a PhD.

For assistance in the field, I thank Derek Eamus, Susan Williams, Bev Terry, Mike Zeppel, Cate Macinnis-Ng, Grant Hose, Liz Crawford, Mal Johnston, Kath Poole and Melinda Collett.

I give deep thanks to my friends, Susan and Greg, for allowing me to turn up to their house at any time of day or night. Hannah (and her sister) also gave me a much needed sense of perspective.

My favourite community, Small Boat, Big Sea, was always there for me, subtly reminding me of priorities, and providing an extremely significant support network, without which I would have floundered.

Thanks to my parents, and grandmother, who encouraged and convinced me that I could do anything I wanted if I worked hard enough at it, which I have never ceased to believe.

Most of all, Michael Zeppel has provided boundless patience, kindness and emotional support, understanding and offering support while my mind and or/body has been elsewhere.

Abstract

This thesis presents the results of work undertaken to examine daily, seasonal and annual patterns of water use by a native remnant woodland in temperate Australia. The focus of the study was on the two dominant tree species of the woodland, but limited assessments of understorey and soil evapotranspiration were also undertaken. One of dominant species was *Eucalyptus crebra*, a broad-leaved tree and the other was *Callitris glaucophylla* a needle-leaved tree. At the start of the study, much of the eastern seaboard of Australia experienced a severe and prolonged drought but towards the end of the field work, rainfall at the site was significantly larger than the long-term average. This provided a fortuitous opportunity to compare the responses of vegetation water use to drought and non-drought periods.

The study was conducted on the Liverpool Plains, of western New South Wales, Australia. Principle methods applied were (a) use of heat-pulse technology to measure rates of sap flow through trees; (b) open-top chambers to measure understorey and soil evapotranspiration; (c) application of the Penman-Monteith equation to estimate canopy conductance and transpiration rates; (d) two methods to scale spatially from measurements of individual trees to estimates of stand water use; (e) three methods to scale temporally from measurements conducted over a few weeks each year to provide annual estimates of stand water use; (f) a simple water balance was constructed to approximate the rate of deep drainage of water (rate of recharge). An annual water budget for the site was estimated for the drought and post-drought periods.

The relationship between tree water use and diameter at breast height (DBH) was similar for the two species in each season, but the relationship for both species differed between seasons and years. In contrast, the relationships amongst DBH, sapwood area and leaf area differed between species at all times. This suggests that

the same rate of water use by the eucalypt and *Callitris* (at a common size) was achieved through different mechanisms.

Daily rates of stand water use showed significant intra- and inter-seasonal variation, with the lowest rates observed in winter and largest rates in summer. A simple model based upon solar radiation and vapour-pressure deficit was able to account for approximately 80 % of variation of stand water use under summer conditions with wet soil.

Estimates of stand water use derived from the Penman-Monteith equation generally agreed well with those based upon measurements of sap velocity, with a slope of the regression of the two estimates being 1.03.

In the drought-year, stand water use was approximately 59 % of rainfall and recharge was approximately 2% but in the post-drought year, when rainfall doubled compared to the drought year, stand water use was also 59 % whereas recharge was 4 %. This showed that despite the impact of an extensive and pronounced drought, the trees were able to rapidly adjust to more favourable conditions and maintain a low rate of recharge.

These results are discussed in relation to the management of water resources for human consumptive use and in relation to the development of dryland salinity across Australian landscapes that have been cleared of trees.

Table of Contents

Acknowledgements	
Table of Contents	
List of Figures	
List of Plates	
List of Abbreviations	xviii
Chapter One: Introduction	1
1.0 Understanding the role of forests in water resource management	1
1.1 Knowledge of tree water relations assists in managing water resources	1
1.2 Whole tree water use	4
1.3 Spatial scaling from whole tree to stand water use	5
1.4 Temporal scaling	7
Chapter Two: Vegetation Structure	11
2.0 Introduction	11
2.1 Methods	12
2.1.1 Site description	12
2.1.2 Site description - soils	13
2.1.3 Climate	16
2.1.4 Soil moisture	16
2.1.5 Tree size and basal area	16
2.1.6 Leaf area index	17
2.1.7 Understorey leaf area	17
2.1.8 Foliage projective cover	18
2.1.9 Above ground biomass and carbon estimates	18
22 D 1	1.0

	2.2.1 Climate and rainfall	18
	2.2.2 Species present	22
	2.2.3 Soil moisture	22
	2.2.4 Basal area	22
	2.2.5 Foliage projective cover (FPC) and leaf area index (LAI)	22
	2.3 Discussion	28
	2.3.1 Leaf area index of other Australian temperate woodlands	29
	2.3.2 Influence of rainfall and soil water content on LAI	29
	2.3.3 Conclusion	30
Cl	napter Three: Relationships among tree size and tree water use, sapwood area, leaf	
ar	ea, Huber values and decay curves	31
	3.0 Introduction	31
	3.1 Methods	33
	3.1.1 Leaf area	33
	3.1.2 Sapwood area and depth	34
	3.1.3 Sap flow measurement	34
	3.1.4 Radial sapflow profiles	35
	3.1.5 Wound estimation	35
	3.1.6 Statistical analysis	35
	3.1.7 Huber value	36
	3.1.8 Half time and decay curves	36
	3.2 Results	36
	3.2.1 Tree size <i>versus</i> water use – does the relationship vary between species?	36
	3.2.2 Does the relationship between tree size and water use vary seasonally?	39
	3.2.3 Tree size <i>versus</i> leaf area	39
	3.2.4 Tree size <i>versus</i> sapwood area	40
	3.2.5 Leaf water potential	42
	3.2.6 Huber value	45
	3.2.7 Half time of decay curve after rain	45

	3.2.8 Canopy conductance, canopy decoupling and transpiration rate	. 49
	3.2.9 Sap velocity <i>versus</i> tree size	. 49
	3.3 Discussion	. 49
	3.3.1 Weather and leaf water potential	. 49
	3.3.2 Tree size, leaf area and sapwood area vary with season and species	. 50
	3.3.3 Huber values	. 51
	3.3.4 Tree size <i>versus</i> water use – does the relationship vary between species and	
	seasonally?	. 52
	3.3.5 Decay curve	. 53
	3.3.6 Sap velocity and tree size	. 55
	3.3.7 Decoupling, maximum canopy conductance and transpiration rate	. 55
	3.3.8 Conclusion	. 58
Ch	apter Four: The influence of vapour pressure deficit and solar radiation on sap	
	velocity	59
	4.0 Introduction	59
	4.1 Methods	61
	4.1.1 Site description and climate	61
	4.1.2 Sap flow measurement	61
	4.1.3 Data analyses	62
	4.2 Results	62
	4.2.1 Climate and soil moisture	62
	4.2.2 The relationship between sap velocity, VPD and solar radiation	63
	4.3 Discussion	71
	4.3.1 Hysteresis in water use-VPD relationships	72
	4.3.2 Hysteresis in winter and summer	74
	4.3.3 Hysteresis in the light response curve of sap velocity	
	4.3.4 The relationship between sap velocity, solar radiation and VPD	
	4 3 5 Conclusion	. 76

Chapter Five: Stomatal conductance, transpiration and the Penman-Monteith equation	77
5.0 Introduction	77
5.1 Methods	79
5.1.1 Stomatal conductance	79
5.1.2 Weather data	80
5.1.3 Heat pulse estimates of transpiration	80
5.1.4 Leaf area	80
5.1.5 Time lag	81
5.1.6 Penman-Monteith Equation	81
5.1.7 Three phase response of stomatal conductance to transpiration	82
5.1.8 Estimating daily stand transpiration using a simple model and the Penman-	
Monteith equation	83
5.2 Results	86
5.2.1 A comparison of E _{hp} and E _{PM}	86
5.2.2. The impact of soil water content on the correlation between E_{PM} and E_{hp}	87
5.2.3 The three phase response of stomatal conductance to transpiration in the field	92
5.2.4 The relationship between VPD, G _s and E _{hp}	94
5.2.5 The three phase response of stomatal conductance to transpiration – between	
tree variation	96
5.3 Discussion	98
5.3.1 E _{PM} compared with E _{hp}	98
$5.3.2~E_{PM}$ compared with E_{hp} as soil water content declines, and in different seasons	99
5.3.3 Adjusting for time lag	
5.3.4 The three phase of G _s to transpiration is shown in the field	100
5.3.5 The influence of VPD, soil water content, and radiation on stomatal	
conductance	101
5.3.6 Conclusion	102
Chapter Six – Spatially scaled estimates of tree and understorey evapotranspiration	104
6.0 Introduction	

	6.1 Methods	107
	6.1.1 Individual tree water use	107
	6.1.2 Spatial scaling – the DBH versus Q method	107
	6.1.3 Spatial scaling – the E _{saxsv} method	108
	6.1.4 Daily estimates of understorey evapotranspiration (E _u)	109
	6.2 Results	110
	6.2.1 Stand water use calculated using the DBH <i>versus</i> Q method	110
	6.2.2 Stand water use calculated using the E _{saxsv} method	111
	6.2.3 A comparison of methods	113
	6.2.4 Understorey E _u	118
	6.2.5 Forest Evapotranspiration (trees and understorey)	118
	6.3 Discussion	121
	6.3.1 Understorey water use	121
	6.3.2 What proportion does E _u contribute to total evapotranspiration?	122
	6.3.3 Stand water use in winter and summer	122
	6.3.4 Stand water use and basal area	123
	6.3.5 A comparison of methods.	123
	6.3.6 Conclusion.	124
Ch	napter Seven - Temporal scaling of evapotranspiration - annual estimates	125
	7.0 Introduction	125
	7.1 Methods	128
	7.1.1 The E _{saxsv} method	128
	7.1.2 Penman-Monteith estimates of daily stand transpiration (E _{PM})	128
	7.1.3 Predicting transpiration from climate and soil moisture (E _m)	129
	7.1.4 Plantation forest	129
	7.1.5 Pasture evapotranspiration estimated by the Bowen Ratio method	130
	7.2 Results	130
	7.2.1 The DBH versus Q method	130
	7.2.2 The E _{saxsv} method	130

	$1.2.3$ A comparison of stand transpiration estimated using the E_{saxsv} method and	
	using the E _{PM} method.	137
	7.2.4 Relationships among daily transpiration, radiation and VPD	139
	7.2.5 The 'broken stick' relationship	142
	7.2.6 A simple model to predict transpiration (E _m)	143
	7.2.7 Comparing the three methods	146
	7.2.8 Annual stand transpiration	146
	7.3. Discussion	150
	7.3.1 Comparing methods of temporal scaling	150
	7.3.2 Seasonal patterns	150
	7.3.3 Benefits and limitations of the E _{saxsv} method	151
	7.3.4 Benefits and limitations of the E _{PM} method	151
	7.3.5 Benefits and limitations of the E _m method	154
	7.3.6 The influence of radiation, VPD and soil moisture on Em	154
	7.3.7 Annual stand water use	156
	7.3.8 Conclusion	158
Ch	apter Eight - Discussion	159
	8.0 Introduction	159
	8.1 Relationships among DBH, sapwood area and leaf area	160
	8.2 Relationships between weather and sap velocity.	163
	8.3 Spatial and temporal scaling	165
	8.4 Annual water balance, reforestation and salinity	167
	8.5 Limitations of this thesis	169
	8.6 Future research directions	169
	8.7 Conclusions	171
Cha	apter Nine - References	173

List of Figures

Fig. 2. 1. Location of study area	1
Fig. 2. 2. Daily patterns of radiation and VPD for summer 2002/3 and winter 200219	9
Fig. 2. 3. Seasonal patterns of radiation, potential evaporation, VPD, and soil moisture content	С
Fig. 2. 4. Rainfall over the study period, from June 2002 to December 2004, compared with 50 year average rainfall.	1
Fig. 2. 5. Frequency histograms of <i>E. crebra</i> (a) and <i>C. glaucophylla</i> (b)24	4
Fig. 2. 6. Seasonal patterns of leaf area index in <i>E. crebra/C. glaucophylla</i> remnant forest	5
Fig. 3. 1. The relationship between DBH (cm) and daily total tree water use (L d-1) for (a) winter 2002, (b) summer 2002/3, (c) winter 2003, and (d) summer 2003/437	7
Fig. 3. 2. The relationship between DBH (cm) and total tree leaf area (m2) during (a) summer 2003 and (b) winter 2003 and (c) summer 2004	8
Fig. 3. 3. The relationship between DBH (cm) and sapwood area (cm2)40	С
Fig. 3. 4. The diurnal time course of leaf water potential for <i>E. crebra</i> (closed circles) and <i>C. glaucophylla</i> (open circles) during summer 2002/3, winter 2003 and summer 2003/4	1
Fig. 3. 5. The relationship between Huber value and tree height (m)	3
Fig. 3. 6. A seasonal comparison of Huber value for both species	4

Fig. 3. 7. The relationship between decay curve half-time after a rain event and soil water storage (a), and tree size (b)
Fig. 3. 8. Seasonal changes of maximum stomatal conductance (a) and Omega (b) of <i>E. crebra</i> and <i>C. glaucophylla</i>
Fig. 3. 9. Seasonal changes of maximum transpiration (c) of <i>E. crebra</i> and <i>C. glaucophylla</i>
Fig. 3. 10. The relationship between sap velocity and tree size (DBH, cm) in (a) summer 2002/3 plus summer 2003/4 and (b) winter 200348
Fig. 4. 1a, and b. The relationship between vapour pressure deficit (kPa) and sap velocity (cm h ⁻¹)
Fig. 4. 1 c and d. The relationship between vapour pressure deficit (kPa) and sap velocity (cm h ⁻¹)65
Fig. 4. 2. The daily pattern of tree water use (Q) over four consecutive days during drought, in winter67
Fig. 4. 3. Hysteresis area is strongly related to maximum daily VPD
Fig. 4. 4. A three dimensional linear plot of sap velocity for <i>Callitris glaucophylla</i> 69
Fig. 4. 5. A three dimensional linear plot of sap velocity for <i>Eucalyptus crebra</i> 70
Fig. 5. 1. The relationship between E_{hp} and E_{PM} for summer 2003/4 and winter 2003.
Fig. 5. 2. The relationship between E_{hp} and E_{PM} during summer 2004/5 for 3 trees89
Fig. 5. 3. The relationship between E_{hp} and E_{PM} for 4 trees during winter 200390

Fig. 5. 4. The impact of soil water content on E_{PM} as a predictor of E_{hp}	91
Fig. 5. 5.The three phase response in the diurnal relationship between G_c and E_{hp} , over four sampling periods during daylight.	93
Fig. 5. 6. The influence of VPD on the three phase response – the 3D relationship between VPD, $E_h p$ and G_c .	95
Fig. 5. 7. The relationship between G_c and E_{hp} for trees of different size	97
Fig. 6. 1. Seasonal estimates of stand water use using the DBH vs Q method. Seasonal means with s.e. bars are shown.	111
Fig. 6. 2. Daily stand water use (mm d^{-1}) from December 2002 to January 2005, estimated using the E_{saxsv} method. Grey columns show daily rainfall (mm)	115
Fig. 6. 3. The relationship between transpiration estimates of the plantation and the remnant forest (using the E_{saxsv} method).	116
Fig. 6. 4. A comparison of stand water use during winter 2003 (a) and summer 2003/4 (b).	117
Fig. 6. 5. Daily stand water use (mm d ⁻¹) over 2 years.	118
Fig. 6. 6. Understorey evapotranspiration of three ground cover types.	120
Fig. 7. 1. Estimates of stand transpiration using the DBH versus Q method	132
Fig. 7. 2. A comparison of daily transpiration estimated using the E_{saxsv} and the DBH versus Q methods.	134
Fig. 7. 3. A comparison of E_{saxsv} of the remnant woodland with daily transpiration of a nearby plantation and pasture.	135
Fig. 7. 4. A comparison of F _m and F _{cavey} from December 2002 to January 2005	136

Fig. 7. 5. A comparison of daily estimates of E _{pm} and E _{saxsv} , showing daily patterns over summer 2004
Fig. 7. 6. The relationship between daily transpiration estimated using E_{pm} and $E_{\text{saxsv.}}$
Fig. 7. 7. The relationship between daily transpiration and radiation
Fig. 7. 8. The relationship between daily transpiration and VPD14
Fig. 7. 9. The two groups of relationships between VPD and daily transpiration under different soil water scenarios
Fig. 7. 10. The 'broken stick' relationship between E_{saxsv} and E_{pm} and soil moisture content.
Fig. 7. 11. The optimal relationship among VPD, radiation and E_{saxsv} , for warm conditions and wet soil
Fig. 7. 12. The response surface describing the predicted relationship (E_m) among E_{saxsv} , radiation and VPD
Fig. 7. 13. Daily estimates of transpiration calculated using 3 methods (E_{saxsv} , E_{PM} , and E_{m})
Fig. 7. 14. The relationship between E_m and $E_{saxsy}(a)$, and between E_m and $E_{pm}(b)$

List of Tables

Table 2.1. Parameters measured at each site.
Table 2.2. Basal area of the trees present at the site
Table 2.3. Foliage projective cover of the ground cover in the remnant vegetation at Paringa, comparing winter and summer
Table 2.4. Leaf area index of the remnant vegetation at Paringa during winter and summer
Table 2.5. Basal area, above-ground biomass and carbon at Paringa and Pine Ridge State Forest
Table 5. 1 Abbreviations and derivations used in this chapter86
Table 7. 1 Comparison of E _{saxsv} method with DBH <i>versus</i> Q method
Table 7. 2. Annual rainfall and transpiration estimated using E_{saxsv} , E_{pm} , and E_{m} methods, and % of available rainfall.
Table 7. 3. A comparison of annual stand water use within a range of forest types158

List of Plates

Plate 2. 1. Paringa site from 12 m above ground in crane during summer 2003/415
Plate 2. 2. Pine Ridge State Forest study area during summer 2003/415
Plate 6. 1. The open top chamber set up in the pasture between the plantation and the remnant forest. The plantation is visible in the background112

List of Abbreviations

° C Degrees Celsius

ANCOVA Analysis of Co-variance

ANOVA Analysis of Variance

cm Centimetre

DBH Diameter at Breast Height (cm)

E_{hp} Transpiration estimated using the heat pulse system (mm³ water day⁻¹

mm² leaf area)

 $E_{\mbox{\tiny PM}}$ transpiration estimated using the Penman Monteith equation

E_{SAxSV} Stand transpiration estimated by multiplying the stand sapwood area by

sap velocity (mm³ water day⁻¹ mm⁻² ground area)

E_u Understorey evapotranspiration (mm³ water day¹¹/hr¹ mm⁻² ground

area)

G_a Aerodynamic conductance

g_s Leaf scale stomatal conductance

G_s Canopy scale stomatal conductance

G_c Stomatal canopy conductance at the canopy scale.

G_{smax} Maximum stomatal conductance at the tree scale

ha Hectare

h Hour

 J_s Sap flux (cm³ day⁻¹ cm⁻²)

kPa Kilo Pascal

LA Leaf Area (m²)

LAI Leaf Area Index (m² m⁻², unitless)

m Meter

mm Millimetre

MPa Mega-Pascal

NMM Neutron Moisture Meter

Q sap flow $(m^3 day^{-1})$

s.e. standard error

VPD Vapour Pressure Deficit (kPa)