

## **TITLE PAGE**

**Title of the article:** Pharmacists' interventions on clinical asthma outcomes. A systematic review

### **Authors:**

Victoria Garcia-Cardenas  
Graduate School of Health  
University of Technology Sydney, Sydney  
Australia

Carol Armour  
Woolcock Institute of Medical Research  
The University of Sydney, and  
Sydney Local Health District, Sydney  
Australia

Shalom I Benrimoj  
Graduate School of Health  
University of Technology Sydney, Sydney  
Australia

Fernando Martinez-Martinez  
Pharmaceutical Care Research Group  
University of Granada  
Spain

Inajara Rotta.  
Hospital Pharmacy Department, Hospital de Clínicas  
Federal University of Paraná, Curitiba  
Brazil

Fernando Fernandez-Llimos  
Research Institute for Medicines (iMed.Ulisboa), Department of Social Pharmacy  
Faculty of Pharmacy, University of Lisbon, Lisbon  
Portugal

Corresponding author:  
Victoria Garcia-Cardenas  
Graduate School of Health  
University of Technology Sydney  
City Campus, Broadway, Building 7, Lvl 4, Room 41  
PO Box 123, Broadway NSW 2007 Australia  
[Victoria.GarciaCardenas@uts.edu.au](mailto:Victoria.GarciaCardenas@uts.edu.au)  
+61 2 9514 9297

### **120-character summary:**

A recent systematic review finds that pharmacists have a positive impact on various clinical asthma outcomes

## **ABSTRACT**

This study aimed to evaluate the impact of pharmacists' interventions on clinical asthma outcomes on adult patients and to identify the outcome indicators used.

PubMed, Scopus, WoS and Scielo were searched. Studies addressing pharmacists' interventions on adult asthma patients reporting clinical asthma outcomes were incorporated.

Eleven clinical outcomes were identified in twenty-one studies. Ten studies measured the impact of the intervention on asthma control. Randomised and non-randomised controlled trials found positive results in percentages of controlled patients and Asthma Control Questionnaire scores. Discordant results were found for Asthma Control Test results. Asthma severity was assessed in four studies. A randomised controlled trial found a significant decrease in the percentage of severe patients, two non-randomised controlled trials found significant improvements in severity scores. Eleven studies reported pulmonary function indicators, showing inconsistent results. Eight studies measured asthma symptoms; three randomised controlled trials and four non-randomised controlled trials showed significant improvements.

Randomised and non-randomised controlled trials generated similar results for most outcomes. Based on the evidence generated by randomised controlled trials, pharmacists' have a positive impact on the percentage of controlled patients, Asthma Control Questionnaire scores, severity and symptoms. Future research should report using the core outcome set of indicators established for asthma (PROSPERO#CRD42014007019).

## **INTRODUCTION:**

According to World Health Organization estimates[1], 235 million people worldwide suffer from asthma, making it a major health problem in industrialised countries. The social impact of asthma is high, with negative clinical, economic and humanistic implications[2] mainly due to ineffective management of the disease.

Good asthma outcomes hinge on the accessibility to effective medications and their appropriate use by patients. Regular reviews with a healthcare provider are an essential component for effective asthma management[3]. Asthma education and training can be delivered effectively by different health care providers such as physicians, nurses or pharmacists. Given that many of the issues associated with suboptimal asthma management are related to the inappropriate use of medications[4, 5], pharmacists are in an excellent position to play an active and positive role in the management of asthma. The change in pharmacists' practice in health care to a more patient centred approach, through the provision of professional pharmacy services, supports and focuses on optimising medicines use and improving health outcomes. Several meta-analyses have shown a positive impact of pharmacists when delivering clinical services for patients with chronic conditions such as diabetes[6] or hypertension[7]. Similarly, a literature review found that community pharmacists can play an effective role in screening for poorly controlled asthma and undiagnosed COPD by delivering management interventions[8]. A narrative review revealed an expanding role in asthma care across different settings[9]. However, no systematic review of pharmacists' impact on asthma outcomes has been found.

Selecting appropriate outcomes when designing any research study is crucial when reporting the results of the research, since it allows analysing the effects of different interventions in ways that minimise bias[10]. However, there seems to be a high variability in the literature when reporting the effects of interventions on asthma patients. Difficulties caused by the heterogeneity of outcome measurements are common. This heterogeneity has direct implications when comparing and analysing the evidence available. However this problem could be addressed by the design of a 'core outcome set', which is an agreed minimum set of outcomes or outcome measures (The COMET Initiative: <http://www.comet-initiative.org>)[11]. This implies a standardisation of the variables that should be measured and reported in all trials in a specific area. Although there is a growing recognition of its relevance and some work has already been undertaken in childhood asthma, to our knowledge no core outcome set has been established for adult asthma patients in community care.

The objective of this systematic review was to evaluate the impact of pharmacists' interventions on clinical asthma outcomes on adult patients and to identify the clinical outcome indicators reported in experimental studies to assess them.

## MATERIAL AND METHODS

A systematic review was undertaken following the methodological and reporting standards recommended by PRISMA[12] and AMSTAR[13]. A literature search was conducted in August 2015. Neither publication date nor publication type filters were used. Studies assessing pharmacist interventions on adult asthma patients reporting clinical asthma outcomes as a result of the intervention provided were included. The studies eligible were those published or at least with an abstract and written with the Latin alphabet. Searches were conducted in PubMed, Scopus, Web of Science and Scielo. The queries used are described in Table 1. Duplicates records were removed.

<b>Table 1. Search strategies used in the literature retrieval</b>
<b>PubMed:</b>
("Medication Adherence"[MH] ("Patient Compliance"[MH] AND "Drug Therapy"[MH]) OR "Patient education as topic" OR "Program Evaluation"[MH] OR "Outcome and Process Assessment (Health Care)"[MH] OR "Educational Measurement" [MH] OR "Patient Care Management" [MH] OR "Preventive Health Services" [MH] OR "Health Behavior" [MH] OR "Follow-Up Studies"[MH]) AND ((Pharmacists OR Pharmacists [MH] OR "Pharmaceutical Services"[MH] OR Pharmacies OR Pharmacies [MH])) AND ((Asthma OR Asthma [MH] OR "Anti-Asthmatic Agents"[MH] OR "Adrenergic beta-2 Receptor Agonists"[MH])) AND HASABSTRACT
<b>Scopus:</b>
KEY (Pharmacist OR Pharmacy OR "community pharmacy" OR "hospital pharmacy" OR "pharmaceutical care") AND KEY (asthma OR "antiasthmatic agent") AND KEY ("disease severity" OR "forced expiratory volume" OR "forced vital capacity" OR "outcome assessment" OR spirometry OR "disease control" OR "treatment outcome" OR "point of care testing" OR "patient compliance" OR "patient education" OR "patient care" OR "medication compliance" OR "patient counseling" OR counseling OR "Asthma Control Questionnaire" OR "Asthma Control Test" OR "asthma therapy assessment questionnaire" OR "Asthma Control Scoring System")
<b>Web of Science:</b>
(TS=(pharmacist* NEAR intervention)) AND (TS=asthma)
<b>Scielo:</b>
((asma) AND (farmaceutico*)) OR (asthma AND pharmacist*)

The literature selection process was undertaken and discussed between two experts on asthma services (FFL, VGC). To identify potentially relevant articles, a screening of records retrieved from the search was performed by reviewing titles and abstracts. This process was over-inclusive. Obviously irrelevant records were removed. Potentially relevant articles were retrieved and multiple reports of the same study were linked together. Full-text papers not discarded in the screening were read, and studies were excluded according to the following criteria: (1) non-experimental studies, (2) studies with asthma patients younger than 18 years old, (3) studies not reporting a clinical asthma outcome (e.g. asthma control, asthma severity, asthma symptoms, peak expiratory flow, forced expiratory volume in the first second, or other pulmonary function indicator) (4) studies in which the educational intervention was not exclusively provided by a pharmacist. The reference lists of the retrieved papers were reviewed for potentially additional relevant studies.

Relevant information from all included studies was gathered in a pre-designed and piloted data extraction form. The following information was extracted: - Source: Study ID, citation and contact details - Eligibility: Confirmation of inclusion criteria - Objective - Methods: Study design, study groups, follow-up time, sequence generation, allocation sequence concealment, blinding - Participants: Total number of patients, total number of practices, setting, country, Inclusion/exclusion criteria. - Interventions: Number of intervention groups, intervention details, outcomes, outcomes definition, method of assessment, characteristics of the method of assessment (If ad-hoc). - Results: Number of patients and practices allocated to each group, sample size, missing participants, subgroup analysis. - Summary of data: summary of results for each clinical asthma outcome assessed - Conclusions - Miscellaneous: Funding source, references to other relevant studies, reviewer's comments.

Following AMSTAR recommendations, data synthesis and conclusions were formulated taking into consideration the epidemiological design of the studies[13]. The systematic review was registered in the PROSPERO -International Prospective Register of Systematic Reviews- database (registration#CRD42014007019), where a detailed protocol of the review can be found.

## **RESULTS**

Initially, 1194 different potential articles were retrieved from the databases used. After screening by title and abstract, 68 of them were selected for full-text review. In the second step, 41 were excluded for the following reasons (papers could be excluded due to more than one criterion): six were non-experimental studies, twenty-two included a population under eighteen years old, in ten the intervention was not delivered exclusively by a pharmacist and nineteen did

not evaluate any clinical asthma outcome. Twenty-four papers corresponding to twenty-one studies were included (Figure 1). Fourteen studies were conducted in a community pharmacy setting[14-27], two in a hospital setting[28, 29], two in a community clinic setting[30, 31], one in an outpatient medical centre[32], one in an antenatal outpatient clinic[33] and one was conducted by telephone[34]. Seven studies were conducted under a randomised controlled trial design (RCT)[16, 17, 22, 28, 29, 33, 34], two under a cluster randomised controlled trial (C-RCT) design[14, 19], two under a cluster randomised trial design (C-RT)[15, 26], two under a cluster controlled trial design[24, 25] and eight under a quasi-experimental study with no control group design[18, 20, 21, 23, 27, 30-32]. The follow-up period varied across the studies, ranging between one month[20, 26], two months[28], three months[16, 29], five months[18], six months[14, 15, 17, 19, 22, 23, 27, 31, 33, 34], nine months[25], twelve months[21, 24] and twenty-four months[30]. The follow-up period was not specified in one of the studies retrieved[32]. The main characteristics of the studies are summarised in the supplementary material.

### **Impact of pharmacist interventions on clinical asthma outcomes**

#### **Current asthma control**

Ten studies measured the impact of a pharmacist intervention on current asthma control as a main outcome[15, 17, 19, 20, 22, 25-27, 33, 34]. It was mainly assessed using validated instruments, such as the Asthma Control Questionnaire (ACQ) and the Asthma Control Test (ACT). Four studies reported a change in the number of controlled asthma patients[15, 19, 22, 27] and ten studies reported a change in either ACQ[15, 19, 20, 25, 26, 33] or ACT [17, 22, 27, 34] scores.

Five studies that measured current asthma control as a main outcome used either a RCT or a C-RCT design[17, 19, 22, 33, 34]. The two RCTs/C-RCTs that measured the change in the number of controlled asthma patients, reported an improvement in the percentage of patients considered to have good asthma control[19, 22]. Results showed that the difference in the percentage of controlled patients between study groups after the provision of the intervention was 12.1% ( $p=0.028$ ) and 7.7% (no  $p$ -value provided) respectively. One of the studies found that patients in the intervention group were almost three times more likely to have their asthma controlled than patients in the control group (OR 3.06, 95% CI:1.63-5.73;  $p<0.001$ ). The two RCT/C-RCT assessing the change in ACQ scores also found a positive impact, since both statistically and clinically significant reductions in patient's ACQ scores were observed[19, 33]. One study found a difference in ACQ scores between study groups after the provision of the intervention of 0.41

points ( $p<0.001$ )[19], whereas in another the difference was equal to 0.60 ( $p<0.001$ )[33]. Only one of the three RCTs evaluating the change in ACT scores found significant improvements in the intergroup comparisons[17]. This study found a difference in asthma control scores between study groups after the intervention equal to 2.6 ( $p<0.01$ ).

The other five studies that measured current asthma control used different research designs[15, 20, 25-27]. Two of them assessed the change in the number of controlled patients, signalling positive findings[15, 27]. One study assessed the impact of two different interventions, reporting a significant improvement in the percentage of patients considered to have good asthma control after both interventions were delivered (increase of 32% and 38% respectively, no p-value provided for the intragroup comparisons)[15]. Four studies reported the change in ACQ scores[15, 20, 25, 26]. These four found significant reductions in ACQ scores, ranging from 0.56 and 0.57 (no p-value provided), 0.4 ( $p<0.001$ ), 0.23 ( $p=0.02$ ), and 0.4 ( $p=0.003$ ) respectively. One study reported statistical and clinical significant reductions, since 48% of patients demonstrated a reduction greater or equal to 0.5 in their ACQ scores[15].

#### Asthma Severity

The impact of pharmacy interventions on asthma severity was assessed in four studies[14, 21, 23, 24]. The methods of assessment varied; two studies used the criteria established by the German Asthma Guidelines[21, 24], one used a tool based on the National Asthma Council Australia severity score[14], whereas another based the assessment on the criteria established by the Australian Asthma Management Handbook[23]. In terms of the impact of the pharmacist intervention, one study reported the change in the number of patients suffering from severe asthma[14], and three studies used the change in mean asthma severity scores as their outcome indicator.

The only RCT assessing the impact of a pharmacist intervention on asthma severity reported a significant decrease in the percentage of patients having severe asthma[14]. This study reported a significant decrease in the proportion of severe patients from 87.9% to 52.7% ( $p<0.001$ ), while the control group remained unchanged. The authors also found that patients in the intervention group were more likely to change from the severe to the not severe category than patients in the control group (OR 2.68,  $p<0.001$ ).

Three further studies assessing asthma severity as a main outcome indicator used a non-randomised design. Two studies reported significant decreases in mean asthma severity scores [21, 23] (-0.3,  $p<0.002$  and -0.3,  $p<0.001$ , respectively), and one did not find any significant

difference[24].

## Pulmonary function

Eleven studies reported some measurement of pulmonary function as an outcome indicator of the intervention designed[14, 18, 21-24, 27-30, 32]. Seven reported changes in Peak Expiratory Flow (PEF) values[18, 21-24, 29, 30], six in Forced Expiratory Volume in the First second (FEV1) values[14, 21, 24, 28, 32, 35], one in Vital Capacity (VC)[21], one in FEV1 % VC (FEV1 expressed as a percentage of the VC, Tiffeneau index)[21], one in Forced Vital Capacity (FVC)[32] and one in FEV1/FVC ratio[14]. Methods of evaluation and reporting of outcome indicators diverged. PEF values were either measured at the pharmacy[18, 21, 24, 29, 30] or self-measured by the patient[22-24]. Results were expressed as PEF rate[18, 21, 24, 29], percentage of maximum predicted PEF[22], peak flow index[23] and number of patients below 70% and 85% of optimal PEFR[30]. In terms of FEV1 values, they were measured by a pharmacist[14] or by a physician[21]. This information remained unknown in four studies[24, 27, 28, 32]. Results were reported as percentage predicted[14, 27], as an absolute number[21, 28], as a mean percentage change[32], as a percentage change from baseline[24] and as a percentage of the VC[21]. The study assessing VC reported it in absolute values[21] whereas the one measuring FVC, as mean percentage change[32]. The only study measuring FEV1/FVC reported it as a percentage of predicted value[14].

Four out of the ten studies evaluating any pulmonary function outcome indicator used a RCT design[14, 22, 28, 29]. No studies reported improvements in percentage of maximum PEF[22] or PEF rate[29]. However, one RCT did report significant improvements in FEV1 values after the intervention[28] (difference of 0.20 L between study groups, no p-value provided). One C-RCT evaluated the change in percentage of predicted FEV1/FVC, with no change after the follow-up [14].

Results from non-RCT showed two studies reporting improvements in PEF rates[18, 21]. In the first study, PEF rates ranging from 0.13 to 0.12L/min (no p-value provided) were found. In the second one, an improvement of 0.35L/min ( $p < 0.001$ ) was reported. Only one study was identified as using peak flow index as an outcome indicator [23], with results showing significant improvements after the intervention [from  $82.7\% \pm 8.2\%$  to  $87.4\% \pm 8.9\%$  ( $p < 0.001$ )]. In terms of other outcomes, one study showed significant improvements in the percentage of predicted FEV1 [from  $46.6\% \pm 0.09$  to  $70.4\% \pm 0.10$  ( $p < 0.05$ )][27]. One study reported significant improvements of mean percentage changes in FEV1 values ( $18.5L \pm 1.5$  in the intervention group vs  $5.2 \pm 1.0$  in the comparison group, no p-value provided) but no differences in FVC[32]. Two



studies did not find any effect on other pulmonary function indicators such as VC, FVC or FEV%VC[21, 32]

#### Asthma symptoms

Eight studies reported the impact of the pharmacist intervention on asthma symptoms. Three of them assessed the occurrence of general asthma symptoms[16, 21, 30], evaluated with the validated North England Asthma Symptoms Scale, with a self-reported measure and with an ad-hoc questionnaire. This outcome was reported as a mean symptoms score[16], as a self-scored punctuation[21] and as a mean number of symptoms suffered in the previous week[30]. Four studies assessed the occurrence of nocturnal asthma symptoms as outcome indicators, including mean nocturnal episodes of asthma[31], mean frequency of nocturnal asthma symptoms[29], number of nocturnal awakenings due to asthma[22], and sleep disturbances [18]. Two studies used a self-reported card/diary to assess asthma nocturnal episodes[22, 29], whereas the method of assessment was not specified in two studies[18, 31]. Two studies assessed dyspnoea severity[21, 24], rated by the patient's physician through the Medical Research Council dyspnoea scale.

Amongst the three RCTs[16, 22, 29], one showed a significant improvement in the mean asthma symptoms score[16], with a mean difference in asthma symptoms scores between study groups after the intervention equal to 7 ( $p<0.001$ ). Two studies reported positive outcomes in terms of nocturnal asthma symptoms[22, 29]; one reported a difference in mean change of nocturnal awakenings between study groups equal to 3.5 ( $p=0.004$ ) and the second found that patients in the intervention group had a greater significant decrease in the mean frequency of nocturnal symptoms than patients in the control group after 20 and 22 weeks of follow-up (no mean values provided,  $p<0.05$ ).

Of the five non-randomised studies assessing the impact on asthma symptoms, one study found a reduction in the number of patients with moderate or severe symptoms[30] (no p-value provided). Two studies reported positive findings in terms of nocturnal asthma symptoms[18, 31], and one in terms of dyspnoea severity scores[21].

Table 2 shows a summary of the pharmacists' impact on the different asthma outcomes identified and the heterogeneity in both the outcomes assessed and the method of reporting.

Table 2. Summary of findings of the impact of pharmacists' interventions on asthma outcomes				
Outcome		Reporting method	Findings in RCT and C-RCT	Findings in other research designs
Current asthma control		% Controlled patients	(+)[19, 22]**	(+)[15, 27]**
		ACQ Scores	(+)[19, 33]	(+)[15, 20, 25, 26]
		ACT Scores	(+)[17] (+)[22, 34]**	(+)[27]
Asthma severity		% Severe patients	(+)[14]	NEA
		Asthma severity score	NEA	(+)[21, 23] (+)[24]
Pulmonary function	PEF	PEF rate	(+)[29]	(+)[18, 21] (+)[24]
		Percentage of maximum predicted PEF	(+)[22]	NEA
		Peak Flow Index	NEA	(+)[23]
		PEF values below 85% of optimal PEF	NEA	(+)[30]**
		PEF values under 70% of optimal PEF	NEA	(+)[30]**
	FEV1	Percentage of predicted personal best	(+)[14]	(+)[27]
		FEV1 (absolute number)	(+)[28]**	(+)[21]**
		Mean percentage change	NEA	(+)[32]
		Percentage change of FEV1 from baseline	NEA	(+)[24]
	VC	Absolute value	NEA	(+)[21]
	FVC	Mean percentage change	NEA	(+)[32]**
	FEV1/FVC	Percentage of FEV1/FVC predicted value	(+)[14]	NEA
	FEV1%VC	FEV1%VC (FEV1 expressed as a percentage of the VC, Tiffeneau Index)	NEA	(+)[21]
Asthma symptoms	General symptoms	Mean asthma symptoms score	(+)[16]	NEA
		Self rated score	NEA	(+)[21]
		Mean number of symptoms in previous week	NEA	(+)[30]**
	Nocturnal symptoms	Mean nocturnal episodes of asthma	NEA	(+)[31]
		Mean frequency of nocturnal asthma symptoms	(+)[29]	NEA
		Number of nocturnal awakenings due to asthma	(+)[22]	NEA
		Number of patients with sleep disturbances	NEA	(+)[18]**
	Dyspnoea	Dyspnoea severity score	NEA	(+)[21] (+)[24]
<p>RCT: Randomised controlled trial, C-RCT: Cluster randomised controlled trial            (+): Positive findings, (+-): Neutral findings            NEA: No evidence available            ** no p-value provided            ACQ Asthma Control Questionnaire; ACT: Asthma Control Test; PEF: Peak Expiratory Flow; FEV1: Forced Expiratory Volume in the first second;            FVC: Forced Vital Capacity; VC: Vital Capacity</p>				

## **Meta-analysis**

The use of statistical techniques was attempted to integrate and summarize the results reported for clinical asthma outcomes. However, a meta-analysis could not be performed due to the lack of some statistical parameters needed as well as the variability in the different outcomes assessed and epidemiological designs used. Although nine studies satisfied all the requirements for meta-analysis, it was not undertaken as the high heterogeneity would have generated evidence of poor quality (Table 3).

Table 3. Individual assessment of each study regarding suitability for inclusion in meta-analysis		
Continuous outcomes		
Outcome assessed	Study	Justification
ACQ scores	Garcia-Cardenas <i>et al</i> [19]	Lack of standard deviation of mean change for control group
	Armour <i>et al</i> (2013)[15]	Lack of control group
	Giraud <i>et al</i> [20]	Lack of control group
	Smith <i>et al</i> [25]	Lack of standard deviation of mean change for both study groups
	Lim <i>et al</i> [33]	Appropriate for meta-analysis
	Toumas-Sehata <i>et al</i> [26]	Lack of control group
ACT scores	Mehuys <i>et al</i> [22]	Appropriate for meta-analysis
	Young <i>et al</i> [34]	Lack of standard deviation of mean difference
	Bereznicki <i>et al</i> [17]	Lack of baseline results for control group. Lack of mean change and standard deviation of both study groups
Asthma severity	Mangiapane <i>et al</i> [21]	Lack of control group
	Saini <i>et al</i> [23]	Lack of standard deviation of mean for both groups
	Schulz <i>et al</i> [24]	Lack of standard deviation of mean for both groups
PEF	Mangiapane <i>et al</i> [21]	Lack of control group
	Saini <i>et al</i> [23]	PEF values only in intervention group
	Mehuys <i>et al</i> [22]	Appropriate for meta-analysis
	Abdelhamid <i>et al</i> [29]	Lack of outcome values before and after in both study groups
	Petkova <i>et al</i> [18]	Lack of control group
	Schulz <i>et al</i> [24]	Lack of standard deviation of mean change for both study groups
	Narhi <i>et al</i> [30]	Lack of control group
FEV1	Mangiapane <i>et al</i> [21]	Lack of control group
	Kumar <i>et al</i> [28]	Lack of standard deviation of mean change for both study groups
	Schulz <i>et al</i> [24]	Appropriate for meta-analysis
	Armour <i>et al</i> (2007)[14]	Appropriate for meta-analysis
	de Tullio <i>et al</i> [32]	Lack of standard deviation of mean change and p value for both study groups
	Zanghelini <i>et al</i> [27]	Lack of control group
VC	Mangiapane <i>et al</i> [21]	Appropriate for meta-analysis
FVC	de Tullio <i>et al</i> 1987[32]	Results not reported
FEV1/FVC	Armour <i>et al</i> [14]	Appropriate for meta-analysis
FEV1%VC	Mangiapane <i>et al</i> [21]	Appropriate for meta-analysis
Asthma symptoms	Barbanel <i>et al</i> [16]	Appropriate for meta-analysis
	Mehuys <i>et al</i> [22]	Appropriate for meta-analysis
	Petkova <i>et al</i> [18]	Lack of control group
	Narhi <i>et al</i> [30]	Lack of control group
	Odegard <i>et al</i> [31]	Lack of control group
	Abdelhamid <i>et al</i> [29]	Appropriate for meta-analysis
	Mangiapane <i>et al</i> [21]	Lack of control group
	Schulz <i>et al</i> [24]	Lack of standard deviation of mean change for both study groups
Dichotomous outcomes		
Asthma control	Mehuys <i>et al</i> [22]	Appropriate for meta-analysis
	Armour <i>et al</i> [15]	Appropriate for meta-analysis
	Garcia-Cardenas <i>et al</i> [19]	Appropriate for meta-analysis
	Zanghelini <i>et al</i> [27]	Lack of control group
Asthma severity	Armour <i>et al</i> (2007)[14]	Appropriate for meta-analysis

ACQ: Asthma Control Questionnaire; ACT: Asthma Control Test; PEF: Peak Expiratory Flow; FEV1: Forced Expiratory Volume in the first second; FVC: Forced Vital Capacity; VC: Vital Capacity

## DISCUSSION

This systematic review identified twenty-one studies assessing the impact of pharmacists' interventions on asthma outcomes in adult patients. A large variety of outcomes were used to demonstrate such impact, including different measures of asthma control, asthma severity, pulmonary function and asthma symptoms.

RCTs/C-RCTs generated similar results to those generated by non-RCT for most of the outcomes assessed. Nine out of ten studies assessing asthma control revealed a positive impact after the provision of the pharmacist's intervention. For example, all studies reporting changes in the percentage of controlled asthma patients, found a positive association between pharmacist's intervention and current asthma control [15, 19, 22]. Similar results were observed in terms of ACQ scores, with all six studies reporting significant improvements for this outcome [15, 19, 20, 25, 26, 33]. Two of the four studies measuring ACT scores found a similar trend [17, 27]. Studies assessing any measure of asthma severity also supported the potential role of pharmacists on asthma management. Pharmacists contributed to a significant reduction in the percentage of severe patients and asthma severity scores in three of the four studies assessing this outcome [14, 21, 23, 24]. Both RCT and non-RCT assessing pulmonary function measures, failed to demonstrate a significant impact on VC, FVC, FEV1/FVC and FEV1%VC. However, improvements were identified in terms of PEF and FEV1 [18, 21, 23, 27, 28, 30, 32]. Regardless of the study design, pharmacists were found to have a positive impact on different measures of asthma symptoms. For example, all seven studies assessing general or nocturnal symptoms as their main outcome reported improvements after the provision of the intervention [16, 18, 21, 22, 29-31]. It is evident that both RCTs/C-RCTs and non-RCT yielded similar trends, highlighting the potential role of pharmacists in asthma management. However, appropriately designed, conducted and reported RCTs/C-RCTs represent the gold standard in assessing healthcare interventions [36]. Future research aimed at assessing pharmacists' interventions on asthma outcomes should utilize these research designs. Nevertheless, further experimental designs can be considered if a conventional RCT design is not feasible [37]. The use of other research designs would also be of great interest, allowing the assessment of the interventions' elements. It is worthwhile mentioning that amongst the included studies, many different interventions with several interacting components were tested. This heterogeneity may have impacted the outcomes achieved, and may account for the differences between significant and non-significant findings. As the effect of the different intervention elements was not independently assessed in any of the studies, those with the biggest impact on clinical asthma outcomes remain unknown.

Within the twenty-one included studies, eleven different clinical outcomes and twenty-six different reporting systems were identified, hindering the synthesis of the evidence available. As in other conditions [38, 39], it is necessary to agree on a core outcome set to be measured and reported in all the studies assessing the impact of health interventions on adult asthma patients. The selection of appropriate outcomes to measure allows the analysis of the effects of different interventions and minimises bias[10]. This provides not only a reliable comparison of results across different studies, but also an evaluation of the consistency of the research findings when translated into clinical practice. The core outcome set for asthma interventions must be established in accordance with updated evidence based reports, such as the Global Initiative for Asthma (GINA). In its latter update, GINA recommends the assessment of asthma control based on two different domains: symptom control (previously known as current clinical control) and future risk of adverse outcomes (including risk factors for exacerbations, fixed airflow limitation and medication side-effects)[3]. This implies that both domains should always be assessed and reported separately. The application of this approach to assess the impact of pharmacists' interventions seems to be feasible. We suggest that studies assessing pharmacist interventions in clinical trials always report at least symptom control and specific indicators of risk of adverse outcomes as a core outcome set, described as follows. The assessment of symptoms control should be done using validated tools, such as the ACQ and the ACT. These instruments have already been used in several intervention studies with positive results[15, 17, 19, 20, 25-27, 33, 34]. In terms of a future risk of adverse outcomes assessment; at least short-acting beta2-agonist use, adherence to inhaled corticosteroids, inhaler technique, exposure to triggers, and potentially FEV1, should be measured and reported. None of the studies included in this review reported all the aforementioned outcomes. This might be explained by the fact that first asthma severity, and then current clinical control, have traditionally been the ultimate outcomes of asthma management.

A meta-analysis of the studies was attempted to estimate the pooled effects of pharmacists' interventions on asthma outcomes. However, some factors impeded its performance, such as the variability on the different outcomes assessed and the reporting systems used, together with lack of statistical parameters. Different guidelines for reporting experimental studies such as CONSORT[35], have been developed to assist authors in writing manuscripts, and journal editors and peer reviewers in evaluating them for publication. This has undoubtedly helped to increase the quality of data available in the scientific community. However, whilst key statistical parameters are missing or high variability remains common, identifying a real effect of an intervention will not be possible.

A potential limitation of this systematic review is that some studies undertaken in the early 2000's, assessing the role of pharmacists on asthma management were retrieved in our search strategy but not included in the analysis[40-42]. These papers included children as well as adults, and we reviewed studies which dealt with adults only. Although our query intended to retrieve all published studies evaluating the effect of pharmacists' interventions in clinical asthma outcomes, publication bias may have occurred because no grey literature was included. However, since no papers were retrieved from additional references, it seems that high-sensitivity queries were used. This may also be explained due to the number of databases used, covering most of the published data in our research area. Due to the low number of RCTs/C-RCTs identified, studies with different experimental research designs were included. This impeded performing a risk of bias assessment. However, following AMSTAR recommendations, this was taken into account for the data synthesis and formulation of the conclusions of this review[13]. Moreover, the potential differing quality of the studies included did not affect the main objective of our review.

In conclusion, the evidence of pharmacists' interventions on clinical asthma outcomes in adult patients has been evaluated using heterogeneous outcomes, including different measures of asthma control, asthma severity, pulmonary function, and asthma symptoms. Based on the evidence generated by RCTs/C-RCT, pharmacists' interventions have a positive impact on the percentage of asthma-controlled patients, ACQ scores, asthma severity and asthma symptoms. Inconsistent impact has been found in terms of ACT scores and pulmonary function indicators. Future research evaluating the impact of pharmacists' interventions on clinical asthma outcomes should report using a core outcome set of indicators established for this condition, based on GINA recommendations.

#### **COMPETING INTERESTS**

None.

#### **FUNDING**

None.

## REFERENCE LIST

1. World Health Organisation (WHO). Asthma fact sheet 307 [Internet]. World Health Organisation; 2013 Nov [cited 2015 Jan 22]. Available from: <http://www.who.int/mediacentre/factsheets/fs307/en/>.
2. Braman SS. The global burden of asthma. *Chest* 2006;130(1 Suppl):4s-12s.
3. Global Initiative for asthma (GINA). Global Strategy for Asthma Management and Prevention; 2015 [cited 2015 Apr 5]. 148p. Available from: <http://www.ginasthma.org>.
4. Haughney J, Price D, Kaplan A, Chrystyn H, Horne R, May N, et al. Achieving asthma control in practice: understanding the reasons for poor control. *Respir Med* 2008;102(12):1681-93.
5. Horne R, Price D, Cleland J, Costa R, Covey D, Gruffydd-Jones K, et al. Can asthma control be improved by understanding the patient's perspective? *BMC Pulm Med* 2007;7:8.
6. Machado M, Bajcar J, Guzzo GC, Einarson TR. Sensitivity of patient outcomes to pharmacist interventions. Part I: systematic review and meta-analysis in diabetes management. *Ann Pharmacother* 2007;41(10):1569-82.
7. Machado M, Bajcar J, Guzzo GC, Einarson TR. Sensitivity of patient outcomes to pharmacist interventions. Part II: Systematic review and meta-analysis in hypertension management. *Ann Pharmacother* 2007;41(11):1770-81.
8. Fathima M, Naik-Panvelkar P, Saini B, Armour CL. The role of community pharmacists in screening and subsequent management of chronic respiratory diseases: a systematic review. *Pharm Pract (Granada)* 2013;11(4):228-45.
9. Benavides S, Rodriguez JC, Maniscalco-Feichtl M. Pharmacist involvement in improving asthma outcomes in various healthcare settings: 1997 to present. *Ann Pharmacother* 2009;43(1):85-97.
10. Williamson PR, Altman DG, Blazeby JM, Clarke M, Devane D, Gargon E, et al. Developing core outcome sets for clinical trials: issues to consider. *Trials* 2012;13:132.
11. Gargon E, Williamson PR, Altman DG, Blazeby JM, Clarke M. The COMET Initiative database: progress and activities from 2011 to 2013. *Trials* 2014;15:279.
12. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med* 2009;6(7):e1000100.
13. Shea BJ, Grimshaw JM, Wells GA, Boers M, Andersson N, Hamel C, et al. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol* 2007;7:10.
14. Armour C, Bosnic-Anticevich S, Brilliant M, Burton D, Emmerton L, Krass I, et al. Pharmacy Asthma Care Program (PACP) improves outcomes for patients in the community. *Thorax* 2007;62(6):496-502.
15. Armour CL, Reddel HK, LeMay KS, Saini B, Smith LD, Bosnic-Anticevich SZ, et al. Feasibility and effectiveness of an evidence-based asthma service in Australian community pharmacies: a pragmatic cluster randomized trial. *J Asthma* 2013;50(3):302-9.



16. Barbanel D, Eldridge S, Griffiths C. Can a self-management programme delivered by a community pharmacist improve asthma control? A randomised trial. *Thorax* 2003;58(10):851-4.
17. Bereznicki BJ, Peterson GM, Jackson SL, Walters H, Fitzmaurice K, Gee P. Pharmacist-initiated general practitioner referral of patients with suboptimal asthma management. *Pharm World Sci* 2008;30(6):869-75.
18. Petkova E. Evaluation of the impact of a pharmaceutical care program on patients with asthma. *Ankara Universitesi Eczacilik Fakultesi Dergisi* 2005;34(4):251-62.
19. Garcia-Cardenas V, Sabater-Hernandez D, Kenny P, Martinez-Martinez F, Faus MJ, Benrimoj SI. Effect of a pharmacist intervention on asthma control. A cluster randomised trial. *Respir Med* 2013;107(9):1346-55.
20. Giraud V, Allaert FA, Roche N. Inhaler technique and asthma: feasibility and acceptability of training by pharmacists. *Respir Med* 2011;105(12):1815-22.
21. Mangiapane S, Schulz M, Muhlig S, Ihle P, Schubert I, Waldmann HC. Community pharmacy-based pharmaceutical care for asthma patients. *Ann Pharmacother* 2005;39(11):1817-22.
22. Mehuys E, Van Bortel L, De Bolle L, Van Tongelen I, Annemans L, Remon JP, et al. Effectiveness of pharmacist intervention for asthma control improvement. *Eur Respir J* 2008;31(4):790-9.
23. Saini B, Krass I, Armour C. Development, implementation, and evaluation of a community pharmacy-based asthma care model. *Ann Pharmacother* 2004;38(11):1954-60.
24. Schulz M, Verheyen F, Muhlig S, Muller JM, Muhlbauer K, Knop-Schneickert E, et al. Pharmaceutical care services for asthma patients: a controlled intervention study. *J Clin Pharmacol* 2001;41(6):668-76.
25. Smith L, Bosnic-Anticevich SZ, Mitchell B, Saini B, Krass I, Armour C. Treating asthma with a self-management model of illness behaviour in an Australian community pharmacy setting. *Soc Sci Med* 2007;64(7):1501-11.
26. Toumas-Shehata M, Price D, Basheti IA, Bosnic-Anticevich S. Exploring the role of quantitative feedback in inhaler technique education: a cluster-randomised, two-arm, parallel-group, repeated-measures study. *NPJ Prim Care Respir Med* 2014;24:14071.
27. Zanghelini F, Filho J, Carvalho A, da Silva A. Impact of pharmaceutical care on lung function in patients with severe asthma. *Revista de Ciencias Farmaceuticas Basica e Aplicada* 2013;34(3):379-86.
28. Anjan Kumar D, Adepu R, Parthasarathi G, Mahesh P. Impact of Community Pharmacist Provided Patient Education in Asthma Patients on Treatment Outcomes- A Study. *Indian J Educ Res* 2009;43(2):125-33.
29. Abdelhamid E, Awad A, Gismallah A. Evaluation of a hospital pharmacy-based pharmaceutical care services for asthma patients. *Pharmacy Practice* 2008;6(1):25-32.
30. Narhi U, Airaksinen M, Tanskanen P, Erlund H. Therapeutic outcomes monitoring by community pharmacists for improving clinical outcomes in asthma. *J Clin Pharm Ther* 2000;25(3):177-83.

31. Odegard PS, Lam A, Chun A, Blough D, Li MA, Wu J. Pharmacist provision of language-appropriate education for Asian patients with asthma. *J Am Pharm Assoc* (2003) 2004;44(4):472-7.
32. De Tullio PL, Corson ME. Effect of pharmacist counseling on ambulatory patients' use of aerosolized bronchodilators. *Am J Hosp Pharm* 1987;44(8):1802-6.
33. Lim AS, Stewart K, Abramson MJ, Walker SP, Smith CL, George J. Multidisciplinary Approach to Management of Maternal Asthma (MAMMA): a randomized controlled trial. *Chest* 2014;145(5):1046-54.
34. Young HN, Havican SN, Griesbach S, Thorpe JM, Chewing BA, Sorkness CA. Patient and pharmacist telephonic encounters (PARTE) in an underserved rural patient population with asthma: results of a pilot study. *Telemed J E Health* 2012;18(6):427-33.
35. Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *Bmj* 2010;340:c332.
36. Sedgwick P. What is a non-randomised controlled trial? *Bmj* 2014;348:g4115.
37. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *Bmj* 2008;337:a1655.
38. Benstoem C, Moza A, Autschbach R, Stoppe C, Goetzenich A. Evaluating outcomes used in cardiothoracic surgery interventional research: a systematic review of reviews to develop a core outcome set. *PLoS One* 2015;10(4):e0122204.
39. Sinha IP, Gallagher R, Williamson PR, Smyth RL. Development of a core outcome set for clinical trials in childhood asthma: a survey of clinicians, parents, and young people. *Trials* 2012;13:103.
40. Cordina M, McElnay JC, Hughes CM. Assessment of a community pharmacy-based program for patients with asthma. *Pharmacotherapy* 2001;21(10):1196-203.
41. Herborg H, Soendergaard B, Froekjaer B, Fonnesbaek L, Jorgensen T, Hepler CD, et al. Improving drug therapy for patients with asthma--part 1: Patient outcomes. *J Am Pharm Assoc (Wash)* 2001;41(4):539-50.
42. McLean W, Gillis J, Waller R. The BC Community Pharmacy Asthma Study: A study of clinical, economic and holistic outcomes influenced by an asthma care protocol provided by specially trained community pharmacists in British Columbia. *Can Respir J* 2003;10(4):195-202.