

ON SOME RECIPROCAL SUMS OF BROUSSEAU: AN ALTERNATIVE APPROACH TO THAT OF CARLITZ

R. S. Melham

Department of Math. Sciences, University of Technology, Sydney

PO Box 123, Broadway, NSW 2007, Australia

(Submitted November 2000)

1. INTRODUCTION

In [2], it was shown that

$$\sum_{n=1}^{\infty} \frac{1}{F_n} = 3 + \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{F_n F_{n+1} F_{n+2}}, \quad (1.1)$$

and, using the same approach, it can be shown that

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{F_n} = 1 - \sum_{n=1}^{\infty} \frac{1}{F_n F_{n+1} F_{n+2}}. \quad (1.2)$$

Let m be a positive integer, and define the sums

$$S(1, \dots, m) = \sum_{n=1}^{\infty} \frac{1}{F_n F_{n+1} \cdots F_{n+m}}, \quad m \geq 1, \quad (1.3)$$

and

$$T(1, \dots, m) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{F_n F_{n+1} \cdots F_{n+m}}, \quad m \geq 1. \quad (1.4)$$

In [1] (see equations 20, 21, and 22), Brousseau proved that

$$T(1, 2) = \frac{5}{12} - \frac{3}{2} S(1, 2, 3, 4), \quad (1.5)$$

$$S(1, 2, 3, 4) = \frac{97}{2640} + \frac{40}{11} T(1, 2, 3, 4, 5, 6), \quad (1.6)$$

and

$$T(1, 2, 3, 4, 5, 6) = \frac{589}{1900080} - \frac{273}{29} S(1, 2, 3, 4, 5, 6, 7, 8). \quad (1.7)$$

As an application, he computed the value of the sum $\sum_{n=1}^{\infty} \frac{1}{F_n}$ to twenty-five decimal places.

Our aim in this paper is to establish explicit formulas that extend (1.5)-(1.7). Specifically, we obtain formulas

$$T(1, \dots, m) = r_1 + r_2 S(1, \dots, m, m+1, m+2), \quad m \geq 1, \quad (1.8)$$

and

$$S(1, \dots, m) = r_3 + r_4 T(1, \dots, m, m+1, m+2), \quad m \geq 1, \quad (1.9)$$

where the r_i are rational numbers that depend on m . Among other things, Carlitz [3] attacked the same problem with the use of generating functions and Fibonomial coefficients. Here we provide an alternative and more transparent approach, with the use of only simple identities that involve the Fibonacci numbers.

2. PRELIMINARY RESULTS

We require the following results:

$$\sum_{n=1}^{\infty} \frac{1}{F_n F_{n+2}} = 1; \tag{2.1}$$

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{F_n F_{n+2}} = -1 + 2T(1); \tag{2.2}$$

$$\sum_{n=1}^{\infty} \frac{1}{F_n F_{n+1} F_{n+3}} = -\frac{1}{4} + S(1, 2); \tag{2.3}$$

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{F_n F_{n+1} F_{n+3}} = \frac{1}{4}; \tag{2.4}$$

$$\sum_{n=1}^{\infty} \frac{1}{F_n F_{n+1} \cdots F_{n+m-1} F_{n+m+1}} = \frac{(-1)^{m-1}}{F_1 \cdots F_m F_{m+1}^2} + \frac{(-1)^m + F_m}{F_{m+1}} S(1, 2, \dots, m), \quad m \geq 1; \tag{2.5}$$

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{F_n F_{n+1} \cdots F_{n+m-1} F_{n+m+1}} = \frac{(-1)^m}{F_1 \cdots F_m F_{m+1}^2} + \frac{(-1)^{m-1} + F_m}{F_{m+1}} T(1, 2, \dots, m), \quad m \geq 1; \tag{2.6}$$

$$\begin{aligned} & F_{n+m} F_{n+m+2} - F_n F_{n+m+1} \\ &= (-1)^{m-1} F_{m+1} F_{n+m+1} F_{n+m+2} + [1 + (-1)^m F_{m+2}] F_{n+m} F_{n+m+2} - (-1)^{n-1} F_{m+2}; \end{aligned} \tag{2.7}$$

$$\begin{aligned} & F_{n+m} F_{n+m+2} + F_n F_{n+m+1} \\ &= (-1)^m F_{m+1} F_{n+m+1} F_{n+m+2} + [1 + (-1)^{m-1} F_{m+2}] F_{n+m} F_{n+m+2} + (-1)^{n-1} F_{m+2}. \end{aligned} \tag{2.8}$$

Formulas (2.1)-(2.3) can be obtained from [1]. More precisely: (2.1) occurs as (4); (2.2) follows if we use (3) to evaluate

$$\sum_{n=1}^{\infty} \frac{1}{F_n F_{n+k}}$$

for $k = 1$ and $k = 2$; and (2.3) is the first entry in Table III. Formula (2.4) follows from (2.4) in [4]. Again, turning to [4], we see that identity (3.3) therein and its counterpart for T yield (2.5) and (2.6) for $m \geq 3$. We can verify the validity of (2.5) and (2.6) for $m = 1$ and 2 by simply substituting these values and comparing the outcomes with (2.1)-(2.4). Finally, (2.7) and (2.8) can be established with the use of the Binet form for F_n .

3. THE RESULTS

Our results are contained in the theorem that follows.

Theorem: Let $m \geq 1$ be an integer. Then

$$S(1, \dots, m) = \frac{1}{(1 + (-1)^{m-1} - L_{m+1})} \left[\frac{(-1)^{m-1} F_{m+2} - F_{2m+3}}{F_1 \cdots F_{m+2}} - F_{m+1} F_{m+2} T(1, \dots, m+2) \right], \tag{3.1}$$

$$T(1, \dots, m) = \frac{1}{(1 + (-1)^{m-1} + L_{m+1})} \left[\frac{(-1)^{m-1} F_{m+2} + F_{2m+3}}{F_1 \cdots F_{m+2}} - F_{m+1} F_{m+2} S(1, \dots, m+2) \right]. \tag{3.2}$$

Proof: Let $m \geq 1$ be an integer. Then, due to telescoping, we have

$$\sum_{n=1}^{\infty} \left(\frac{1}{F_n \cdots F_{n+m-1} F_{n+m+1}} - \frac{1}{F_{n+1} \cdots F_{n+m} F_{n+m+2}} \right) = \frac{1}{F_1 \cdots F_m F_{m+2}}. \tag{3.3}$$

Alternatively, with the use of (2.7), this sum can be written as

$$\begin{aligned} & \sum_{n=1}^{\infty} \frac{F_{n+m} F_{n+m+2} - F_n F_{n+m+1}}{F_n \cdots F_{n+m+2}} \\ &= \sum_{n=1}^{\infty} \frac{(-1)^{m-1} F_{m+1} F_{n+m+1} F_{n+m+2} + [1 + (-1)^m F_{m+2}] F_{n+m} F_{n+m+2} - (-1)^{n-1} F_{m+2}}{F_n \cdots F_{n+m+2}} \\ &= (-1)^{m-1} F_{m+1} S(1, \dots, m) + [1 + (-1)^m F_{m+2}] \sum_{n=1}^{\infty} \frac{1}{F_n \cdots F_{n+m-1} F_{n+m+1}} - F_{m+2} T(1, \dots, m+2). \end{aligned} \tag{3.4}$$

Finally, after using (2.5) to substitute for

$$\sum_{n=1}^{\infty} \frac{1}{F_n F_{n+1} \cdots F_{n+m-1} F_{n+m+1}},$$

we equate the right sides of (3.3) and (3.4), and then solve for $S(1, \dots, m)$ to obtain (3.1). In the course of the algebraic manipulations, we make use of the well-known identities $L_n = F_{n-1} + F_{n+1}$, $F_n^2 + F_{n+1}^2 = F_{2n+1}$, and $F_{n-1} F_{n+1} - F_n^2 = (-1)^n$.

Since the proof of (3.2) is similar, we merely give an outline. To begin, we have

$$\sum_{n=1}^{\infty} (-1)^{n-1} \left(\frac{1}{F_n \cdots F_{n+m-1} F_{n+m+1}} + \frac{1}{F_{n+1} \cdots F_{n+m} F_{n+m+2}} \right) = \frac{1}{F_1 \cdots F_m F_{m+2}}. \tag{3.5}$$

Next, we write the left side of (3.5) as

$$\sum_{n=1}^{\infty} (-1)^{n-1} \left(\frac{F_{n+m} F_{n+m+2} + F_n F_{n+m+1}}{F_n \cdots F_{n+m+2}} \right). \tag{3.6}$$

Finally, we make use of (2.8) and (2.6), and then solve for $T(1, \dots, m)$. This completes the proof of the theorem. \square

If we substitute $m = 4$ into (3.1), we obtain (1.6). Likewise, if we substitute $m = 2$ and $m = 6$ into (3.2), we obtain (1.5) and (1.7), respectively.

4. CONCLUDING COMMENTS

Our results (3.1) and (3.2) do not produce (1.1) and (1.2), which, as stated in the Introduction, can be arrived at independently. Interestingly, due to our alternative approach, our main results are more simply stated than the corresponding results in [3]. See, for example, (5.8) in [3]. Incidentally, there is a typographical error in the last formula on page 464, where $-\frac{2}{3}$ should be replaced by $\frac{2}{3}$.

Finally, we refer the interested reader to the recent paper [5], where Rabinowitz discusses algorithmic aspects of certain finite reciprocal sums.

REFERENCES

1. Brother Alfred Brousseau. "Summation of Infinite Fibonacci Series." *The Fibonacci Quarterly* **7.2** (1969):143-68.
2. L. Carlitz. Solution to Problem H-10 (proposed by R. L. Graham). *The Fibonacci Quarterly* **1.4** (1963):49.
3. L. Carlitz. "Reduction Formulas for Fibonacci Summations." *The Fibonacci Quarterly* **9.5** (1971):449-66, 510-11.
4. R. S. Melham. "Reduction Formulas for the Summation of Reciprocals in Certain Second-order Recurring Sequences." *The Fibonacci Quarterly* **40.1** (2002):71-75.
5. Stanley Rabinowitz. "Algorithmic Summation of Reciprocals of Products of Fibonacci Numbers." *The Fibonacci Quarterly* **37.2** (1999):122-27.

AMS Classification Numbers: 11B39, 40A99





The Fibonacci Quarterly

THE OFFICIAL JOURNAL OF THE FIBONACCI ASSOCIATION

TABLE OF CONTENTS

Referees	2
The Interval Associated with a Fibonacci Number <i>Takao Komatsu</i>	3
Some Identities Involving the Powers of the Generalized Fibonacci Number <i>Feng-Zhen Zhao and Tianming Wang</i>	7
Complex Factorizations of the Fibonacci and Lucas Numbers <i>Nathan D. Cahill, John R. D'Errico, and John P. Spence</i>	13
Inequalities among Related Pairs of Fibonacci Numbers <i>K.T. Atanassov, R. Knott, K. Ozeki, A.G. Shannon, and L. Szalay</i>	20
A Summation Formula for Power Series Using Eulerian Fractions <i>Xinghua Wang and Leetsch C. Hsu</i>	23
q -Fibonacci Polynomials	<i>Johann Cigler</i> 31
On Lacunary Recurrences	<i>Paul Thomas Young</i> 41
Applications of Matrix Theory to Congruence Properties of k^{th} -Order F-L Sequences	<i>Chizhong Zhou</i> 48
On Some Reciprocal Sums of Brousseau: An Alternative Approach to That of Carlitz	<i>R.S. Melham</i> 59
Some Fractals in Goldpoint Geometry	<i>J.C. Turner</i> 63
Divisibility Properties by Multisection	<i>Tamás Lengyel</i> 72
The Sum of the Squares of Two Generalized Fibonacci Numbers	<i>F.T. Howard</i> 80
Elementary Problems and Solutions	<i>Edited by Russ Euler and Jawad Sadek</i> 85
Advanced Problems and Solutions	<i>Edited by Raymond E. Whitney</i> 91

VOLUME 41

FEBRUARY 2003

NUMBER 1

Find solutions for:

- ACADEMIC
- GOVERNMENT
- NON-PROFIT
- CORPORATE

- PRODUCTS
- CONFERENCES AND EVENTS

- TRAINING
- SUPPORT
- RESEARCH SERVICES
- JOURNAL LISTS
- JOURNAL SELECTION PROCESS
- ISI LINKS
- ISI ESSAYS
- HOT RESEARCH

- REGIONS

Business Website GO

SEARCH RESULTS

ISI Master Journal List
SEARCH RESULTS

Search Terms: 0015-0517
Total journals found: 1

The following title(s) matched your request:

Journals 1-1 (of 1) [Navigation icons]

FORMAT FOR PRINT

FIBONACCI QUARTERLY

Quarterly
ISSN: 0015-0517
FIBONACCI ASSOC, C/O PATTY SOLSAA, PO BOX 320, AURORA, USA, SD, 57002-0320

Journals 1-1 (of 1) [Navigation icons]

FORMAT FOR PRINT