

# Training Interventions for Improved Deceleration Ability in Adult Team-Based Field Sports Athletes: A Systematic Review and Meta-Analysis of the Literature

Louise Amy Marvin<sup>1,2</sup>, Paul Read<sup>3,4,5,7</sup>, Blake McLean<sup>1,2</sup>, Sam Palmer<sup>1,2</sup> & Job Fransen<sup>6</sup>

<sup>1</sup>School of Sport, Exercise & Rehabilitation, Faculty of Health, University of Technology Sydney (UTS), Moore Park, NSW, Australia, <sup>2</sup>Human Performance Research Centre, University of Technology Sydney (UTS), Moore Park, NSW, Australia, <sup>3</sup>Institute of Sport Exercise and Health, London, UK, <sup>4</sup>Division of Surgery & Interventional Science, University College London, UK, <sup>5</sup>School of Sport and Exercise Sciences, University of Gloucestershire, UK, <sup>6</sup>Charles Sturt University, Port Macquarie, NSW, Australia, <sup>7</sup>Faculty of Sport, Allied Health and Performance Science, St Mary's University, London, United Kingdom

## ABSTRACT

**Background:** Deceleration is a fundamental component of multidirectional speed by which athletes reduce the velocity of their centre of mass to stop or execute changes of direction following acceleration or running at a constant velocity. Enhancing deceleration abilities is crucial for athletes as successfully executing horizontal deceleration has important implications for match outcomes in sports requiring rapid multidirectional movements. However, specific training interventions targeting deceleration are scarce. The purpose of this systematic review and meta-analysis was to examine the effects of training interventions on deceleration performance in adult team-based field and court sports athletes.

**Methods:** A systematic literature search was conducted through electronic databases, SPORTdiscus, PubMed, and Web of Science from inception to February 2022, and re-run in May

2023. The search terms were related to different training interventions and kinetic, kinematic, and performance outcomes related to deceleration performance. Studies were included if they consisted of a randomised controlled trial which investigated the effects of training on deceleration-specific outcome measures in adult team-based field and court sports athletes. Risk of bias was assessed using the revised Cochrane risk-of-bias tool for randomised trials (RoB2). Post-intervention effect sizes (Hedge's *g*) were calculated between the intervention and control groups and a meta-analysis was performed using a random effects model.

**Results:** Twelve studies were included, with 29 deceleration-specific outcomes measured in a total of 381 participants. There was inconsistency in methodological designs, including control group types, length and type of interventions and in reported deceleration-specific outcome measures. Across all observations of deceleration performance

measures there was a standardised mean difference of -0.04 (95% CI: -0.50, 0.42), favouring control groups, indicating little effect of training on deceleration performance. For secondary outcomes related to deceleration, for kinetics (SMD = -0.29, 95% CI = -0.83, 0.25,  $I^2$  = 55%,  $p$  < 0.01), joint kinetics (SMD = -0.20, 95% CI = -1.01, 0.61,  $I^2$  = 68%,  $p$  < 0.01) and muscle activation (SMD = -0.10, 95% CI = -0.44, 0.24,  $I^2$  = 19%,  $p$  = 0.28) had greater effects for control groups, whereas joint kinematics (SMD = 0.07, 95% CI = -0.11, 0.24,  $I^2$  = 0%,  $p$  = 0.77), favoured the intervention groups.

**Conclusion:** For the deceleration-specific outcomes reported in the included studies, training was not likely to produce a performance improvement in participants compared to control groups. However, due to methodological inconsistencies between studies and observed high risk of bias, the results should be interpreted with caution. More rigorous research methods should be included in the future to address areas that may introduce potential biases. Future research should address the differences in the type, timing, frequency, and duration of the implemented training interventions for improving deceleration performance, and in the reported deceleration-specific outcome measures.

**Registration:** This systematic review was registered on the Open Science Framework (<https://osf.io/cmwwr>) (<https://doi.org/10.17605/OSF.IO/CMWBR>)

**Keywords:** Deceleration, systematic review, meta-analysis, athletic performance, training, field sports, team-based sports

## INTRODUCTION

Deceleration is an essential component of multidirectional speed whereby an athlete slows down their centre of mass following acceleration or running at a constant velocity to come to a stop, slow down, and/or change direction [1, 2]. In sports, such as soccer, basketball, and rugby, frequent and intense accelerations and decelerations are crucial elements of competitive match play and have an important impact on performance outcomes and match results [3, 4].

While considerable research has investigated the trainability, physiology, and influence on performance of linear speed, acceleration, change-of-direction (COD) and agility [5-10], deceleration is rarely studied as an explicit skill [4]. However,

recent research has highlighted the importance of horizontal deceleration [11] as this ability enables athletes to reduce momentum during very short time frames and distances to successfully evade or pursue opponents. Therefore, targeting deceleration capacities and motor skills [12] with effective training methods is warranted [4].

To-date intervention trials aimed at enhancing deceleration performance have not been widely examined [9]. Few studies have included deceleration-specific outcome measures or addressed how deceleration occurs in response to sports specific stimuli [4, 12, 13]. Most research has included indirect measures of deceleration performance. These have included the coordination patterns employed by the performer during the final steps to reduce velocity prior to coming to a complete stop. Additionally analyses of the preparatory phase, initiated 50 milliseconds before ground contact, and the ensuing loading phase, defined by the interaction of the foot with the ground during the final foot contact of a COD, which requires momentarily coming to zero velocity, have been used [8]. Outcome measures have included joint kinetics and kinematics of the hip, knee, and ankle; kinetics; and muscle activation patterns measured by surface electromyography (EMG). Deterministic models have identified predictors of deceleration ability, including eccentric strength, power, and reactive strength of the lower limbs [3]. Therefore enhancing these physical capacities may be beneficial to improve attributes associated with executing the skill. Additionally, athletes could benefit from being exposed to challenges which enhance the perceptual-cognitive skills essential for successful unplanned braking [4] emphasising the cognitive aspect of trainability. Acknowledging the complementary nature of both approaches, it is essential to recognise that training for effective deceleration performance involves a combination of improving physical capacities and refining perceptual-cognitive skills. However, despite these theorised avenues to improve deceleration performance, there is no clear evidence for the effectiveness of such training methods for specifically improving deceleration ability.

Therefore, the aim of this systematic review and meta-analysis is to critically evaluate and summarise the results of studies examining the effectiveness of different training interventions to specifically improve deceleration ability in adult team-based field and court sports athletes. The review also aims to appraise the methodological

approach of the included studies and determine the effectiveness of training interventions compared to normal or “as usual” training practices commonly employed by the respective sports. By examining interventions against established training norms, this review seeks to provide insights into the specific impact of targeted interventions on deceleration performance. The findings of our review aim to contribute to the development of evidence-based training recommendations to enhance deceleration performance and reduced injury risk in team-based field and court sports athletes.

## METHODS

The systematic review was performed according to the guidelines for the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [14] and the protocol registered with the Open Science Framework (<https://doi.org/10.17605/OSF.IO/CMWBR>, 16 February 2022).

### Searches

A comprehensive search of the electronic bibliographic databases SPORTdiscus, PubMed, and Web of Science was conducted in February 2022 by one author (LM). The search strategy is outlined in Figure 1. No restrictions were placed on publication date or language

The keywords used in the database search were:

- “Team sport\*\*” OR team-sport\* OR “multi-direction\*\*” OR multidirection\* OR “field sport\*\*” OR “field-based sport\*\*” OR “intermittent sport\*\*” OR soccer OR football\* OR futsal OR rugby OR “American football” OR “Australian rules football” OR “Australian football” OR AFL OR basketball OR handball OR netball OR hockey OR “invasion sports” OR lacrosse OR hurling OR “Gaelic football”

AND

- Train\* OR interven\* OR exercis\* OR “plyometric training” OR plyometrics OR “power training” OR “strength training” OR “resistance training” OR “speed training” OR practic\* OR practis\* OR program\* OR “agility training”

AND

- Accelerat\* OR decelerat\* OR agility OR

“change of direction” OR COD OR “side step” OR side-step OR reactive OR unplanned OR unanticipated OR manoeuvre\* OR maneuver\* OR cutting OR brak\* OR “negative accelerat\*” OR turn\* OR stop\*

The reference lists of all included articles were then manually searched to identify additional articles for inclusion in the review. Citation chaining was performed using Google Scholar and Web of Science. A revised search was completed in the same databases up until June 2023 and eligible studies included for data extraction.

### Eligibility Criteria

The eligibility criteria were specified following a scoping search of relevant studies. The included studies met the following population, intervention, comparator, outcome, and study design (PICOS) criteria.

All original journal articles identified in the initial search were included in this review if they met the following criteria:

- Population: trained adult ( $\geq 18$  years) male and female athletes from any team-based field and court sports. Team-based field and court sports were defined as team sports played on a court or field as a supporting surface such as volleyball, basketball, or Australian football.
- Intervention: any training program (e.g. strength, plyometrics, agility) that was not part of participants’ standard or as usual training.
- Comparator: studies had to compare to either a passive control condition (no training or as usual training) or active control condition (performing a different type of training that would be considered standard in the field) (examples of such training include traditional resistance training, or sprint and agility training).
- Outcome: a kinetic, kinematic, or performance-related measure specifically measuring deceleration ability, assessed pre- and post-intervention. Examples of such measures are peak and average deceleration, joint angles and moments, ground reaction forces (GRFs), etc during the final foot contact of a COD or deceleration.
- Study design: only randomised controlled trials (RCT) with a pre- and post-test design were considered.

The population of trained adult male and female team sports athletes was chosen to limit possible

confounding effects of age and training status on deceleration ability that may be evident with the inclusion of children/youth athletes. Trained athletes referred to anyone with a history of regular participation in training and/or competing in their chosen sport, indicating relevant skill development and familiarity with the demands of their respective sports. Sex, age, and sport were not specified as it was necessary to gather a wide scope of data of a range of training interventions aimed at improving deceleration ability in all team-based field and court sports where deceleration is important for performance.

Studies were excluded from the review if they were not an RCT, did not have a control group, or if measures of deceleration were not reported as primary or secondary outcomes. Further exclusionary criteria were letters, reviews, and books. Studies from grey literature or those published in languages other than English were not excluded from the analysis.

### Study Selection

All articles retrieved through the search were exported into the web-based software platform Covidence (Veritas Health Innovation, <https://www.covidence.org/>) where all duplicates were subsequently removed. Two authors (LM & SP) independently screened titles and abstracts for records identified during the initial search strategy. Studies included for full text review were then independently screened by the same two reviewers for extraction. At each stage of the study selection and extraction process the reviewers were blinded to each other's decisions. In the case of disagreement, decisions were made through discussion, and resolved by input from an additional independent screener (JF). In both screening stages, Cohen's kappa statistics were used to calculate the inter-rater agreement for study inclusion.

### Data Extraction

Data were extracted and collated by one author (LM) using a customised excel spreadsheet (Online Resource 1). The information extracted from each eligible study included the publication details, population characteristics (number of participants, age (mean  $\pm$  standard deviation [SD]), sex, sport), duration, frequency, and type of intervention and comparison, reported deceleration-specific outcome measures, statistical analyses used, and results. Descriptive statistics, including mean

and standard deviations for outcome measures, confidence intervals, and effect sizes were extracted and reported. If the pre- and post-intervention test data were displayed in figures, or data were not provided within the manuscript [15-18] the corresponding authors were contacted via email for further information. Where authors were unable to provide data [17], or did not respond [15, 16] these studies were not included in the final meta-analysis. All eligible effect sizes from each study were extracted, resulting in more than one effect size per study.

In the analysis of reported measures of deceleration, five main dependent variables were considered separately: performance outcomes, kinetics, joint kinetics, joint kinematics, and muscle activation patterns. Across the reported outcomes, some outcomes are positively scored (i.e., higher values are related to better deceleration performance) such as in peak and average deceleration, peak vertical and posterior ground reaction forces, and knee flexion angles, while other outcomes such as knee valgus angles and moments, and ankle plantarflexion angle are scored negatively, and lower values would be deemed optimal (see Table 1). For this reason, each of the dependent variables was analysed separately.

### Data Analysis

Where possible, effect sizes and their 95% confidence intervals (CI) were calculated for all performance outcome measures of studies included in the review. Effect sizes were calculated between the intervention and control groups using the formula:

$$\frac{(M_1 - M_2)}{SD_{pooled}}$$

Where  $M_1$  and  $M_2$  are the post-test means for the intervention and control groups. To obtain the pooled standard deviation, the following formula was used:

$$\sqrt{\frac{(n_1 - 1) \times SD_1^2 + (n_2 - 1) \times SD_2^2}{(n_1 + n_2 - 2)}}$$

Where  $n_1$  and  $n_2$  are the sample sizes from the intervention and control groups respectively, and  $SD_1$  and  $SD_2$  are their respective standard deviations. Hedge's  $g$  correction factor was applied to all effect sizes to correct for small sample bias.

Post-intervention results were used due to the assumption that since these were RCTs there would be limited between-group differences prior to the intervention since they were randomised. When not



**Table 1.** Definition of all outcome variables assessed and their relevance to deceleration performance)

Outcome	Dependent Variable	Definition	Relevance to Deceleration
Ankle plantar / dorsi flexion	Joint kinematics	Joint angle of ankle during weight acceptance phase of final step	Lower values (more dorsiflexion) more beneficial for performance as ankle flexion contributes to lowering centre of mass to effectively orientate braking forces [19]
Average deceleration	Performance outcomes	Average of all instantaneous deceleration values ( $\text{m.s}^{-2}$ )	Direct measurement of ability to decelerate, higher numbers indicate greater ability to decelerate more rapidly and in shorter amount of time [20].
Hip abduction	Joint kinematics	Hip joint angle, relative to the trunk reference frame, with adduction-abduction	Lower values more beneficial for performance as too high values likely to be inefficient and increase risk of injury [18]
Hip adduction moment	Joint kinetics	Peak torque/rotational force about the hip joint resulting from forces acting to adduct the thigh	Lower values hypothesised to be more beneficial for performance [21] and reduce risk of injury
Hip flexion	Joint kinematics	Hip joint angle, relative to the trunk reference frame, with in flexion/extension	Higher values more beneficial for performance as hip flexion contributes to lowering centre of mass to effectively orientate braking forces [19]
Hip flexion moment	Joint kinetics	Peak torque/rotational force about the hip joint resulting from forces acting to flex the thigh	Higher values more beneficial for performance as this would be indicative of strong activation of hip flexor muscles [11]
Hip internal rotation moment valgus	Joint kinetics	Peak torque/rotational force about the hip joint resulting from forces acting to internally rotate and adduct the thigh	Higher values more beneficial for performance as lower values would be associated with a more stable and controlled deceleration rather than rapid, forceful braking [11]
Initial deceleration velocity	Performance outcomes	Running speed immediately prior to deceleration ( $\text{m.s}^{-1}$ )	Higher values indicate a better ability to decelerate more quickly from higher velocities, which indicates greater deceleration ability [13]
Knee abduction moment	Joint kinetics	Peak torque/rotational force about knee joint resulting from forces acting to abduct the knee	Higher values more beneficial for performance as lower values would be associated with a more stable and controlled deceleration rather than rapid, forceful braking [11]
Knee extension moment	Joint kinetics	Peak torque/rotational force about knee joint resulting from forces acting to extend the knee	Higher values more beneficial for performance as knee extensor muscles contract eccentrically to decelerate as fast as possible in a stable position [12, 22, 23]
Knee flexion at peak GRF	Joint kinematics	Angle of flexion about the knee joint taken at the time of peak impact GRF	Higher values more beneficial for deceleration performances as knee flexion contributes to lowering centre of mass to effectively orientate braking forces [19]
Knee flexion range of motion	Joint kinematics	Calculated as the difference between maximum knee flexion angle and the knee flexion angle at initial ground contact	Higher values more beneficial for deceleration performances as knee flexion contributes to lowering centre of mass to effectively orientate braking forces [19]
Knee rotation moment	Joint kinetics	Peak torque/rotational force about knee joint resulting from forces acting to externally rotate the knee	Higher values more beneficial for performance as lower values would be associated with a more stable and controlled deceleration rather than rapid, forceful braking. [24]

Outcome	Dependent Variable	Definition	Relevance to Deceleration
Knee valgus moment	Joint kinetics	Peak torque/rotational force about the knee joint that causes the lower leg to move inward towards the midline of the body	Lower values more beneficial to deceleration performance due to decreasing risk for ACL injury [25] and better alignment and control of the knee joint
Knee valgus varus	Joint kinematics	Valgus (positive values) refers to the angulation of the knee joint inward, where the proximal part of the lower leg deviates towards the body's midline relative to the thigh, and the distal part is more lateral. Varus (negative values) refers to angulation of the knee joint where the lower leg deviates outward away from the body's midline relative to the thigh.	Lower values (indicating less valgus) more beneficial to deceleration performance due to decreasing risk for ACL injury [38] and better alignment and control of the knee joint
Loading phase lateral hamstrings activation	Muscle activation	Mean amplitude of muscle activity of the lateral hamstrings assessed using surface EMG while the foot was in contact with the ground	Higher values more beneficial for performance as adequate activation is required during final foot contact while they contract eccentrically to absorb impact and control knee joint, and act to limit excessive anterior tibial translation which could contribute to ACL injury risk [26]
Loading phase medial hamstrings activation	Muscle activation	Mean amplitude of muscle activity of the medial hamstrings assessed using surface EMG while the foot was in contact with the ground	Higher values more beneficial for performance as adequate activation is required during final foot contact while they contract eccentrically to absorb impact and control knee joint, and act to limit excessive anterior tibial translation which could contribute to ACL injury risk [26]
Loading phase rectus femoris activation	Muscle activation	Mean amplitude of muscle activity of the rectus femoris assessed using surface EMG while the foot was in contact with the ground	Higher values more beneficial for performance as adequate activation is required during final foot contact while they contract eccentrically to absorb impact and control knee joint, and act to limit excessive anterior tibial translation which could contribute to ACL injury risk [26]
Loading phase vastus medialis activation	Muscle activation	Mean amplitude of muscle activity of the vastus medialis assessed using surface EMG while the foot was in contact with the ground	Higher values more beneficial for performance as quadriceps help to stabilise and control leg when decelerating from high speeds and control descent of the body during deceleration [22]
Peak deceleration	Performance outcomes	Peak value of instantaneous deceleration (m.s <sup>-2</sup> )	Direct measurement of ability to decelerate, higher numbers indicate greater ability to decelerate more rapidly and in shorter amount of time [20]
Peak knee Flexion	Joint kinematics	Maximum angle of flexion of the knee joint	Higher values more beneficial for performance as knee flexion contributes to lowering centre of mass to effectively orientate braking forces [27]
Peak posterior GRF	Kinetics	Maximum value of the ground-reaction forces exerted in the backward direction with foot contact with the ground collected during final foot contact using force plates and normalized to body weight.	Higher values more beneficial for performance as they indicate more force being exerted in the opposite direction of movement helping to slow down the athlete's momentum and enable more effective deceleration [7]
Peak vertical GRF	Kinetics	Maximum value of the ground reaction forces exerted vertically with foot contact with the ground collected during final foot contact using force plates and normalized to body weight.	Higher values more beneficial for performance indicate the athlete is using the ground effectively to brake their forward momentum indicating a more rapid and forceful deceleration [7].

Outcome	Dependent Variable	Definition	Relevance to Deceleration
Preparatory phase lateral hamstrings activation	Muscle activation	Mean amplitude of muscle activity of the lateral hamstrings assessed using surface EMG 50ms before final foot contact with the ground	Higher values more beneficial for performance as they act to resist forward momentum of the leg and control the descent of the limb prior to ground contact. Also act to limit excessive anterior tibial translation which could contribute to ACL injury risk [26]
Preparatory phase medial hamstrings activation	Muscle activation	Mean amplitude of muscle activity of the medial hamstrings assessed using surface EMG 50ms before final foot contact with the ground	Higher values more beneficial for performance as they act to resist forward momentum of the leg and control the descent of the limb prior to ground contact. Also act to limit excessive anterior tibial translation which could contribute to ACL injury risk [26]
Preparatory phase rectus femoris activation	Muscle activation	Mean amplitude of muscle activity of the rectus femoris assessed using surface EMG 50ms before final foot contact with the ground	Higher values more beneficial for performance as they act to resist forward momentum of the leg and control the descent of the limb prior to ground contact. Also act to limit excessive anterior tibial translation which could contribute to ACL injury risk [26]
Preparatory phase vastus medialis activation	Muscle activation	Mean amplitude of muscle activity of the vastus medialis assessed using surface EMG 50ms before final foot contact with the ground	Higher values more beneficial for performance as quadriceps work synergistically with hamstrings to prepare leg for ground contact [22]
Tibial anterior shear force	Joint kinetics	Maximum shear force exerted in the anterior direction to the proximal tibia pulling the tibia forward relative to the femur during ground contact	Higher values more beneficial for performance as they indicate a more rapid or forceful deceleration as athlete generates significant braking force through the tibia. Can pose risk of knee injury [25]
Tibial internal rotation moment	Joint kinetics	Peak torque/rotational force acting around the tibia where it inwardly rotates towards the body's midline	Higher values more beneficial for performance as lower values would be associated with a more stable and controlled deceleration rather than rapid, forceful braking [25]

reported, the SD was estimated from the standard error (SE) of the mean, 95% CI, or p-value as per the methods suggested in the Cochrane handbook [27]. SDs were estimated for one study [28] from the 95% CIs provided using the formula:

$$\sqrt{N} \times \frac{(\text{upper limit} - \text{lower limit})}{(2 \times 2.0639)}$$

Where N is the sample size of the group, and 2.0639 is the t-statistic for a 95% CI from a small sample size.

A meta-analysis was performed in R (version 4.3.0) using meta package and applying a multi-level random effects model, clustered according to individual studies, to estimate the pooled effect of training on deceleration ability. The random effects model was applied due to high level of variance expected between studies due to sampling errors and differences between studies in intervention protocols (intervention length, type, session duration, and frequency). The multi-level model was used to deal with the problem of dependencies arising from the

inclusion of multiple effect sizes per study which likely share contributors of variance and may have even more similarities because they belong to the same participants. Following Cochrane guidelines, for any study that included two intervention groups and one control group [29], the sample size in the control group was evenly divided so a comparison could be made to each intervention, and as the results are continuous outcomes, only the total number of participants was divided up and the means and standard deviations left unchanged [30]. Variance of the distribution of the true effect sizes ( $\tau^2$ ) was used to measure heterogeneity [31] and was calculated using the restricted maximum likelihood estimator. Prediction intervals were also calculated to aid in the interpretation of heterogeneity by estimating the likely effect that could be expected in future studies. Knapp-Hartung adjustments [32] were used to calculate the confidence intervals around the pooled effect. All analyses used two-sided tests, and  $p < 0.05$  considered statistically significant. Results are displayed as the mean difference between

pre- and post-intervention scores for both the intervention and control groups, with 95% confidence intervals.

### *Risk of Bias*

The revised Cochrane risk of bias tool for randomised trials (RoB2) [33] was used to assess characteristics including the randomisation process, blinding, deviations from intended interventions, availability of outcome data, appropriateness of measurement methods, and reporting of results and analyses. This tool was used because it is a specialised methodological instrument allowing for the systematic assessment of the risk of bias in randomised trials. One reviewer (LM) independently performed quality and risk of bias assessments on each of the included articles to assess and weight the quality of the included studies with respect to the research question, and accuracy was assessed by a second author (JF). Disagreements between reviewers' judgements were resolved through discussion or by additional input from a 3<sup>rd</sup> independent reviewer (BM).

## RESULTS

### *Study Selection*

The search strategy and results are shown in Figure 1. The literature search yielded 13,348 studies. The initial search was intentionally kept broad to not introduce systematic bias by excluding for example studies which measured deceleration outcomes, but did not focus on deceleration per se. Once duplicates were removed, the titles and abstracts of 8,797 articles were screened for inclusion. From these, 8,521 studies were excluded and the full texts from the remaining 276 were assessed for eligibility. The percentage agreement between the two screeners for title and abstract screening was 97% with kappa 0.61. This level of inter-rater reliability is moderate-to-substantial [34, 35], likely because many publications don't specify in the title or abstract whether any deceleration components were reported separately to other measures of agility or change of direction, and title and abstract were screened independently hence erred on the side of inclusion to introduce as little systematic bias as possible. Of the 276 full-text articles assessed, 264 were excluded (Online Resource 2). The remaining 12 met the inclusion criteria and were included in the analysis for this systematic review. The percentage agreement between the two reviewers for

studies included in the review was 94%, with kappa 0.53. Upon review and consultation with a third screener it was found that the moderate-to-substantial kappa score was due to the difficulty in discerning which results were true measures of deceleration performance. This difficulty stemmed from inconsistencies in terminology and methodology across studies, as well as variations in how deceleration was defined and measured. For instance, some studies referred to deceleration as part of a broader assessment of agility or COD, making it challenging to isolate and extract relevant data specific to deceleration. Additionally, the lack of standardised protocols for assessing deceleration further compounded the issue, with studies employing different testing procedures, equipment, and performance metrics. Consequently, there was ambiguity surrounding which outcomes truly reflected deceleration ability, leading to discrepancies in interpretation among reviewers. Of the included studies, 3 were unable to be used for the meta-analysis due to missing or incomplete data [15-17], leaving 9 studies for the meta-analysis.


### *Study Characteristics*

The descriptive results of the included studies are presented in Table 2. All were RCTs with pre- and post-intervention measures reported, including 380 participants with ages ranging from 15 to 34 years. The experience levels of included participants ranged from recreational ( $n = 124$ ) to professional ( $n = 23$ ), with most ( $n = 176$ ) reported to have some-to-moderate level of experience. The most common sport included was basketball ( $n = 4$ ) with 87 participants [16, 26, 36, 37], while other sports examined were soccer ( $n = 2$ ) with 44 participants [17, 29], AFL ( $n = 1$ ) with 34 participants [11], American football ( $n = 1$ ) with 21 participants [28], Gaelic football ( $n = 1$ ) with 31 participants [18], rugby ( $n = 1$ ) with 30 participants [38], and volleyball ( $n = 1$ ) with 9 participants [36]. Two studies with 124 participants did not report the specific sports [21, 24].

### *Methodological approaches*

One study [16] had a passive control group which only completed the testing procedures, while seven had control groups who performed their regular training [18, 21, 24, 26, 29, 36, 38]. The remaining control groups performed some form of standard training that is common in their sport such as traditional resistance exercise [17, 28, 38] or straight-line running and acceleration drills [15].





**Figure 1.** PRISMA flow diagram of systematic search and included studies

Strength or resistance training that would not be considered standard practice was the most common intervention ( $n = 6$ ) [21, 24, 29, 36, 37, 39]. Other interventions included technique instruction or feedback protocols ( $n = 3$ ) [15, 16, 21], balance or stability exercises ( $n = 3$ ) [15, 18, 26], plyometric training ( $n = 2$ ) [17, 36], agility training ( $n = 1$ ) [26], and small-sided games training ( $n = 1$ ) [29].

The duration of the interventions ranged from 2 - 28 weeks, and 10 - 46 sessions were performed. One study [16] was classified as short-term ( $< 2$  weeks), six studies were classified as intermediate-term (4 - 6 weeks) [17, 18, 26, 28, 29, 36], and five studies [15, 21, 24, 37, 38] were classified as long term ( $> 6$  weeks). Sessions were performed twice-weekly in five studies [15, 17, 29, 37, 38], thrice-weekly in five studies [18, 21, 24, 28, 36], four times per week in one study [25], and daily in one study [16]. One study [15] also had participants start with two sessions per week for 18 weeks, before condensing training to one session per week for the remaining ten weeks of the study.

### Outcome measures

Twenty-nine separate outcome measures specific to deceleration performance were identified in the included studies. There were five distinct themes in the outcome measures used by the included stud-

ies: joint kinematics, performance outcomes, joint kinetics, kinetics, and muscle activation patterns (Table 1). Joint kinematics were the most analysed, with seven studies [15-18, 21, 24, 25]. Performance outcomes [15, 17, 29, 37, 38], joint kinetics [15, 18, 21, 24, 28], and kinetics [18, 21, 24, 26, 36] were analysed in five studies each. While only one study included measures of muscle activation patterns [26].

### Effectiveness of interventions

#### Primary Outcome

Given the differences in the scales and variables used to measure deceleration performance, standardised mean differences were used to compare the between-groups differences post-intervention period. Figure 2 reports a summary of the meta-analyses performed for each deceleration-specific performance measure reported in this review. Across all observations of deceleration performance measures, when clustered by study, the implementation of a training intervention had on average a smaller effect ( $SMD = -0.04$ ,  $95\% CI = -0.50, 0.42$ ) than for control groups. The estimated variance of the true effect sizes ( $\tau^2$ ) was found to be  $<0.0001\%$ , and  $I^2 0\%$ , indicating no, or very low, heterogeneity but was not statistically significant. The prediction interval ranged from  $SMD = -0.52$  to  $0.43$ .

**Table 2.** Data extraction and descriptive results of included studies

Study	Sport	Participants	Sex	Population Characteristics	Intervention Group	Control Group	Testing Protocol	Deceleration-Specific Outcomes	Hedge's g Intervention vs. Control (95% CI)
Arede et al (2022)	Basketball	20	Males	All (mean $\pm$ SD) Age: 19.45 $\pm$ 4.36yrs (15-34) Height: 1.83 $\pm$ 0.09m Body mass: 86.36 $\pm$ 17.20kg Level: U/18s to amateur seniors	n = 10 Strength training 2 sessions/week for 10 weeks	n = 10 Training as usual	-V-cut test (25m sprint with 45° COD every 5m) - COD speed test - Game situation (5v5 full court simulation)	Deceleration properties: - Total decelerations (n.min <sup>-1</sup> ) - Total high intensity decelerations (n/min) - Peak deceleration (m.s <sup>-2</sup> )	- Peak deceleration: -0.53 (-0.75, -0.32)
Donnelly et al (2012)	AFL	34	Males	Intervention (mean $\pm$ SD) - Age 21.0 $\pm$ 3.3yrs - Height 1.86 $\pm$ 0.08m - Mass 81.2 $\pm$ 10.0kg  Control (mean $\pm$ SD) - Age 20.9 $\pm$ 3.1yrs - Height 1.84 $\pm$ 0.08m - Mass 80.6 $\pm$ 10.2kg	n = 14 Balance & technique training 2 sessions/week for 18 weeks, then 1 session/week for 10 weeks	n = 20 Acceleration & straight-line running 2 sessions/week for 18 weeks, then 1 session/week for 10 weeks	- Motion analysis of knee joint bio-mechanics during random series of pre-planned & unplanned straight running, cross-over cuts, and side-step running tasks	- Mean pre-contact velocity - Mean COD angle & velocity - Mean knee flexion & knee flexion ROM - Mean peak externally applied flexion, valgus, and internal rotation knee moments	Not Reported
Herman et al (2009)	Not reported	58	Females	Intervention (mean $\pm$ SD) - Height 1.67 $\pm$ 0.07m - Age 22.5 $\pm$ 2.3yrs - Mass 64.1 $\pm$ 9.1kg  Control (mean $\pm$ SD) - Height 1.66 $\pm$ 0.06m - Age 22.5 $\pm$ 3.8yrs - Mass 62.1 $\pm$ 7.3kg  Recreational	n = 29 Strength training + feedback protocol 3 sessions/week for 9 weeks	n = 29 Feedback protocol only (no strength training) 3 sessions/week for 9 weeks	Motion analysis of stop-jump task	Kinetic variables (sampled at peak anterior tibial shear force): - Peak proximal tibial anterior shear force - Vertical GRF - Knee valgus & extension moments - Hip adduction moment  Kinematic variables (sampled at peak anterior tibial shear force): - Knee & hip flexion angles - Knee valgus angle - Hip abduction angle	- Peak proximal tibial anterior shear force: -0.35 (-0.39, -0.31) - Vertical GRF: 0.12 (0.01, 0.23) - Knee flexion: 0.00 (-1.43, 1.43) - Knee valgus/varus: 0.16 (-0.74, 1.16) - Hip flexion: -0.05 (-2.42, 2.31) - Hip abduction: 0.35 (-1.93, 2.62) - Knee extension moment: -0.06 (-0.08, -0.05) - Knee valgus moment: 0.29 (0.28, 0.30) - Hip adduction moment: 0.44 (0.40, 0.47)


Study	Sport	Participants	Sex	Population Characteristics	Intervention Group	Control Group	Testing Protocol	Deceleration-Specific Outcomes	Hedge's g Intervention vs. Control (95% CI)
Herman et al (2008)	Not reported	66	Females	<p>Intervention (mean <math>\pm</math> SD)</p> <ul style="list-style-type: none"> <li>- Height 1.67 <math>\pm</math> 0.07m</li> <li>- Age 22.47 <math>\pm</math> 2.25yrs</li> <li>- Mass 63.52 <math>\pm</math> 9.18kg</li> </ul> <p>Control (mean <math>\pm</math> SD)</p> <ul style="list-style-type: none"> <li>- Height 1.66 <math>\pm</math> 0.06m</li> <li>- Age 22.53 <math>\pm</math> 3.81yrs</li> <li>- Mass 61.05 <math>\pm</math> 8.35kg</li> </ul> <p>Recreational</p>	n = 33 Strength training 3 sessions/week for 9 weeks	n = 33 Training as usual	Motion analysis of stop-jump task	<p>Kinetic variables (sampled at peak anterior tibial shear force):</p> <ul style="list-style-type: none"> <li>- Peak proximal tibial anterior shear force</li> <li>- Vertical GRF</li> <li>- Knee valgus &amp; extension moments</li> <li>- Hip adduction &amp; IR valgus moments</li> </ul> <p>Kinematic variables (sampled at peak anterior tibial shear force):</p> <ul style="list-style-type: none"> <li>- Knee &amp; hip flexion angles</li> <li>- Knee valgus angle</li> </ul>	<ul style="list-style-type: none"> <li>- Peak proximal tibial anterior shear force: 0.44 (0.40, 0.49)</li> <li>- Vertical GRF: 0.08 (-0.07, 0.23)</li> <li>- Knee flexion: 0.08 (-1.72, 1.88)</li> <li>- Knee valgus/varus: -0.27 (-1.15, 0.62)</li> <li>- Hip flexion: 0.01 (-2.84, 2.85)</li> <li>- Knee extension moment: 0.36 (0.35, 0.38)</li> <li>- Knee valgus moment: 0.24 (0.22, 0.25)</li> <li>- Hip adduction moment: 0.37 (0.36, 0.39)</li> <li>- Hip IR moment valgus: -0.14 (-0.15, -0.14)</li> </ul>
Jamison et al (2012)	American football	21	Males	<p>Trunk Stabilisation (mean <math>\pm</math> SD)</p> <ul style="list-style-type: none"> <li>- Height 1.82 <math>\pm</math> 0.06m</li> <li>- Age 20.5 <math>\pm</math> 1.2yrs</li> <li>- Mass 82.8 <math>\pm</math> 7.6kg</li> </ul> <p>Resistance Training (mean <math>\pm</math> SD)</p> <ul style="list-style-type: none"> <li>- Height 1.81 <math>\pm</math> 0.05m</li> <li>- Age 20.3 <math>\pm</math> 1.5yrs</li> <li>- Mass 82.9 <math>\pm</math> 5.4kg</li> </ul>	n = 10 Trunk stabilisation + strength training 3 sessions/week for 6 weeks	n = 11 Strength training only 3 sessions/week for 6 weeks	Motion analysis of biomechanical loading during 45° unanticipated run-to-cut manoeuvre	<p>Peak moments:</p> <ul style="list-style-type: none"> <li>- Knee abduction</li> <li>- Tibial IR</li> </ul>	<ul style="list-style-type: none"> <li>- Peak knee abduction moment: -1.86 (-2.27, -1.45)</li> <li>- Peak tibial IR moment: -1.31 (-1.40, -1.21)</li> </ul>
Kato et al (2006)	Basketball	10	Females	<p>University students All (mean <math>\pm</math> SD)</p> <ul style="list-style-type: none"> <li>- Height 1.62 <math>\pm</math> 0.03m</li> <li>- Mass 55.4 <math>\pm</math> 4.6kg</li> <li>- Age 20.4 <math>\pm</math> 0.5yrs</li> <li>- 11.7 <math>\pm</math> 1.3yrs of athletic hx</li> </ul>	n = 5 Technique instruction 7 sessions/week for 2 weeks	n = 5 Passive control group	Motion analysis of stop-motion task	<ul style="list-style-type: none"> <li>- Knee flexion angle</li> <li>- Angle on anterior surface</li> <li>- Torsion angle of two sides</li> <li>- 2-week compliance with exercise instructions</li> <li>- Shot success (success rate of 10 shots)</li> </ul>	Not Reported

Study	Sport	Participants	Sex	Population Characteristics	Intervention Group	Control Group	Testing Protocol	Deceleration-Specific Outcomes	Hedge's g Intervention vs. Control (95% CI)
Nevado-Garrosa et al (2021)	Soccer	23	Females	U23 female soccer players from the 2nd team of a professional Spanish club All (mean $\pm$ SD) - Age $19.3 \pm 1.7$ (16-23) - Mass $53.9 \pm 7.3$ kg - Height $1.61 \pm 0.06$ m	n = 8 Eccentric over-load training (EOT) 2 sessions/week for 5 weeks n = 8 Small-sided games training (SSG) 2 sessions/week for 5 weeks	n = 7 Training as usual	6v6+1 training task	Maximum decelerations (DCC max) The average of all the values was calculated as average DCC. Initial velocity deceleration (Initial V DCC), the running speed immediately before the start of deceleration	EOT vs. Control - Max. deceleration: 0.37 (0.18, 0.57) - Avg. deceleration: -0.48 (-0.56, -0.41) - Initial velocity of deceleration: 0.46 (0.09, 0.84) SSG vs. Control - Max. deceleration: -0.45 (-0.64, -0.26) - Avg. deceleration: -0.24 (-0.31, -0.17) - Initial velocity of deceleration: 0.30 (-0.05, 0.65)
Raedergard et al (2020)	Soccer	21	Males	Strength (mean $\pm$ SD) - Height $1.81 \pm 0.06$ m - Age $22.2 \pm 3$ yrs - Mass $77.1 \pm 7.2$ kg Plyometric (mean $\pm$ SD) - Height $1.82 \pm 0.06$ m - Age $22.6 \pm 2.6$ yrs - Mass $82.5 \pm 7.3$ kg	n = 11 Plyometric training 2 sessions/week for 6 weeks	n = 10 Strength training 2 sessions/week for 6 weeks	- Motion analysis of COD task; either 4m or 20m approach with cut angles of 45°, 90°, 135° & 180°	- COM displacement - Contact time - Hip flexion & abduction	Not Reported
Whyte et al (2018)	Gaelic football	31	Males	Intervention (mean $\pm$ SD) - Age $22.05 \pm 1.47$ years - Height $1.81 \pm 0.06$ m - Mass $78.5 \pm 8.35$ kg Control (mean $\pm$ SD) - Age $21.76 \pm 1.59$ years - Height $1.80 \pm 0.06$ m - Mass $79.13 \pm 10.24$ kg	n = 16 Core stability training 3 sessions/week for 6 weeks	n = 15 Training as usual	- Motion analysis of anticipated & un-anticipated cross-over & side-cutting manoeuvres	Sagittal, frontal, & transverse plane: - Kinematics of the trunk, hip, knee, and ankle - Internal moments of the hip, knee, and ankle - GRFs during first 30% of stance phase	Side-cutting - Knee valgus moment: -0.18 (-3.16, 2.81) - Hip flexion/extension moment: 1.12 (-5.11, 7.34) - Knee rotation moment: -0.43 (-1.24, 0.37) - Anterior/posterior GRF: -1.23 (-1.77, -0.70) & -0.92 (-1.33, -0.52) Crossover-cutting - Ankle plantar/dorsiflexion: 0.37 (-4.00, 4.73) - Anterior/posterior GRF: -0.81 (-1.25, -0.36); -1.42 (-2.24, -0.60); & -1.01 (-2.02, 0.01)



Study	Sport	Participants	Sex	Population Characteristics	Intervention Group	Control Group	Testing Protocol	Deceleration-Specific Outcomes	Hedge's g Intervention vs. Control (95% CI)
Wilderman et al (2009)	Basketball	30	Females	Intervention (mean $\pm$ SD) - Age $21.07 \pm 3.62$ years - Height $1.71 \pm 0.06$ m - Mass $67.58 \pm 7.71$ kg  Control (mean $\pm$ SD) - Age $21.07 \pm 1.83$ years - Height $1.71 \pm 0.04$ m - Mass $65.13 \pm 7.14$ kg	n = 15 Agility training 4 sessions/week for 6 weeks	n = 15 Training as usual	- Side-step pivot manoeuvre - 4m run then 45° in opposite direction to plant leg, wearing electromagnetic tracking sensors	Preparatory & loading phase activation (% MVIC) of: - Vastus Medialis Oblique - Rectus femoris - Medial hamstrings - Lateral hamstrings  Knee flexion (degrees) - At initial ground contact - Maximum - Displacement  Peak vertical GRF normalised to body weight	Loading phase activation (% MVIC) of: - Vastus Medialis oblique: 0.28 (-64.02 - 64.59) - Rectus femoris: -0.54 (-29.57, 28.48) - Medial hamstrings: 0.70 (-16.43, 17.82) - Lateral hamstrings: -0.02 (-23.44, 23.41)  Knee flexion (degrees) - At initial ground contact: 0.19 (-4.74, 5.12) - Maximum: 0.23 (-3.95, 4.42) - Displacement: 0.01 (-2.20, 2.23)  Peak vertical GRF: -0.75 (-0.91, -0.59)
Winwood et al (2015)	Rugby	30	Males	Strongman (mean $\pm$ SD) - Age $23.4 \pm 5.6$ years - Height $1.80 \pm 0.07$ m - Mass $91.2 \pm 14.8$ kg - $3.9 \pm 2.3$ years resistance training experience  Traditional (mean $\pm$ SD) - Age $22.5 \pm 3.4$ years - Height $1.81 \pm 0.06$ m - Mass $93.7 \pm 12.3$ kg - $4.7 \pm 3.3$ years resistance training experience	n = 15 Strongman training 2 sessions/week for 7 weeks	n = 15 Traditional strength training 2 sessions/week for 7 weeks	505 COD test	505 COD -5 to -2m time	505 COD deceleration: -0.05 (-0.06, -0.04)

Study	Sport	Participants	Sex	Population Characteristics	Intervention Group	Control Group	Testing Protocol	Deceleration-Specific Outcomes	Hedge's g Intervention vs. Control (95% CI)
Yang et al (2018)	Basketball Volleyball	36	Males Females	17 Male basketball players, 10 female basketball players, 1 male volleyball player, 8 female volleyball players	n = 18 Strength & plyometric training 3 sessions/week for 4 weeks	n = 18 Training as usual	- Motion analysis of stop-jump & side-cutting tasks	o Knee flexion angle at peak impact posterior GRF o Peak impact posterior and vertical GRF	Knee flexion angle at peak impact posterior GRF - Male, stop-jump: 0.88 (-2.95, 4.72) - Male, side-cutting: -0.86 (-3.99, 2.28) - Female, stop-jump: 0.18 (-3.49, 3.84) - Female, side-cutting: -0.61 (-4.64, 3.41)
				Intervention (mean ± SD) - Height 1.87 ± 0.09m (male), 1.78 ± 0.04m (female) - Mass 82.45 ± 16.54 kg (male), 70.21 ± 4.4kg (female)					Peak impact posterior GRF - Male, stop-jump: 0.10 (0.00, 0.20) - Male, side-cutting: 0.07 (-0.14, 0.28) - Female, stop-jump: 0.26 (0.11, 0.41) - Female, side-cutting: 0.55 (0.37, 0.73)
				Control (mean ± SD) - Height 1.89 ± 0.04m (male), 1.77 ± 0.07m (female) - Mass 79.04 ± 8.15kg (male), 65.96 ± 7.83kg (female)					Peak impact vertical GRF - Male, stop-jump: -0.38 (-0.63, -0.13) - Male, side-cutting: 0.18 (-0.24, 0.60) - Female, stop-jump: 0.25 (-0.10, 0.61) - Female, side-cutting: 0.22 (-0.20, 0.63)



**Figure 2.** Forest plot displaying the standardised mean differences (SMD) and 95% confidence intervals (CIs) for all observations of all deceleration-specific performance measures. Negative values favour control groups, positive values favour interventions. Size of the box refers to the weight of the study in relation to the pooled estimate.

### Secondary Outcomes

An analysis of the secondary outcomes related to deceleration performance, clustered according to study, can be seen in figures 3 through 6, and all effect sizes calculated can be found in Online Resources 3 and 4.

Training interventions resulted in a slightly greater, but non-significant effect for measures of joint kinematics (SMD = 0.07, 95% CI = -0.11, 0.24,  $P = 0\%$ ,  $p = 0.77$ ). Whereas for kinetics (SMD = -0.29, 95% CI = -0.83, 0.25,  $P = 55\%$ ,  $p < 0.01$ ), joint kinetics (SMD = -0.20, 95% CI = -1.01, 0.61,  $P = 68\%$ ,  $p < 0.01$ ) and muscle activation (SMD = -0.10, 95% CI = -0.44, 0.24,  $P = 19\%$ ,  $p = 0.28$ ) the effect for training interventions was less than for control groups.


### Quality assessment

A summary of the quality assessment results is available in Figure 7, with detailed results made available in Online Resource 5. A high overall risk of bias was observed in all studies included in this review. With respect to the randomisation sequence, one study was subject to high risk of bias due to lack of random sequence generation and lack of allocation concealment [29]. One study was con-

sidered low risk [18], while the rest presented some concerns due to a lack of information regarding the randomisation process. Nine studies were high risk for deviations from intended interventions due to a low compliance rate with the intervention or an inappropriate analysis used to estimate the effect of assignment to intervention and potential for this to substantially bias the studies' findings [15-18, 28, 29, 36-38]. One study was considered low risk of bias [26] while the remaining two had some concerns [21, 24]. Three of the studies were observed to be low risk for having all or nearly all available outcome data [16, 18, 38], while nine were high risk for missing outcome data due to high dropouts or a lack of information [15, 17, 21, 23, 24, 26, 28, 29, 36, 37]. All studies presented a low risk of bias for measurement of the outcome as all measurements were objective and all participants performed the same assessments and at the same time periods in all included studies. Finally, all studies presented some concerns when addressing bias in selection of the reported result as it was not reported if the data were analysed against a pre-specified plan.


### DISCUSSION

The purpose of this systematic review was to crit-




**Figure 3.** Forest plot displaying the standardised mean differences (SMD) and 95% confidence intervals (CIs) for joint kinetics outcome variables. Negative values favour control groups, positive values favour interventions. Size of the box refers to the weight of the study in relation to the pooled estimate.






**Figure 4.** Forest plot displaying the standardised mean differences (SMD) and 95% confidence intervals (CIs) for joint kinematics outcome variables. Negative values favour control groups, positive values favour interventions. Size of the box refers to the weight of the study in relation to the pooled estimate. ROM: range of motion; GRF: ground reaction force



**Figure 5.** Forest plot displaying the standardised mean differences (SMD) and 95% confidence intervals (CIs) for kinetics outcome variables. Negative values favour control groups, positive values favour interventions. Size of the box refers to the weight of the study in relation to the pooled estimate. GRF: ground reaction force.




**Figure 6.** Forest plot displaying the standardised mean differences (SMD) and 95% confidence intervals (CIs) for muscle activation outcome variables. Negative values favour control groups, positive values favour interventions. Size of the box refers to the weight of the study in relation to the pooled estimate.

ically evaluate and summarise the results of studies examining the effectiveness of different training interventions to improve measures of deceleration performance in adult team-based field and court sports athletes, compared to control groups. By assessing the effectiveness of training interventions utilised (i.e. type of training, timing, frequency, and duration) against performing no training, sport-typical training, or continuing “as usual training”, this review also aimed to guide sporting practitioners who wish to improve their athletes’ deceleration performance through evidence-based practice. All RCTs that measured deceleration performance, kinetics, or kinematics during the final deceleration steps to reduce velocity prior to coming to a complete stop or the final foot contact of a severe COD which required coming to zero velocity were included. The main finding from our meta-analysis was that across the reported deceleration-specific outcomes there was an overall small effect which favoured control groups over the implementation of training interventions.

In analysing the primary outcomes related to deceleration performance (Figure 2), we showed no, or very low, heterogeneity with a  $\tau^2$  value of  $< 0.0001$  suggesting the studies in the meta-analyses for these primary outcomes were relatively homogeneous in their effect sizes. This is supported by the  $I^2$  value of 0% indicating that any variability in the observed effects is likely due chance or random sampling variation. Irrespective of the study designs used (all RCTs), there was large variability in the type, timing, frequency, and duration of the intervention protocols. Different forms of training were

used, such as strength, balance, plyometric, agility, and small-sided games training. The duration of these interventions also ranged widely, from as short as 2 weeks to as long as 28 weeks, with varying session frequencies. Furthermore, there was a lack of consistency in reporting of the results, with 29 separate deceleration-specific outcomes reported across the included studies. This meant there were few observations for each outcome variable, most likely influencing the interpretation of the results of our meta-analysis. For the primary outcomes of deceleration performance the effect for intervention versus control groups was inconclusive, with an SMD = -0.04 (95% CI -0.50, 0.42). While this was surprising, it may be explained by the fact that none of the training interventions were deceleration-specific, targeting stopping, changes of direction or deceleration mechanics. Instead, the interventions included strength training [37, 38], eccentric overload training [29], and small-sided games training [28] in which some outcome measures could be related to deceleration performance. Additionally, there were few observations of average and peak deceleration performance to contribute to the meta-analysis. The prediction interval ranged from SMD = -0.52 to 0.43, indicating the possibility for some future studies to find a larger effect for control groups based on present evidence. However, the interval is quite broad, indicating our findings are unable to elucidate the true effect and further research is required [39].

As a secondary outcome of deceleration performance, kinetic measures (Figure 5) were used in some studies [18, 21, 24, 26, 36]. The ability to de-



**Figure 7.** Summary of risk of bias assessment results

celerate effectively requires the application of rapid braking forces to reduce forward momentum [40]. Peak posterior ground reaction force refers to the maximum force exerted by an athlete against the ground in the backward direction during activities such as stopping, changing direction, or slowing down. Dos'Santos et al. demonstrated a substantial association between horizontal braking forces (HBF) and faster 505 COD times [7]. Athletes who exhibited higher HBF during specific phases, particularly in the penultimate step and the final ground contact, demonstrated superior performance in COD tasks. This underscores the significance of effective braking forces, particularly in the horizontal plane, for deceleration performance. Here, a negative SMD of -0.29 was observed, showing a smaller effect in the intervention group compared to the control group. However this difference was not statistically significant, indicating that training interventions did not produce a meaningful difference in deceleration performance compared to control groups. Peak posterior GRF was lower following the training interventions, however the wide confidence intervals prevent a definitive conclusion about the true effect size. Peak vertical GRF remained relatively unchanged, indicating little difference between intervention & control groups. Greater braking forces would be expected after successful training interventions to enable the execution of decelerations in shorter time frames and distances [11] provided the intervention was applied effectively and cued appropriately to ensure training transfer such as Dos'Santos et al. who provided external verbal coaching cues such as "slam on the brakes" and "push/punch the ground away" to promote rapid braking and faster COD performance [41]. Since the results of this meta-analysis show that training interventions had no effect on increasing braking forces when compared to control groups, an analysis of the content of the intervention such as exercise selection, intensity, volume, and the integration of sport-specific cues, could shed light on the factors influencing the observed outcomes.

The remaining secondary outcomes (joint kinetics and kinematics, and muscle activation patterns) were indirect measures of deceleration performance relating to coordination patterns employed by the performer during the final steps to reduce velocity prior to coming to a complete stop or the final foot contact of a severe COD which required coming to momentary zero velocity. Some of the technical movement characteristics are associated with reducing injury risk (e.g. knee valgus and tibial internal rotation) while others are related to effective decel-

eration performance (e.g. knee and hip flexion) [37, 42, 43]. Across the measures of joint kinetics (Figure 3), there was an overall small but non-significant effect for the control groups compared to training interventions. However there was considerable heterogeneity indicative of significant variability among the studies' results. Knee abduction moment, tibial internal rotation moment, knee rotation moment, and hip internal rotation moment valgus were lower following training interventions than in control groups which would be conducive to a reduced risk of injury. The relationship between moments and deceleration performance is complex, and it is important to consider that sometimes safe movement mechanics and performance can be in conflict [42]. Larger moments, indicative of a more forceful deceleration, may not always align with improved performance. The effectiveness of deceleration can depend on the specific strategy employed. For instance, a wide lateral foot plant strategically employed to evade opponents can increase knee abduction moment, but athletes may employ strategies to mitigate this and reduce this moment without compromising performance [7, 8]. Deceleration performance involves various factors and strategies and should be considered in light of the specific demands and objectives within the sporting context. Hip flexion moment was significantly greater for the intervention groups which would also favour better deceleration performance as the strong hip flexor muscles would be activated to rapidly flex the thigh allowing for quick steps to slow down [11, 18]. High angular velocities produced by these moments acting about the joints are typically observed during decelerations where rapid triple flexion of the hip, knee, and ankle are required to effectively orientate braking forces [19, 45]. Hip adduction moment and knee valgus moments saw greater effects following interventions than for controls, however attributing a straightforward interpretation of these values may oversimplify the complex nature of deceleration strategies especially in the sporting context. While lower values for these kinetic measures would be considered more optimal for reducing high-risk technique deficits [19, 21, 24], it is important to recognise the context-dependent nature of these mechanics. In agility and COD, the movement strategy will emerge within the constraints imposed by the task. For example, a wider lateral foot plant during deceleration is a strategy athletes frequently employ when evading opponents and executing more pronounced changes of direction (greater than 70°) and is associated with an increase in valgus and knee abduction moments. This highlights the limitations in just measuring deceleration or COD mechanics without consid-



ering the task and environmental constraints under which they occur. A comprehensive understanding of deceleration mechanics requires a consideration of these contextual factors.

There was no substantial effect of training on measures of joint kinematics (Figure 4) compared to control groups. Both hip abduction and ankle plantarflexion were slightly higher for the training intervention groups, although this was non-significant. These would be detrimental to deceleration performance due to their associations with injury risk through increasing knee varus loading and potential ACL strain [18]. Additionally, lower ankle plantarflexion angles are typically observed in better decelerators [19]. However, the wide confidence intervals about these results indicate a degree of uncertainty and caution should be taken when drawing conclusions. For hip flexion and measures of knee flexion, greater values, within a specific bandwidth, would indicate better deceleration performance as they are necessary to lower the athlete's centre of mass to increase stability and place the athlete in a better position to produce more horizontally orientated braking force [19]. However it should be noted that there is an optimal range for these angles, as excessive knee flexion may be associated with a more compliant strategy, potentially hindering deceleration performance. Effective deceleration into COD requires a balance between knee flexion, centre of mass, and leg stiffness. Differences between intervention and control groups were inconclusive.

Regarding muscle activation patterns (Figure 6) there was also little difference between intervention and control groups. Muscle activation, as measured through surface EMG, refers to the electrical activity generated within skeletal muscles during contraction. Unlike muscle contraction, which generally implies the shortening of muscle fibres resulting in movement, muscle activation encompasses the entire process of neuromuscular activity, including the initiation and transmission of electrical signals from the nervous system to the muscle fibres [45]. During running deceleration, a high degree of muscle activation is required to attenuate the large external moments generated during forceful ground contact and allow for the required external braking impulse to be generated to reduce horizontal momentum [22, 44]. The results of this meta-analysis showed that hamstring and quadricep activation during the preparatory phase favoured control groups, potentially indicating a detriment to deceleration performance. Within the literature it is typically suggested that plyometrics can increase muscle pre-activation

[46-48]. The findings from this meta-analysis indicating little difference in preparatory phase muscle activation patterns prompts consideration of the specificity and representativeness of the employed training interventions. Training for improved deceleration performance should aim to replicate the demands of these movements, such as those encountered during dynamic, sport-specific scenarios. Training specificity and representativeness may be essential to effectively enhance appropriate muscle pre-activation and consequently improve deceleration performance. On the other hand, loading phase medial hamstring and quadriceps activation were higher for the intervention groups suggesting benefits to deceleration as the hamstring act eccentrically to absorb impact and control the knee joint during high knee flexion angular velocities [22], while the quadriceps stabilise and control the leg and descent of the body during deceleration. However, wide confidence intervals were observed for all observations of muscle activation patterns which indicates substantial variability in the reported data and a need for caution when interpreting these findings.

There was an overall high risk of bias observed in all studies included in this review (Figure 7) suggesting caution should be applied when interpreting the findings from our review. The majority of these biases were due to lack of reporting in the randomisation process, low compliance rate with the intervention or an inappropriate analysis used to estimate the effect of assignment to intervention, and selection of the reported result. Compliance rates were not described by all studies, and where explanations were provided for the low compliance rates, it was attributed to scheduling conflicts, transport issues, or lower extremity injuries. Notably, only one study was at low risk of bias regarding the randomisation process, with the remaining raising concerns due to a lack of transparency in reporting of the process. Concerns regarding deviations from intended interventions primarily arose from low compliance rates and inappropriate analyses used to estimate the intervention effects. These deviations, present in nine studies, potentially introduced substantial biases. Additionally, issues related to outcome data were common, with nine studies exhibiting high risk due to significant dropouts or insufficient data documentation which raises questions about the completeness of the reported results, impacting the overall confidence in the findings. All studies displayed a low risk of bias in outcome measurement, as objective assessments were used across participants and time periods. However, there were concerns in addressing bias related to the selection of report-

ed results, as the analysis against a pre-specified plan was not consistently reported. Future research investigating the effectiveness of training interventions for improving deceleration performance should be done with more rigorous methodological designs and more complete reporting within the manuscript to address these areas.

## CONCLUSION

The findings of this systematic review and meta-analysis highlight several critical areas that warrant further research including the trainability of deceleration in team-based field and court sports athletes. While there is a plethora of literature investigating the trainability of linear speed, acceleration, COD, and agility in team-based field and court sports athletes [9, 10], deceleration as an explicit skill has not been so widely researched [4]. As such, there are few high quality RCTs specifically examining the effectiveness of training interventions for specifically enhancing deceleration performance. Future research should also include more rigorous methodological reporting to address areas that may introduce potential biases. This could involve providing more detail regarding randomisation processes, ensuring intervention protocols are adequately described, and accurately reporting participant compliance.

Since movement outcomes are a product of both skill and physical capacity, the lack of specificity and representativeness of the training interventions employed may compromise training transfer. More integration is needed of more detailed and sport-specific training protocols that closely replicate the dynamic and context-specific demands of deceleration performed in sporting scenarios. Robust training designs tailored explicitly to deceleration would contribute to a greater understanding of effective training strategies. Implementing detailed guidelines for intervention design, clearly outlining the components and structure of training programs, would contribute to increased methodological consistency across studies and improved comparability. Standardised protocols for interventions, including consistent type, timing, frequency, and duration, of training are recommended. Additionally, collaboration among researchers in the field could lead to the development of consensus guidelines for deceleration intervention protocols to promote a more consistent and comprehensive approach to studying deceleration performance.

Furthermore, future research should prioritise standardisation of the reported outcomes specifically related to deceleration ability. Crucial to this process would be achieving consensus among researchers regarding the most appropriate objective measures of deceleration ability that are reliable, valid, and precise and establishing gold standard protocols for assessing deceleration performance. This would provide a foundation for meaningful comparisons and facilitate comprehensive evaluations of deceleration performance across different interventions and athlete populations.

The purpose of this systematic review and meta-analysis was to critically examine the current evidence to determine effectiveness of training interventions aimed at improving deceleration performance in team-based field and court sports athletes. It is pertinent to acknowledge that very few studies have specifically targeted this construct and measured relevant outcomes directly related to deceleration ability. This scarcity, coupled with a high risk of bias reported across all studies, leads to an overall uncertainty regarding the effectiveness of training interventions for improving deceleration performance. The findings from this review revealed that, for the deceleration-specific outcomes reported in the included studies, training interventions did not demonstrate a significant performance improvement when compared to control groups. Despite the results of this systematic review and meta-analysis not supporting the effectiveness of training interventions for improving deceleration performance, these null/negative findings should be interpreted within the context of the study limitations. The limited number of included studies, inconsistencies in experimental designs and reported outcomes, and the observed high risk of bias demonstrate that more research and greater methodological transparency are needed to provide a comprehensive understanding of training to improve deceleration performance.

## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable

## CONSENT FOR PUBLICATION

Not applicable

## AVAILABILITY OF DATA AND MATERIAL

The datasets generated and/or analysed during the current study are available in the Open Science Framework repository. [https://osf.io/sg4qu/?view\\_only=c958a5ef838d4d2ebedfc718bc5c37ba](https://osf.io/sg4qu/?view_only=c958a5ef838d4d2ebedfc718bc5c37ba)

## COMPETING INTERESTS

Louise A. Marvin, Job Fransen, Paul Read, Blake McLean, and Sam Palmer declare that they have no conflicts of interest relevant to the content of this review.

## FUNDING

Not applicable

## AUTHORS' CONTRIBUTIONS

LM, JF, BM, & PR conceived the idea and design of the systematic review and meta-analysis and contributed to the development of the review protocol and analysis plan. LM ran the literature searches. LM and SP screened titles, abstracts, and full texts for inclusion, while JF acted as an additional independent screener to resolve disagreements. LM extracted and collated relevant data from the selected studies, conducted statistical analyses, and interpreted the results. JF contributed to the quality and risk of bias assessment. JF and BM provided guidance around the statistical analyses and interpretation of results. LM drafted the initial manuscript. JF, BM, and PR provided critical feedback and revisions to the manuscript. All authors read and approved the final manuscript.

## ACKNOWLEDGEMENTS

Not applicable

## AUTHORS' INFORMATION

Not applicable

## LIST OF ABBREVIATIONS

ACL: Anterior cruciate ligament  
COD: Change of direction

COM: Centre of mass  
CI: Confidence interval  
EMG: Electromyography  
EOT: Eccentric overload training  
GRF: Ground reaction force  
HBF: Horizontal braking force  
IR: Internal rotation  
MVIC: Maximal voluntary isometric contraction  
PICOS: Population, intervention, comparator, outcome, and study design  
PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses  
RCT: Randomised controlled trial  
RoB2: Cochrane risk-of-bias tool for randomised trials  
ROM: Range of motion  
SD: standard deviation  
SMD: standardised mean difference  
SSG: Small-sided games  
VGRF: Vertical ground reaction force

## REFERENCES

1. Hewitt J, Cronin J, Button C, & Hume P. Understanding deceleration in sport. *Strength Cond J*. 2011. 33(11): 47-52.
2. Lockie RG, Schultz AB, Callaghan SJ, Jeffries MD. The effects of traditional and enforced stopping speed and agility training on multidirectional speed and athletic function. *J. Strength Cond. Res*. 2014. 28(6): 1538-1551. <https://doi.org/10.1519/JSC.0000000000000309>
3. Harper DJ, Carling C, Kiely J. High-intensity acceleration and deceleration demands in elite team sports competitive match play: A systematic review and meta-analysis of observational studies. *Sports Med*. 2019. 49. <https://doi.org/10.1007/s40279-019-01170-1>
4. Harper DJ, Kiely J. Damaging nature of accelerations: Do we adequately prepare players? *BMJ Open Sport & Exercise Medicine*. 2018, 4(e000379). <https://doi.org/10.1136/bmjsem-2018-000379>
5. Ramírez-Campillo R, Vergara-Pedrerós M, Henríquez-Olguín C, Martínez-Salazar C, Alvarez C, Nakamura FY, et al. Effects of plyometric training on maximal-intensity exercise and endurance in male and female soccer players. *J Sport Sci*. 2016;34(8):687–93. <https://doi.org/10.1080/02640414.2015.1068439>
6. Spinks CD, Murphy AJ, Spinks WL, Lockie RG. The effects of resisted sprint training on acceleration performance and kinematics in soccer, rugby union, and Australian football players. *J Strength Cond. Res*. 2007;21(1):77–85. <https://doi.org/10.1519/00124278-200702000-00015>
7. Dos'Santos T, McBurnie A, Comfort P, Jones PA. The effects of six-weeks change of direction speed and

- technique modification training on cutting performance and movement quality in male youth soccer players. *Sports* 2019;7:205. <https://doi.org/10.3390/sports7090205>
8. Dos'Santos T, Thomas C, McBurnie A, Comfort P, Jones PA. Change of Direction Speed and Technique Modification Training Improves 180° Turning Performance, Kinetics, and Kinematics. *Sports*. 2021;9(6):73. <https://doi.org/10.3390/sports9060073>
9. Brughelli M, Cronin J, Levin G, Chaouachi A. Understanding change of direction ability in sport. A review of resistance training studies. *Sports Med*. 2008;38(12): 1045-1063.
10. Young WB, Farrow D, Pyne D, McGregor W, Handke T. Validity and reliability of agility tests in junior Australian Football players. *J Strength Cond. Res*. 2011; 25(12): 3399-3403. <https://doi.org/10.1519/JSC.0B013E318215FA1C>
11. Harper DJ, McBurnie AJ, Dos'Santos T, Eriksrud O, Evans M, Cohen DD., Rhodes D, Carling C, Kiely J. Biomechanical and neuromuscular performance requirements of horizontal deceleration: A review with implications for random intermittent multi-directional sports. *Sports Med*. 2022;52(10): 2321-2354. <https://doi.org/10.1007/s40279-022-01693-0>
12. Gilchrist HL. Does the implementation of rapid deceleration training improve change of direction performance in rugby players? [Master's Thesis, Auckland University of Technology]. 2018. <http://hdl.handle.net/10292/12159>
13. Graham-Smith P, Rumpf M, Jones P. (2018, September 10-14). Assessment of Deceleration Ability and Relationship to Approach Speed and Eccentric Strength. 36th Conference of the International Society of Biomechanics in Sports, Auckland, New Zealand.
14. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021; 372 :n71. <https://doi.org/10.1136/bmj.n71>
15. Donnelly CJ, Elliott BC, Doyle TLA, et al. Changes in knee joint biomechanics following balance and technique training and a season of Australian football. *Br. J. Sports Med*. 2012;46:917-922. <https://doi.org/10.1136/bjsports-2011-090829>
16. Kato S, Urabe Y, Kawamura K. The effect of dynamic alignment control training in movement pattern during the stop action. *Japanese Journal of Clinical Sports Medicine*. 2006;14(1): 13-19.
17. Raedergard HG, Falch NF, & van den Tillaar R. Effects of strength vs. plyometric training on change of direction performance in experienced soccer players. *Sports*, 2020;8(11), 144. <https://doi.org/10.3390/sports8110144>
18. Whyte EF, Richter C, O'Connor S, Moran KA. Effects of a dynamic core stability program on the biomechanics of cutting maneuvers: A randomized controlled trial. *Scand. J. Med. Sci*. 2018;28(2), 452-462. <https://doi.org/10.1111/sms.12931>
19. Dos'Santos T, McBurnie A, Thomas C, Comfort P, Jones PA. Biomechanical determinants of the modified and traditional 505 change of direction speed test. *J Strength Cond Res*. 2020;34:1285-96. <https://doi.org/10.1519/JSC.0000000000003439>
20. Harper DJ, Morin J-B, Carling C, Kiely J. Measuring maximal horizontal deceleration ability using radar technology: Reliability and sensitivity of kinematic and kinetic variables. *Sports Biomech*. 2020;1-17. <https://doi.org/10.1080/14763141.2020.1792968>
21. Herman DC, Onate JA, Weinhold PS, Guskiewicz KM, Garrett WE, & Padua DA. The effects of feedback with and without strength training on lower extremity biomechanics. *Am. J. Sports Med*. 2009;37(7):1301-1308. <https://doi.org/10.1177/0363546509332253>
22. Colby S, Francisco A, Yu B, Kirkendall D, Finch M, Garrett W. Electromyographic and kinematic analysis of cutting maneuvers. Implications for anterior cruciate ligament injury. *Am. J. Sports Med*. 2000;28:234-40
23. Zhang Q, Aurelie L, Foure A, Wong DP, Hauter CA. Relationship between explosive strength capacity of the knee muscles and deceleration performance in female professional soccer players. *Front Physiol*. 2021;12:723041. <https://doi.org/10.3389/fphys.2021.723041>
24. Herman DC, Weinhold PS, Guskiewicz KM, Garrett WE, Padua DA. The effects of strength training on the lower extremity biomechanics of female recreational athletes during a stop-jump task. *Am. J. Sports Med*. 2008;36(4):733-740. <https://doi.org/10.1177/0363546507311602>
25. Hewett TE, Myer GD, Ford KR, Heidt RS, Colosimo AJ, McLean SG, van den Bogert AJ, Paterno MV, Succop P. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med*. 2005; 33: 492-501. <https://doi.org/10.1177/0363546504269591>
26. Wilderman DR, Ross SE, Padua DA. Thigh muscle activity, knee motion, and impact force during side-step pivoting in agility-trained female basketball players. *Journal of Athletic Training*, 2009;44(1):14-25. <https://doi.org/10.4085/1062-6050-44.1.14>
27. Higgins JPT, Li T, Deeks JJ (editors). Chapter 6: Choosing effect measures and computing estimates of effect. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.3 (updated February 2022). Cochrane, 2022. Available from [www.training.cochrane.org/handbook](http://www.training.cochrane.org/handbook).
28. Jamison ST, McNeilan RJ, Young GS, Givens DL, Best TM, Chaudhari AMW. Randomized controlled trial of the effects of a trunk stabilization program on trunk control and knee loading. *Med. Sci. Sports Exerc*. 2012;44(10): 1924-1934. <https://doi.org/10.1249/MSS.0b013e31825a2f61>
29. Nevado-Garrosa F, Torreblanca-Martinez V, Paredes-Hernandez V, del Campo-Vecino J, Balsalo-



- bre-Fernandez C. Effects of an eccentric overload and small-side games training in match accelerations and decelerations performance in female under-23 soccer players. *J. Phys. Educ. Sport.* 2021;21, 3244-3251. <https://doi.org/10.7752/jpes.2021.s6431>
30. Higgins JPT, Eldridge S, Li T (editors). Chapter 23: Including variants on randomized trials. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.4 (updated August 2023). Cochrane, 2023. Available from [www.training.cochrane.org/handbook](http://www.training.cochrane.org/handbook)
  31. Borg DN, O'Brien JL. Comment on "Effects of heat acclimation and acclimatisation on maximal aerobic capacity compared to exercise alone in both thermo-neutral and hot environments: A meta-analysis and meta-regression". *Sports Med.* 2022;52, 1715-1718. <https://doi.org/10.1007/s40279-021-01611-w>
  32. Knapp G, Hartung J. Improved tests for a random effect meta-regression with a single covariate. *Statistics in Medicine*, 2003; 22(17): 2693-2710. <https://doi.org/10.1002/sim.1482>
  33. Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ.* 2019; 366 :l4898 <https://doi.org/10.1136/bmj.l4898>
  34. Kottner J, Audige L, Brorson S, Donner A, Gajewski BJ, Hrobjartsson A, Roberts C, Shoukri M, Streiner DL. Guidelines for reporting reliability and agreement studies (GRRAS) were proposed. *Journal of Clinical Epidemiology.* 2011; 64(1): 96-106. <https://doi.org/10.1016/j.jclinepi.2010.03.002>
  35. McHugh ML. Interrater reliability: the kappa statistic. *Biochemica Medica.* 2012; 22(3), 276-282. <https://doi.org/10.11613/bm.2012.031>
  36. Yang C, Yao W, Garrett WE, Givens DL, Hacke J, Liu H, Yu B. Effects of an intervention program on lower extremity biomechanics in stop-jump and side-cutting tasks. *Am. J. Sports Med.* 2018;46(12):3014-3022. <https://doi.org/10.1177/0363546518793393>
  37. Arede J, Leite N, Tous-Farjado J, Bishop C, Gonzalo-Skok O. (2022) Enhancing high-intensity actions during a basketball game after a strength training program with random recovery times between sets. *J. Strength Cond. Res.* 2022;36(7):1989-1997. <https://doi.org/10.1519/JSC.0000000000004002>
  38. Winwood PW, Cronin JB., Posthumus LR, Finlayson SJ, Gill ND, Keogh JWL. Strongman vs. traditional resistance training effects on muscular function and performance. *J. Strength Cond. Res.* 2015;29(2):429-439. <https://doi.org/10.1519/JSC.0000000000000629>
  39. Schünemann HJ, Vist GE, Higgins JPT, Santesso N, Deeks JJ, Glasziou P, Akl EA, Guyatt GH. Chapter 15: Interpreting results and drawing conclusions. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.4 (updated August 2023). Cochrane, 2023. Available from [www.training.cochrane.org/handbook](http://www.training.cochrane.org/handbook)
  40. Thieschäfer L, Klütz J, Weig J, Dos'Santos T, Büsch D. Development of a cutting technique modification training program and evaluation of its effects on movement quality and cutting performance in male adolescent American football players. *Sports* 2023;11:184. <https://doi.org/10.3390/sports11090184>
  41. Dos'Santos T, Thomas C, Jones PA, Comfort P. Mechanical determinants of faster change of direction speed performance in male athletes. *J Strength Cond Res.* 2017;31(3):696–705. <https://doi.org/10.1519/JSC.0000000000001535>
  42. Fox AS. Change-of-direction biomechanics: Is what's best for anterior cruciate ligament injury prevention also best for performance? *Sports Med.* 2018;48:1799-1807. <https://doi.org/10.1007/s40279-018-0931-3>
  43. Dos'Santos T, Thomas C, Comfort P, Jones PA. Role of the penultimate foot contact during change of direction: Implications on performance and risk of injury. *Strength Cond. J.* 2019, 41: 87-104. <https://doi.org/10.1519/SSC.0000000000000395>
  44. McBurnie AJ, Harper DJ, Jones PA, Dos'Santos T. Deceleration training in team sports: Another potential 'vaccine' for sports-related injury? *Sports Med.* 2021;52:1-12. <https://doi.org/10.1007/s40279-021-01583-x>
  45. Türker H, Sözen H. Surface electromyography in sports and exercise. In: Türker H, editor. *Electrodiagnosis in New Frontiers of Clinical Research.* IntechOpen; 2013. p. 181. <https://doi.org/10.5772/56167>
  46. Behrens M, Mau-Moeller A, Mueller K, Heise S, Gube M, Beuster N, et al. Plyometric training improves voluntary activation and strength during isometric, concentric and eccentric contractions. *J Sci Med Sport.* 2016;19(2):170–6. <https://doi.org/10.1016/j.jsams.2015.01.011>
  47. Behrens M, Mau-Moeller A, Bruhn S. Effect of plyometric training on neural and mechanical properties of the knee extensor muscles. *Int J Sports Med.* 2014;35(2):101–19. <https://doi.org/10.1055/s-0033-1343401>
  48. Chimera NJ, Swanik KA, Swanik CB, Straub SJ. Effects of plyometric training on muscle-activation strategies and performance in female athletes. *J Athl Train.* 2004;39(1):24-31.